

## Original Research Article

### **Effect of budding height on success of patch budding in Aonla (*Emblica officinalis* Gaertn.)**

#### **Abstract**

The present investigation entitled “Studies on effect of budding height on patch budding in Aonla (*Emblica officinalis* Gaertn.)” was carried out on one year old aonla rootstocks which were grown into 5×10 inch size black polybags in 2023 at Main Experiment Station of Horticulture, ANDUA&T Kumarganj Ayodhya. The experiment was laid out in Randomized Block Design comprises five treatment combinations consisting 5, 10, 15, 20 and 25 cm budding height from the ground level. The 15 cm height was treated as control. The observations were recorded on days taken for sprouting of bud, survival of buds after 15 DAB and 30 DAB, total sprouting, total died sprouts, total sprouts survival, length of sprout, Diameter of scion, number of determinate shoots, plant height, diameter of rootstocks just below union, diameter of scion just above union, number of leaves on determinate shoots and leaf chlorophyll content. Minimum days for bud sprouting (15.86 days), scion-shoot died percentage (0%) and maximum survival percentage (100%), total sprouting (90%) and total sprouts survival (100%) observed when patch budding was done at 25 cm budding height. Maximum length (81.56 cm) of sprouted shoot and diameter (6.03 mm) of scion, number of determinate shoots (85.05) and plant height (106.56 cm), rootstocks diameter (10.95 mm) just above union, diameter (7.53 mm) of scion just above union and number of leaves (92.52) were also recorded at 25 cm budding height. The total leaf chlorophyll content (3.31 mg/g) was found to be maximum at 25 cm budding height. It is concluded that the 25 cm budding height found to be the best for the propagation of aonla using patch budding in the month of June under North India condition on the one year old rootstocks grown in the polybags of 5×10 inch size.

**Keywords:** Patch budding, Budding height, Rootstocks, Aonla, *Emblica officinalis*

#### **1. Introduction:**

Aonla (*Emblica officinalis* Gaertn.) is a highly profitable fruit crop that can be cultivated on marginal and saline soils under rainfed conditions in the semi-arid and arid regions. Natural aonla plants can be found growing on hills up to an altitude of 1800 meters above sea level [10]. It is believed to be native to India, Malaysia, Sri Lanka and China, where it thrives and is often found in the wild. In India, it is commercially cultivated in Uttar Pradesh particularly in Pratapgarh, Sultanpur, Azamgarh, Varanasi, Jaunpur, Rai Bareilly, Agra and Bareilly districts. It is

an ideal fruit crop for planting in wastelands because of its drought tolerance, and low susceptibility to pests and diseases. It covers over 105,000 ha area in the country with 1276000 MT production (MoA & FW, 2022). Due to its resilience, high yield, nutritional and medicinal benefits, and versatility in value-added products, aonla has emerged as a significant fruit in the 21<sup>st</sup> century [10] which is highly profitable even with minimal care. Aonla fruits significance comes from its high vitamin C content, second only to Barbados cherry. Aonla contains 500-1500 mg of ascorbic acid per 100g of pulp [1], which is twenty times higher than the amount found in oranges. Aonla holds great commercial importance due to its distinct medicinal and industrial benefits. It is employed in Ayurvedic medicine to prepare Chyavanprash and Triphala. Its cultivation is getting expansion because of its hardy nature, medicinal and therapeutic values.

One of the obstacle to expand aonla cultivation is unavailability of quality planting materials of commercially important cultivars. Although the method of propagating of aonla through patch budding has been standardized [6,11 and 9], but success rate varies significantly across different agroclimatic conditions. Under North Indian conditions, patch budding from mid-May to September offers a good success rate when budding is done on one year old seedling rootstocks of pencil thickness.

The height at which budding occurs is crucial for its success and subsequent growth of scion shoot. An increase in budding height could also lead to a decrease in Trunk Cross-Sectional Area (TCA) over time, affecting tree growth annually due to the dwarfing effect [17]. Very little systematic work on this important aspect has yet been done in different region in India. Therefore, looking to the importance the different height in vegetative propagation, present research was taken to find out the best height for patch budding in aonla to get maximum success, survival and growth performance under Northern India condition.

## **2. Materials and Methods**

The investigation was carried out at Main Experiment Station of Horticulture, Acharya Narendra Deva University of Agriculture & Technology Kumarganj-Ayodhya, India in 2023 on one year old aonla rootstocks which were grown into 5×10-inch size black polybags filled with the similar mixture of soil and Farm Yard Manures. The experiment consisted four replications and five treatments comprising 5, 10, 15, 20 and 25 cm budding height from the ground level. The 15 cm height was treated as control because most of the budder performe patch budding in

aonla at 15 cm height. The scion buds were collected from 25 years old plant of NA-7 variety of aonla and patch budding was done in the month of June. The Vernier Caliper and Meter scale were used to record the diameter and height, respectively. While the observations on sprouting attributes and growth parameters were recorded using following method:

### **2.1 Days taken for sprouting of buds**

The number of days from the budding date to the sprouting of the scion bud was recorded.

### **2.2 Survival of buds after 15 DAB and 30 DAB**

The survival percentage in the nursery was determined by counting the number of live buds out of the total patched buds at 15 and 30 days after budding.

### **2.3 Percentage of total sprouting**

The number of budded plants with green buds that sprouted till July 31st was counted to calculate the bud sprouting percentage. The percentage of bud sprouting was determined using the following formula:

$$\text{Total sprouting (\%)} = \frac{\text{No. of budded plants having sprouted buds}}{\text{Total No. of budded plants}} \times 100$$

### **2.4 Total died sprouted buds:**

The number of sprouted buds that died after sprouting was counted to determine the percentage of sprouted scion buds died. The percentage was calculated using the following formula:

$$\text{Total died sprouted buds (\%)} = \frac{\text{No. of budded plants of died buds after sprouting}}{\text{Total number of plants of sprouted buds}} \times 100$$

### **2.5 Total sprouts survival:**

The percentage of budded plants that ultimately survived was determined by the following formula:

$$\text{Total sprouts survival (\%)} = \frac{\text{Survived plants}}{\text{Budding success plants}} \times 100$$

### **2.6 Leaf chlorophyll content:**

The 50 mg of finely chopped fresh leaf sample was placed in the test tubes to which 10 ml Dimethyl sulphoxide (DMSO) is added and tubes are covered with aluminium foil and kept in an oven at 65°C for 4 hrs. The absorbance of the chlorophyll solution was read at 663 and 645 nm using Spectrophotometer modal LT-2203. DMSO was used to recorded blank reading [2].

The chlorophyll content of leaves was calculated using following formula:

$$\text{Chl 'a'} = 12.7 \times D_{663} - 2.69 \times D_{645} \times \frac{V}{1000 \times W}$$

$$\text{Chl 'b'} = 22.9 \times D_{645} - 4.68 \times D_{663} \times \frac{V}{1000 \times W}$$

$$\text{Total chlorophyll(a + b)} = 20.2 \times D_{645} + 8.02 \times D_{663} \times \frac{V}{1000 \times W}$$

Where, D (Optical density), V (Final volume of DMSO (ml)), W (Weight of plant tissue (gm)), Chl 'a' (Chlorophyll 'a'), Chl 'b' (Chlorophyll 'b'), Unit (mm/g).

### 3. Results and Discussion

#### 3.1 Days taken for sprouting of bud

The data presented in Table 1 show that Minimum days (15.86 days) for bud sprouting were observed when plants were budded at 25 cm height (T<sub>5</sub>) followed by 18.59 days and 19.33 days at 20 cm height (T<sub>4</sub>) and 15 cm budding height (T<sub>2</sub>), respectively. While 5 cm budding height (T<sub>2</sub>) took maximum days (24.37 days) for sprouting whereas, 10 cm budding height (19.50 days) took significantly less days for sprouting in comparison to 5 cm budding height (T<sub>2</sub>). He reported that minimum days (16.40 days) for bud sprouting required in veneer grafting at 20 cm grafting height during the August month in mango [13]. The proximity of the budding point to source of photosynthates might be the reason of less days taken for sprouting of buds at higher height. The published report reveals that relative humidity is a key factor in bud sprouting and higher humidity, leads to early bud sprouts in guava [14]. These reports are in support that days taken for sprouting of bud is also influenced by the plant species and environmental condition because of these reasons the finding of present studies differ with the earlier published works on other fruits.

**Table 1: Effect of budding height on Days taken for sprouting of bud**

Treatments	Days taken for sprouting
	(In days)
T <sub>1</sub> - Control (15 cm)	19.33
T <sub>2</sub> - 5 cm	24.37
T <sub>3</sub> - 10 cm	19.50
T <sub>4</sub> - 20 cm	18.59
T <sub>5</sub> - 25 cm	15.86
<b>SEm ±</b>	<b>0.37</b>
<b>CD at 5%</b>	<b>1.15</b>

**3.2 Survival of bud after 15 DAB and 30 DAB**

The data presented in Table 2 show that maximum 95% and 90% survival percentage as marked green bud was found after 15 DAB at 25 cm budding height (T<sub>5</sub>) and 30 DAB at 25 cm budding height (T<sub>5</sub>), respectively. While minimum 70% and 50% survival percentage were recorded after 15 DAB at 5 cm budding height (T<sub>2</sub>) and 30 DAB at 5 cm budding height (T<sub>2</sub>), respectively. The difference was found to be significant. The findings indicate that survival of buds decreased with time which might be fluctuation in environment and interference of physiological factors in the healing of the plants.

**Table 2: Effect of budding height on survival of bud**

Treatments	Survival of bud (%)	
	15 Days After Budding	30 Days After Budding
T <sub>1</sub> - Control (15 cm)	80	75
T <sub>2</sub> - 5 cm	70	50
T <sub>3</sub> - 10 cm	73	60
T <sub>4</sub> - 20 cm	90	84
T <sub>5</sub> - 25 cm	95	90
<b>SEm ±</b>	<b>1.85</b>	<b>1.44</b>

<b>CD at 5%</b>	<b>5.75</b>	<b>4.48</b>
-----------------	-------------	-------------

### 3.3 Total Sprouting

The data presented in Table 3 reveals that the highest (90%) percentage of sprouting was observed at 25 cm budding height (T<sub>5</sub>) followed by 80% and 70% at 20 cm budding height (T<sub>4</sub>) and 15 cm budding height (T<sub>1</sub>), respectively. While, lowest (45%) success percentage was recorded at 5 cm budding height (T<sub>2</sub>). The effect of budding height was significant on bud sprouting in comparison to control. The result aligns with the discoveries in Bael [16] and in Ber [8]. In addition to proximity of buds to source, the basipetal and acropetal movement of hormones might also have influenced on the sprouting of buds.

### 3.4 Total Died Sprouts

The data presented in Table 3 show that lowest rate (0%) of sprouts scion-shoot died percentage has been observed at 25 cm budding height (T<sub>5</sub>) followed by 6.25% and 15% at 20 cm budding height (T<sub>4</sub>) and 15 cm budding height (T<sub>1</sub>), respectively. While 5 cm budding height (T<sub>2</sub>) has shown highest (30%) sprouts scion-shoot died percentage after 60 days of budding. These scientists observed that after ninety days of grafting, the lowest mortality rate (19.17%) was noted at 6 cm height of rootstock in Mango [7]. The proximity of sink to source of photosynthates, hormones and other healing factors might be the reason of difference in sprouts scion-shoot died percentage.

### 3.5 Total Sprouts Survival

Data presented in Table 3 reveals that maximum (100%) survival percentage was found at 25 cm budding height (T<sub>5</sub>) followed by 93.75% and 85% at 20 cm budding height (T<sub>4</sub>) and 15 cm budding height (T<sub>1</sub>), respectively. While minimum (70%) survival percentage was found at 5 cm height (T<sub>2</sub>). They recorded the highest survival rate (80%) at 15 cm rootstock height in soursop [18]. Earlier work reported that in veneer grafting, the highest survival percentage (78.50%) was recorded in August at 20 cm grafting height in Mango [13]. These reports confirm that height effects the survival percentage of buds and grafts.

**Table 3: Effect of budding height on Sprouting attributes**

Treatments	Sprouting attributes 60 Days After Budding (DAB)		
	Total Sprouting (%)	Total Died Sprouts (%)	Total Sprouts Survival (%)
T <sub>1</sub> - Control (15 cm)	70.00	15.00	85.00
T <sub>2</sub> - 5 cm	45.00	30.00	70.00
T <sub>3</sub> - 10 cm	50.00	20.00	80.00
T <sub>4</sub> - 20 cm	80.00	6.25	93.75
T <sub>5</sub> - 25 cm	90.00	0.00	100
<b>SEm ±</b>	<b>1.42</b>	<b>0.34</b>	<b>1.46</b>
<b>CD at 5%</b>	<b>4.43</b>	<b>1.06</b>	<b>4.55</b>

### 3.6 Length of sprout

The data outlined in Table 4 indicates that maximum (81.56 cm) length of sprouted shoot was found at 25 cm budding height (T<sub>5</sub>) after 90 DAB followed by 79.18 cm and 75.50 cm at 20 cm budding height (T<sub>4</sub>) and 15 cm budding height (T<sub>1</sub>), respectively. Whereas, minimum (72.86 cm) length of sprouted shoot was found at 5 cm budding height (T<sub>2</sub>) after 90 DAB. The maximum (16.27cm) shoot length was reported on 15th September at 60 cm height in mango [4]. The observed maximum (39.72 cm) scion shoot length has also been reported at 15 cm grafting height in Nagpur Mandarin [15]. These reports indicate that budding and grafting height influence the performance of the scion which are in support of present investigation. The distance of budding point from source of photosynthate and phytohormones including the osmotic pressure with gravitational force might be the reason of variation in performance of scion shoots.

**Table 4: Effect of budding height on length (cm) of sprout**

Treatments	Days after budding (DAB)									
	28 DAB	35 DAB	42 DAB	49 DAB	56 DAB	63 DAB	70 DAB	77 DAB	84 DAB	90 DAB
T <sub>1</sub> - Control (15 cm)	5.63	12.53	20.46	27.89	35.14	43.97	51.32	60.50	69.32	75.50
T <sub>2</sub> - 5 cm	2.56	9.87	16.59	22.06	30.35	38.31	47.66	57.51	65.19	72.86
T <sub>3</sub> - 10 cm	4.65	11.76	18.63	25.43	32.52	41.12	49.50	59.11	68.96	77.35
T <sub>4</sub> - 20 cm	6.08	13.87	21.59	29.16	37.65	45.38	53.76	62.15	71.34	79.18
T <sub>5</sub> - 25 cm	6.81	14.56	22.41	30.75	39.33	47.21	56.13	65.42	73.65	81.56
<b>SEm ±</b>	<b>0.11</b>	<b>0.25</b>	<b>0.40</b>	<b>0.54</b>	<b>0.69</b>	<b>0.85</b>	<b>1.01</b>	<b>1.19</b>	<b>1.35</b>	<b>1.50</b>
<b>CD at 5%</b>	<b>0.35</b>	<b>0.78</b>	<b>1.23</b>	<b>1.68</b>	<b>2.16</b>	<b>2.65</b>	<b>3.15</b>	<b>3.70</b>	<b>4.22</b>	<b>4.67</b>

### 3.7 Diameter of scion

Data presented in Table 5 reveals that the diameter of the scion after 90 DAB was maximum (6.03 mm) at 25 cm budding height (T<sub>5</sub>) followed by 5.86 mm and 5.28 mm at 20 cm budding height (T<sub>4</sub>) and 15 cm budding height (T<sub>1</sub>), respectively. While minimum (3.37 mm) diameter of scion was recorded at 5 cm budding height (T<sub>2</sub>). The present results indicate that scion diameter is directly proportional to proximity of budding point to apical buds and leaves on the rootstocks. This might be due to more rapid diversion of food from source to sink.

**Table 5: Effect of budding height on diameter (mm) of scion**

Treatments	Days after budding (DAB)									
	28 DAB	35 DAB	42 DAB	49 DAB	56 DAB	63 DAB	70 DAB	77 DAB	84 DAB	90 DAB
T <sub>1</sub> - Control (15 cm)	0.60	1.18	1.62	2.19	2.74	2.98	3.50	4.12	4.77	5.28
T <sub>2</sub> - 5 cm	0.28	0.75	1.15	1.69	2.05	2.52	2.98	2.52	2.89	3.37
T <sub>3</sub> - 10 cm	0.42	0.96	1.47	1.93	2.38	2.80	3.21	3.66	4.05	4.65
T <sub>4</sub> - 20 cm	0.96	1.48	2.02	2.45	2.87	3.14	3.83	4.39	5.28	5.86
T <sub>5</sub> - 25 cm	1.04	1.63	2.14	2.61	3.09	3.75	4.32	4.91	5.56	6.03
<b>SEm ±</b>	<b>0.02</b>	<b>0.03</b>	<b>0.04</b>	<b>0.04</b>	<b>0.05</b>	<b>0.06</b>	<b>0.07</b>	<b>0.08</b>	<b>0.09</b>	<b>0.10</b>
<b>CD at 5%</b>	<b>0.05</b>	<b>0.08</b>	<b>0.11</b>	<b>0.14</b>	<b>0.16</b>	<b>0.19</b>	<b>0.22</b>	<b>0.25</b>	<b>0.29</b>	<b>0.32</b>

### 3.8 Number of determinate shoots

The data presented in Table 6 recommends that maximum (85.05) number of determinate shoots were recorded at 25 cm budding height (T<sub>5</sub>) after 90 DAB followed by 81.38 and 79.35 at 20 cm budding height (T<sub>4</sub>) and 15 cm budding height (T<sub>1</sub>), respectively. While minimum (72.45) number of determinate shoots were recorded at 5 cm budding height (T<sub>2</sub>). The number of determinate shoots is increased with length of the indeterminate shoots. She recorded that in the second week of September, maximum (5.0) number of determinate shoots were found followed by the budding performed in fourth week of September (3.4) whereas, the minimum (1.5) shoots were observed during the second week of October in aonla [5]. This supports that month of propagation of fruit trees and growth of indeterminate shoot both influence the performance of scion-shoot.

**Table 6: Effect of budding height on number of determinate shoots**

Treatments	Days after budding (DAB)									
	28 DAB	35 DAB	42 DAB	49 DAB	56 DAB	63 DAB	70 DAB	77 DAB	84 DAB	90 DAB
T <sub>1</sub> - Control (15 cm)	4.29	11.31	17.40	25.62	35.54	44.02	52.93	62.87	70.82	79.35
T <sub>2</sub> - 5 cm	2.95	9.33	14.46	22.50	30.75	39.16	47.65	56.36	64.18	72.45
T <sub>3</sub> - 10 cm	3.83	10.75	16.04	23.65	33.48	42.83	51.44	60.31	68.25	76.95
T <sub>4</sub> - 20 cm	4.74	11.52	18.11	26.25	36.19	45.56	55.13	64.19	72.25	81.38
T <sub>5</sub> - 25 cm	5.94	13.08	19.86	28.25	38.56	48.16	57.94	67.30	75.68	85.05
<b>SEm ±</b>	<b>0.09</b>	<b>0.22</b>	<b>0.34</b>	<b>0.47</b>	<b>0.69</b>	<b>0.86</b>	<b>1.04</b>	<b>1.21</b>	<b>1.31</b>	<b>1.54</b>
<b>CD at 5%</b>	<b>0.28</b>	<b>0.69</b>	<b>1.07</b>	<b>1.47</b>	<b>2.14</b>	<b>2.68</b>	<b>3.23</b>	<b>3.79</b>	<b>4.27</b>	<b>4.80</b>

### 3.9 Plant height

The data presented in Table 7 indicate that maximum (106.56 cm) plant height was recorded at 25 cm budding height (T<sub>5</sub>) followed by 99.18 cm and 90.50 cm at 20 cm budding height (T<sub>4</sub>) and 15 cm budding height (T<sub>1</sub>), respectively after 90 DAB. While minimum (77.86 cm) plant height was recorded at 5 cm budding height (T<sub>2</sub>). The published work reveals reported that maximum height (9.63 cm, 10.43 cm, 12.17 cm and 14.86 cm) of the graft was noted at 30, 60, 90, and 120 days after grafting at 15 cm compared to 5 cm and 10 cm [3]. The other researchers observed that maximum (56.18 cm) plant height at 15 cm height of rootstocks in Nagpur Mandarin [15]. These published works are in support that grafting and budding height influenced the performance of plant height after propagation.

**Table 7: Effect of budding height on plant height (cm)**

Treatments	Days after budding (DAB)									
	28 DAB	35 DAB	42 DAB	49 DAB	56 DAB	63 DAB	70 DAB	77 DAB	84 DAB	90 DAB

T <sub>1</sub> - Control (15 cm)	20.63	27.53	35.46	42.89	50.14	58.97	66.32	75.50	84.32	90.50
T <sub>2</sub> - 5 cm	7.56	14.87	21.59	27.59	37.35	43.31	52.66	62.51	70.94	77.86
T <sub>3</sub> - 10 cm	14.65	21.76	28.63	35.43	42.52	51.12	59.50	69.11	78.96	87.35
T <sub>4</sub> - 20 cm	26.08	33.87	41.59	49.16	57.65	65.38	73.76	82.15	91.34	99.18
T <sub>5</sub> - 25 cm	31.81	39.56	47.41	55.75	64.33	72.21	81.13	90.42	98.65	106.56
<b>SEm ±</b>	<b>0.47</b>	<b>0.60</b>	<b>0.74</b>	<b>0.89</b>	<b>1.04</b>	<b>1.19</b>	<b>1.35</b>	<b>1.52</b>	<b>1.69</b>	<b>1.83</b>
<b>CD at 5%</b>	<b>1.46</b>	<b>1.88</b>	<b>2.32</b>	<b>2.76</b>	<b>3.23</b>	<b>3.71</b>	<b>4.20</b>	<b>4.74</b>	<b>5.26</b>	<b>5.71</b>

### 3.10 Diameter of rootstocks just below union

The data presented in Table 8 intimates that rootstocks diameter varied significantly with the budding height. The maximum (10.95 mm) rootstock diameter was recorded at 25 cm budding height (T<sub>5</sub>) followed by 10.43 mm and 9.98 mm at 20 cm budding height (T<sub>4</sub>) and 15 cm budding height (T<sub>1</sub>), respectively after 90 DAB. While minimum (6.80 mm) rootstock diameter was recorded at 5 cm budding height (T<sub>2</sub>). The maximum girth (2.46 cm) below the union at 90 DAG at 25 cm rootstock height and minimum girth (2.27 cm) was noted at 90 DAG at 10 cm height in Jackfruit [12]. Similarly, the maximum stock girth (9.33 mm) was measured at 60 cm grafting height and the minimum stock girth (8.55 mm) was measured at 20 cm rootstock height in Mango [4]. There earlier published works indicate that stock girth influenced by the height of propagation in fruits which are in support of the results of the present investigation.

Treatments	Days after budding (DAB)									
	28 DAB	35 DAB	42 DAB	49 DAB	56 DAB	63 DAB	70 DAB	77 DAB	84 DAB	90 DAB
T <sub>1</sub> - Control	5.96	6.80	7.02	7.56	7.95	8.41	8.93	9.19	9.51	9.98

(15 cm)										
T <sub>2</sub> - 5 cm	4.95	5.25	5.76	6.05	6.52	7.09	7.55	5.80	6.13	6.80
T <sub>3</sub> - 10 cm	5.46	5.87	6.25	6.97	7.33	7.82	8.16	8.45	8.72	9.14
T <sub>4</sub> - 20 cm	6.51	7.07	7.36	7.83	8.12	8.58	8.97	9.56	9.94	10.43
T <sub>5</sub> - 25 cm	6.65	7.44	7.92	8.45	8.99	9.38	9.83	10.06	10.52	10.95
<b>SEm ±</b>	<b>0.12</b>	<b>0.13</b>	<b>0.14</b>	<b>0.15</b>	<b>0.15</b>	<b>0.16</b>	<b>0.17</b>	<b>0.17</b>	<b>0.17</b>	<b>0.18</b>
<b>CD at 5%</b>	<b>0.37</b>	<b>0.41</b>	<b>0.43</b>	<b>0.46</b>	<b>0.48</b>	<b>0.51</b>	<b>0.53</b>	<b>0.51</b>	<b>0.53</b>	<b>0.56</b>

**Table 8: Effect of budding height on diameter of rootstocks just below union**

### 3.11 Diameter of scion just above union

Data pertaining in Table 9 show that maximum (7.53 mm) diameter of scion just above union at 25 cm budding height (T<sub>5</sub>) followed by 7.14 mm and 7.03 mm at 20 cm budding height (T<sub>4</sub>) and 15 cm budding height (T<sub>1</sub>), respectively after 90 DAB. While minimum (4.98 mm) diameter of scion just above union noted at 5 cm budding height (T<sub>2</sub>). The maximum girth (2.80 cm) above the union on 90 DAG was reported at 15 cm rootstock height and minimum girth (2.60 cm) on 90 DAG at 10 cm height in Jackfruit [12]. The maximum diameter (0.90 cm) above 2.5 cm of the graft union at 15 cm grafting height was also reported in Nagpur Mandarin [17]. These published research are in supports of the findings of the present investigation.

**Table 9: Effect of budding height on diameter of scion just above union**

Treatments	Days after budding (DAB)									
	28 DAB	35 DAB	42 DAB	49 DAB	56 DAB	63 DAB	70 DAB	77 DAB	84 DAB	90 DAB
T <sub>1</sub> - Control (15 cm)	1.06	3.10	3.54	4.05	4.50	4.89	5.23	5.78	6.37	7.03
T <sub>2</sub> - 5 cm	0.42	1.06	2.18	2.49	2.74	3.08	3.47	3.93	4.41	4.98

T <sub>3</sub> - 10 cm	0.66	1.40	2.49	2.61	2.93	3.77	3.94	4.29	4.84	5.24
T <sub>4</sub> - 20 cm	1.54	3.78	4.06	4.59	5.01	5.33	5.83	6.04	6.83	7.14
T <sub>5</sub> - 25 cm	2.96	4.08	4.47	4.94	5.19	5.58	6.03	6.50	7.08	7.53
<b>SEm ±</b>	<b>0.04</b>	<b>0.06</b>	<b>0.07</b>	<b>0.08</b>	<b>0.09</b>	<b>0.09</b>	<b>0.10</b>	<b>0.11</b>	<b>0.12</b>	<b>0.13</b>
<b>CD at 5%</b>	<b>0.12</b>	<b>0.20</b>	<b>0.22</b>	<b>0.25</b>	<b>0.27</b>	<b>0.29</b>	<b>0.32</b>	<b>0.34</b>	<b>0.38</b>	<b>0.40</b>

### 3.12 Number of leaves on determinate shoots

The present data in Table 10 express that the maximum (92.52) number of leaves were observed at 25 cm budding height (T<sub>5</sub>) followed by 90.15 and 89.49 at 20 cm budding height (T<sub>4</sub>) and 15 cm budding height (T<sub>1</sub>), respectively after 90 DAB. However, minimum (83.55) number of leaves noted at 5 cm budding height (T<sub>2</sub>). The earlier research also recorded the maximum (16.47) leaves on 30<sup>th</sup> August at 60 cm height and the minimum (8.36) number of leaves at 60 cm heights in Mango [4]. The more number of leaves at higher height might be because of availability of more photosynthates and plant hormones.

**Table 10: Effect of budding height on Number of leaves on determinate shoots**

Treatments	Days after budding (DAB)									
	28 DAB	35 DAB	42 DAB	49 DAB	56 DAB	63 DAB	70 DAB	77 DAB	84 DAB	90 DAB
T <sub>1</sub> - Control (15 cm)	16.25	22.98	28.49	36.38	45.56	54.69	63.72	71.50	80.16	89.49
T <sub>2</sub> - 5 cm	14.32	19.14	25.81	31.23	40.21	48.59	57.86	66.47	75.24	83.55
T <sub>3</sub> - 10 cm	15.50	21.58	27.74	34.89	43.16	51.37	60.52	69.55	77.59	86.02
T <sub>4</sub> - 20 cm	17.18	24.30	30.25	38.73	47.92	56.36	64.54	72.90	81.07	90.15
T <sub>5</sub> - 25 cm	19.05	25.59	32.11	40.60	49.56	57.42	66.93	73.32	84.40	92.52

<b>SEm ±</b>	<b>0.33</b>	<b>0.45</b>	<b>0.57</b>	<b>0.71</b>	<b>0.89</b>	<b>1.05</b>	<b>1.22</b>	<b>1.37</b>	<b>1.55</b>	<b>1.72</b>
<b>CD at 5%</b>	<b>1.01</b>	<b>1.40</b>	<b>1.77</b>	<b>2.24</b>	<b>2.77</b>	<b>3.27</b>	<b>3.81</b>	<b>4.28</b>	<b>4.83</b>	<b>5.35</b>

### 3.13 Leaf chlorophyll content at 60 DAB

The data presented in Table 11 show that maximum (3.31 mg/g) total leaf chlorophyll content was observed at 25 cm budding height (T<sub>5</sub>) followed by 2.86 mg/g and 2.75 mg/g at 20 cm budding height (T<sub>4</sub>) and 15 cm budding height (T<sub>1</sub>), respectively. Whereas, minimum (2.49 mg/g) total leaf chlorophyll content was recorded at 5 cm budding height (T<sub>2</sub>). Maximum (2.30 mg/g) leaf chlorophyll 'a' content was observed at 25 cm budding height (T<sub>5</sub>) followed by 1.94 mg/g and 1.91 mg/g at 20 cm budding height (T<sub>4</sub>) and 15 cm budding height (T<sub>1</sub>), respectively. Whereas, minimum (1.68 mg/g) leaf chlorophyll 'a' content was recorded at 5 cm budding height (T<sub>2</sub>). Maximum (1.01 mg/g) leaf chlorophyll 'b' content was observed at 25 cm budding height (T<sub>5</sub>) followed by 0.93 mg/g and 0.84 mg/g at 20 cm budding height (T<sub>4</sub>) and 15 cm budding height (T<sub>1</sub>), respectively. Whereas, minimum (0.81 mg/g) leaf chlorophyll 'b' content was recorded at 5 cm budding height (T<sub>2</sub>). In present studies observed highest total chlorophyll content in the leaves of scion shoot at 25 cm height which might be due to more exposure to the sun light intensity and more uptake of nutrients due to osmotic pressure built up by transpiration because of more leaves.

**Table 11: Effect of budding height on Leaf chlorophyll content of leaves**

Treatments	Chlorophyll content (mg/g) at 60 DAB		
	Chlorophyll 'a'	Chlorophyll 'b'	Total chlorophyll
T <sub>1</sub> - Control (15 cm)	1.91	0.84	2.75
T <sub>2</sub> - 5 cm	1.68	0.81	2.49
T <sub>3</sub> - 10 cm	1.78	0.83	2.61

T <sub>4</sub> - 20 cm	1.94	0.93	2.86
T <sub>5</sub> - 25 cm	2.30	1.01	3.31
<b>SEm ±</b>	<b>0.04</b>	<b>0.02</b>	<b>0.06</b>
<b>CD at 5%</b>	<b>0.12</b>	<b>0.06</b>	<b>0.17</b>

#### 4. Conclusions

It is concluded on the basis of the result in present investigation that the 25 cm budding height is best for the propagation of aonla using patch budding in the June month under North India condition on the one-year-old rootstocks grown in the polybags of 5×10 inch.

#### Reference:

1. Chauhan, O.P. Srivastava, S., Pandey, P., Rao, G.K. A study on the development of aonla blended sauce. Beverage and Food world. 2005;32:31–33.
2. Hiscox JD, Israelstam GF. A method for the extraction of chlorophyll from leaf tissue without maceration. Canadian journal of botany. 1979;57(12):1332-1334.
3. Jawre, R. Response of mango varieties at different height of grafting on rootstock in polyhouse. Thesis, M.Sc. Jawaharlal Nehru Krishi Vishwa Vidyalaya, Jabalpur. 2012;29-52.
4. Karna, K.A., Varu, K.D., Panda, A.P., Hota, D. Standardization of grafting time and height on success of softwood grafting in mango (*Mangifera indica* L.). J. Pharmacogn. Phytochem. 2017;6(5):2803-2807.
5. Kumari, A. Propagational studies in aonla (*Emblca officinalis gaertn*). Thesis, M.Sc. CCS Haryana Agricultural University Hisar. 2002;23.
6. Nand, D. Budding in aonla (*Phyllunthus emblica*). *Sci. Cult.* 1962;28:486.
7. Nalage, N.A., Magar, S.D., Bhosale, S.S., Mhetre, D.A. Effect of height of rootstock on success of epicotyl grafting in mango (*Mangifera indica* L.) cv. Kesar. *Int. J. Agric. Sci.* 2010;6(1):124-128.

8. Nath, V., Saroj, P.L., Singh, R.S., Bhargav, R., Pareek, O.P. In situ establishment of ber orchard under hot arid ecosystem of Rajasthan. *Ind. J. Horti.* 2000;57:21-26.
9. Pathak, R.K., C.M. Ojha, R. Dwivedi and Hari Om. Studies on the effect of method and duration of budding in aonla. *Indian J. Hort.* 1991;47:207-12.
10. Pathak R K. Status Report on Genetic Resources of Indian Gooseberry-Aonla (*Emblica officinalis* Gaertn) in South and Southeast Asia. IPGRI Office for South Asia National Agriculture Science Centre (NASC) DPS Marg, Pusa Campus, New Delhi, India, 2003;1-96.
11. Pandey, I.C., K.S. Prasad. Propagation of aonla by budding. *Prog. Hort.* 1980;17:345-346.
12. Prajapati, I. Effect of height of rootstock and length of scion on success of epicotyl grafting in jackfruit (*Artocarpus heterophyllus* Lam.) cv. Konkan prolific. Thesis, M.Sc. Navsari Agricultural University, Gujarat. 2011;25-42.
13. Ram, C., Prakash, S., Kumar, V., Kumar, M., Braj, M. 2013. Response of amrapali mango to grafting height, time and techniques of grafting. *Society for Sci. Dev. in Agric. and Tech.* 8: 898-902.
14. Singh G, Gupta S, Mishra R, Singh A. Technique for rapid multiplication of guava (*Psidium guajava* L.). *Acta Horti.* 2007;735:177-183.
15. Thokchom, A., Singh, PKD. Effect of grafting height and scion length on growth of citrus reticulata cv. Nagpur Mandarin grafts. *Int. J. Chem. Stud.* 2018;6(2):2094- 2097.
16. Tripathi, A., Kumar, R. Studies on the effect of method and time of budding in bael (*Aegle marmelos* L.). *Haryana J. Horti. Sci.* 2004;33:195-198.
17. Weibel, A. M., Reighard, G. L. Interstems but not grafting height control vegetative growth of young 'Redhaven' peach trees. In IX International Symposium on Integrating Canopy, Rootstock and Environmental Physiology in Orchard Systems, 2011,903:513-519.
18. Yakubu, F.B., Alaje, V.I., Olaniyi, A.A., Nola, M.O., Odewale, M.A., Fadulu, O.O., Adeniyi, K.K. Influence of scion length and point of attachment on rootstock on survival and growth of Grafted Soursop. *Afr. Crop Sci. J.* 2022;30(1): 77-85.