

Original Research Article

Efficient and inexpensive bio adsorbent for the removal of Safranin: Kinetic and Isotherm study

ABSTRACT

Alstonia Scholaris leaves after treatment with 0.01 M NaOH was used for the adsorptive removal of safranin. As the initial safranin concentration increases from 20 mg/L to 80 mg/L the dye removal increases up to 91 % in 60 min contact time at optimum pH of 8. Pseudo second order model and Freundlich isotherm was followed by the adsorption experimental data.

Key Words: Adsorption, Cotton red, saptaparni, basic red

1. INTRODUCTION

India is a developing country and with the development in the industrial sector like paper, pulp, leather and textile the consumption of synthetic dyes is also increases, though at most care were taken by these industries, some of the dyes and pigments residues likely to be present in the effluents. The presence of synthetic dyes in traces can affect the aquatic and human life [1]. Most of the synthetic dyes have carcinogenic and mutagenic effects [2]. Various chemical, physical and biological methods were used to remove hazardous material from water. Due to the non-biodegradable nature of some of the dyes, the microbial degradation methods have limitation [3]. The **chemical and physical methods** like adsorption [4], oxidation, ozonolysis, chemical and photochemical destruction [5], coagulation, filtration. From the above methods adsorption method is mostly applied due its cost effective and simplicity [6]. Activated charcoal is the best adsorbent but it is not cost effective so researcher preferably uses low cost adsorbent form the waste material like Gmelina aborea [7], pomegranate peel [8] mango leaf powder [9].

Alstonia Scholaris is commonly called as saptaparni or devil plants grown in different parts of India and easily available. Various adsorbent were reported for the removal of safranin such as Snail Shell [10], soybean hull [11], zeolites [12], red mud [13], Citrus reticulata peels[14] etc. Safranin is a cationic dye also known as cotton red, the specification are given in table1




Fig. 1 Chemical structure of Safranin

Table1 Specification of Safranin

Molecular Formula	C ₂₀ H ₁₉ N ₄ Cl
Molecular Weight	350.85
λ max (nm)	521

So in the present adsorption study, the effectiveness of leaves powder after treatment with NaOH was investigated. The data was analyzed for kinetic and isotherm model

2. MATERIALS AND METHODS

All chemicals were purchased from Loba Chemie Pvt Ltd. India and were used without further purification. Elico double beam spectrophotometer SL-210 was used to determine the absorbance. Equiptronics pH meter and Remi stirrer was used. 0.1 M NaOH and 0.1M HCl were used to adjust the pH. The dried leaves of *Alstonia Scholaris* were collected from the institute, grinded and washed with water. The powder was treated with 0.01 M NaOH solution and washed with water till the pH is neutral. The residue was dried in a hot air oven overnight and used for as an adsorbent. By measuring absorption at 521 nm and with the help of standard curve the concentration of Safranin was determined.

The milligram of Safranin dye on adsorbent (q_e) was determined using the following equation

$$q_t = \frac{(C_i - C_t)}{W} v$$

Where C_i and C_t were concentration of Safranin, v is volume of dye solution and w is weight of adsorbent

To determine the optimum condition of adsorption, the adsorption study was performed using batch adsorption method. In this 25 mL dye solution was taken in 100 mL flask and stirred with 0.05 g of adsorbent powder for specific time.

3. Result and Discussion

3.1 Effect of Initial Amount of Adsorbent

The effect of adsorbent dose was assessed by altering the adsorbent dose from 0.02 to 0.15 g. The result shows that the percentage removal of Safranin increases with increase in the initial amount of adsorbent (77 to 89 %) (as shown in Fig 2). The increase in the initial adsorbent dose increases the site for adsorption thus increases the percentage removal.




Fig. 2 Effect of amount of adsorbent

3.2 Effect of Initial pH

The initial H^+ ion concentration (pH) mostly affect the dye adsorption process [15]. It has been observed from experimental result that the increase in pH increases the dye

adsorption, this trend is observed up to pH 8, and further increase in pH decreases the removal. (Fig. 3)




Fig.3. effect of initial pH

3.3 Effect of initial dye concentration and contact time

Batch adsorption study was performed by altering initial dye concentration (20 mg/L, 40 mg/L, 60 mg/L and 80 mg/L) and contact time at optimum pH with 0.1 g of adsorbent. With increase in initial dye concentration and contact time the percentage removal increases up to 91% this may be due to mono layer formation [16]




Fig4. Effect of initial dye concentration and contact time

3.4 Kinetics Studies

The experimental data was studied by applying three kinetic model viz. pseudo first order model, pseudo second order model and Elovic model

The following equation were used

$$\log(q_e - q_t) = \log q_e - \frac{k_1 t}{2.303} \quad (\text{pseudo first order equation}) [16]$$

Where k_1 is the pseudo first order rate constant and the value is determine from a plot of $\log(q_e - q_t)$ versus t .

$$q_t = \frac{1}{\beta} \ln(\alpha\beta) + \frac{1}{\beta} \ln(t) \quad (\text{Elovic model}) [17]$$

The elovic parameters were determined from the plot of q_t versus $\ln t$

$$\frac{t}{q_t} = \frac{1}{q_e^2 k_2} + \frac{t}{q_t} \quad (\text{pseudo second order equation}) [18]$$

K_2 is pseudo second order rate constant, the values were determined from a plot of t/q_t versus t




Fig.5 pseudo second order plot

The above parameters were represented in table 2

Table 2. Parameters for pseudo first-order , pseudo second-order adsorption and Elovich model

Dye Con c. (mg L ⁻¹)	First order				Second order				Elovich model		
	K_1 (min ⁻¹)	$q_{e(\text{exp})}$ (mg g ⁻¹)	$q_{e(\text{cal})}$ (mg g ⁻¹)	R^2	K_2 (min ⁻¹)	$q_{e(\text{exp})}$ (mg g ⁻¹)	$q_{e(\text{cal})}$ (mg g ⁻¹)	R^2	β (mg g ⁻¹)	α (mg g ⁻¹ min ⁻¹)	R^2
20	0.0859	8.158 9	2.6224	0.956 5	0.067 5	8.158 9	8.389 2	0.999 7	1.516 9	4.677 5	0.967 2
40	0.0773	17.30 6	5.6636	0.895	0.026 8	17.30 6	17.85 7	0.999 3	0.721 2	9.767 6	0.942 5
60	0.0660	26.40 9	4.8350 3	0.860 1	0.028 8	26.40 9	26.88 1	0.999 6	0.796 5	61.84 1	0.911
80	0.0700 1	36.77 8	8.1583	0.961 7	0.018 0	36.77 8	37.59 3	0.999 7	0.452 3	40.19 6	0.974 3

It has been observed from the R^2 values that the most suitable kinetic model for present study is the pseudo second order kinetic model. Similar trend was observed in the literature [17-19]

3.5 Adsorption isotherm

The three isotherm model viz. Temkin isotherm, Langmuir isotherm and Freundlich isotherm were applied for the present study.

The following equations were used

$$q_e = b \ln A + b \ln C_e \quad \text{Temkin isotherm [10]}$$

The Temkin parameters A and B were determined from the plot of q_e versus $\ln C_e$

$$\frac{C_e}{q_e} = \frac{C_e}{q_m} + \frac{1}{b q_m} \quad \text{Langmuir isotherm [20]}$$

The Langmuir constant q_m and b were determined from a plot of C_e/q_e versus C_e

$$\log q_e = (1/n)\log C_e + \log k_f \text{ Freundlich isotherm [21]}$$

The values of n and k_f were determined from plot of $\log q_e$ versus $\log C_e$

Table 3 Adsorption isotherm parameters

Langmuir Isotherm			Freundlich Isotherm			Temkin Isotherm		
b (L mg ⁻¹)	q _m (mg g ⁻¹)	R ²	n	K _f (mg g ⁻¹)	R ²	A (L mg ⁻¹)	B (J mole ⁻¹)	R ²
-11.952	4.9504	0.8926	-2.9403	4.6387	0.9938	4.2975	-35.366	0.9711

From the values of R² it has been observed that the present study follows Freundlich isotherm model.

4. Conclusion

The 91% removal of Safranin was observed at optimum pH 8, 80 min contact time with 80 mg/L dye concentration. The present experimental data follows pseudo second order kinetic model and Freundlich isotherm model. Alstonia Scholaris leaves powder after alkali treatment can be used as cheap and efficient adsorbent for the removal of safranin

References

- 1 Chowdhury S, Saha P, Adsorption Kinetic Modeling of Safranin onto Rice Husk Biomatrix Using Pseudo-first- and Pseudo-second-order Kinetic Models: Comparison of Linear and Non-linear Methods, Clean-Soil Air Water(2011) 39, 274, <https://doi.org/10.1002/clen.201000170>
- 2 Mathur N, Bhatnagar P, Mutagenicity assessment of textile dyes from Sanganer (Rajasthan), Journal of Environmental Biology (2007) 28,1,123.
- 3 Garg VK, Gupta R, Bala Yadav A, Kumar R, Dye removal from aqueous solution by adsorption on treated sawdust. Bioresource Technology. (2003), 89, (2):121. doi:10.1016/s0960-8524(03)00058-0. PMID: 12699929.
- 4 Yagub MT., Sen TK, Afroze S, Ang HM, Dye and its Removal from aqueous solution by Adsorption: A review, Advances in Colloid and Interface Science (2014), doi: 10.1016/j.cis.2014.04.002
- 5 Katheresan V, Kansedo J, Lau SY, Efficiency of Various Recent Wastewater Dye Removal Methods: A Review, Journal of Environmental Chemical Engineering (2018), <https://doi.org/10.1016/j.jece.2018.06.060>
- 6 Dutta S, Gupta B, Srivastava SK, Gupta AK, Recent advances on the removal of dyes from wastewater using various adsorbents: a critical review, Materials Advances (2021), 2,4497, <https://doi.org/10.1039/d1ma00354b>
- 7 Bello OS, Alabi EO, Adegoke KA, Adegboyega SA, Inyinbor AA, Dada AO, Rhodamine B dye sequestration using Gmelina aborea leaf powder, Heliyon (2019), 5, <https://doi.org/10.1016/j.heliyon.2019.e02872>
- 8 Ghibate R, Senhaji O, Taouil R, Kinetic and thermodynamic approaches on Rhodamine B adsorption onto pomegranate peel, Case Studies in Chemical and Environmental Engineering (2021),3,100078 <https://doi.org/10.1016/j.cscee.2020.100078>
- 9 Khan TA, Sharma S, Ali I, Adsorption of Rhodamine B dye from aqueous solution onto acid activated mango (Magnifera indica) leaf powder: Equilibrium, kinetic and thermodynamic studies, Journal of Toxicology and Environmental Health Sciences(2011) 3,10,286
- 10 Zwiwer FB, Al-Da'amy MA, Kareem ET, Snail Shell (Rostellariella) as a Low Cost Adsorbent for Safranin Dye Removal from Aqueous Solution, Baghdad Science Journal, 2024, 21(4): 1296-1305, <https://doi.org/10.21123/bsj.2023.8311>
- 11 Chandane V, Singh VK, Adsorption of safranin dye from aqueous solutions using a low-cost agro-waste material soybean hull, Desalination and Water Treatment (2014) <http://dx.doi.org/10.1080/19443994.2014.991758>

- 12 Abukhadra MR, Mohamed AS, Adsorption Removal of Safranin Dye Contaminants from Water Using Various Types of Natural Zeolite, Silicon, <https://doi.org/10.1007/s12633-018-9980-3>
- 13 Sahu MK, Patel RK, Removal of safranin-O dye from aqueous solution using modified red mud: kinetics and equilibrium studies, RSC Advances, (2015), <https://doi.org/10.1039/c5ra15780c>
- 14 Januário EFD, Tay Vidovix TB, de Araújo LA, Beltran LB, Bergamasco R, Vieira AMS, Investigation of Citrus reticulata peels as an efficient and low-cost adsorbent for the removal of safranin orange dye, ENVIRONMENTAL TECHNOLOGY (2021) <https://doi.org/10.1080/09593330.2021.1946601>
- 15 Abukhadra MR, Mohamed AS, Adsorption Removal of Safranin Dye Contaminants from Water Using Various Types of Natural Zeolite, Silicon, <https://doi.org/10.1007/s12633-018-9980-3>
- 16 Kaur S, Rani S, Mahajan RK, Asif M, Gupta VK, Synthesis and adsorption properties of mesoporous material for the removal of dye safranin: Kinetics, equilibrium, and thermodynamics, Journal of Industrial and Engineering Chemistry, (2014), <http://dx.doi.org/10.1016/j.jiec.2014.06.019>
- 17 Ghosh I, Kar S, Chatterjee T, Bar N, Das SK, Adsorptive removal of Safranin-O dye from aqueous medium using coconut coir and its acid-treated forms: Adsorption study, scale-up design, MPR and GA-ANN modeling, Sustainable Chemistry and Pharmacy 19 (2021) 100374, <https://doi.org/10.1016/j.scp.2021.100374>
- 18 S, Faraji M, Yamini Y, Rajabi AA, Fe₃O₄ magnetic nanoparticles modified with sodium dodecyl sulfate for removal of safranin O dye from aqueous solutions, Desalination 270 (2011) 160–165, <http://dx.doi.org/10.1016/j.desal.2010.11.040>
- 19 Addad ME, Regti A, Slimani R, Lazar S, Assessment of the biosorption kinetic and thermodynamic for the removal of safranin dye from aqueous solutions using calcined mussel shells, Journal of Industrial and Engineering Chemistry 20 (2014) 717–724, <http://dx.doi.org/10.1016/j.jiec.2013.05.038>
- 20 Mohamed F, Abukhadra MR, Shaban M, Removal of safranin dye from water using polypyrrole nanofiber/Zn-Fe layered double hydroxide nanocomposite (Ppy NF/Zn-Fe LDH) of enhanced adsorption and photocatalytic properties, Science of the Total Environment 640–641 (2018) 352–363, <https://doi.org/10.1016/j.scitotenv.2018.05.316>
- 21 Ghaedi M, Hajjati S, Mahmudi Z, Tyagi I, Agarwal S, Maity A, Gupta VK, Modeling of competitive ultrasonic assisted removal of the dyes – Methylene blue and Safranin-O using Fe₃O₄ nanoparticles, Chemical Engineering Journal 268 (2015) 28–37, <http://dx.doi.org/10.1016/j.cej.2014.12.090>