

Influence of different planting dates on their survival and growth parameters of two cultivars of jamun (*syzygiumcumini*) c.v Goma Priyanka and Ra Jamun.

Abstract

The present investigation entitled “Influence of different planting dates on their survival and growth parameters of two cultivars of jamun (*syzygiumcumini*) cv Goma Priyanka and Ra Jamun”. The present investigation was carried out in randomized block design with 10 treatment, which were replicated thrice. The treatments were T₁ (Goma priyanka + 30 January), T₂ (Goma priyanka + 15 February), T₃ (Goma priyanka + 1 march), T₄ (Goma priyanka + 15 march), T₅ (Goma priyanka + 30 march), T₆ (Ra jamun + 30 January), T₇ (Ra jamun + 15 February), T₈ (Ra jamun + 1 march), T₉ (Ra jamun + 15 march), T₁₀ (Ra jamun + 30 march). On the basis of our experimental finding it can be concluded that the best date of planting for jamun was 15th of march. The best variety for prayagrajagro climatic condition was found to be Goma Priyanaka. Ultimately on the basis of our overall finding it can be concluded that when Goma Priyanka was planted at 15th march gives the best result in terms of vegetative growth of the jamun plant.

Keywords: Jamun, Date of Planting, Growth, Survivability %.

Introduction

The Jamun, (*Syzygium cumini*), family Myrtaceae, chromosome $2n=40$ is a nutritious fruit with a variety of uses. It is one of the most hardy fruit crops and can easily be grown in neglected and marshy areas where other fruit plants cannot be grown successfully. The fruit is a good source of iron, sugars, minerals, protein and carbohydrates etc. Fully ripened fruits are eaten as fresh fruit and can be processed into beverages like jelly, jam, squash, wine, vinegar and pickles.

Jamun, also known as *Syzygium cumini*, is a tropical evergreen tree native to the Indian subcontinent. It belongs to the Myrtaceae family and can grow up to 30 meters tall. The tree has a dense, spreading crown with glossy, dark green leaves. Its small, fragrant flowers are typically white or cream-colored and are followed by oblong-shaped fruits. The fruit of the jamun tree is a berry, usually ovoid or oblong, with a smooth, dark purple to black skin. Inside, it contains juicy, pinkish-purple flesh surrounding a single, large seed. Jamun fruit has a unique, tangy-sweet flavor with a slightly astringent taste. It is rich in vitamins, minerals, and antioxidants, making it a popular choice in traditional medicine and culinary applications. Jamun trees are often cultivated for their fruit, which is enjoyed fresh, as well as in jams, jellies, juices, and desserts.

The planting date significantly influences the survival and growth of jamun trees. Optimal planting times coincide with the beginning of the rainy season or during mild weather conditions, allowing the young trees to establish root systems before facing harsh environmental stresses. Planting during the rainy season ensures ample moisture for initial growth and reduces transplant shock. Early planting may promote better root establishment and growth, leading to higher survival rates and faster development. Conversely, late planting, particularly during dry or extreme weather conditions, can increase stress on the newly planted trees, resulting in lower survival rates and slower growth. Additionally,

planting during periods of waterlogging or frost can negatively impact root development and overall tree health. Overall, selecting the appropriate planting date is essential for maximizing the survival and growth of jamun trees, ensuring they establish strong root systems and adapt well to their environment, ultimately leading to healthy and productive orchards.

Materials and Methods

The present investigation entitled “Influence of different planting dates on their survival and growth parameters of two cultivars of jamun (*syzygium cumini*) cv Goma Priyanka and Ra Jamun” was carried out during 2022-23 at Horticulture Research Farm, Department of Horticulture, Sam Higginbottom University of Agriculture Technology and sciences, Naini, Prayagraj. The present investigation was carried out in randomized block design with 10 treatments, which were replicated thrice. The treatments were T_1 (Goma priyanka + 30 January), T_2 (Goma priyanka + 15 February), T_3 (Goma priyanka + 1 march), T_4 (Goma priyanka + 15 march), T_5 (Goma priyanka + 30 march), T_6 (Ra jamun + 30 January), T_7 (Ra jamun + 15 February), T_8 (Ra jamun + 1 march), T_9 (Ra jamun + 15 march), T_{10} (Ra jamun + 30 march). The objective of the experiment was to study the best variety suitable for prayagraj and its growth and survival percentage.

Results and Discussion

The maximum Survivability % was observed in T₃ (Goma priyanka + 1 march) with (78.87) % respectively, followed by T₂ (Goma priyanka + 15 February) with (77.54) % which were significantly superior over T₁₀ (Ra jamun + 30 march) with (68.23) %. Early planting allows for establishment before harsh conditions, potentially increasing survival rates. Experimentation across varied planting dates and monitoring survival rates can provide insights into the optimal timing for jamun plantation, aiding in maximizing survival percentages and overall crop success.

The minimum Mortality % was observed in T₃ (Goma priyanka + 1 march) with (21.13) respectively, followed by T₂ (Goma priyanka + 15 February) with (22.46) % which were significantly superior over T₁₀ (Ra jamun + 30 march) with (31.77) %. The influence of different planting dates on the Mortality percentage of jamun (*Syzygiumcumini*) varies due to climatic conditions, soil moisture, and growth stages. Planting during optimal conditions, such as the onset of the rainy season, typically results in higher Mortality rates due to favorable moisture levels and reduced stress

The maximum Plant height was observed in T₃ (Goma priyanka + 1 march) with (31.16) cm followed by T₂ (Goma priyanka + 15 February) with (30.42) cm which were significantly superior over T₁₀ (Ra jamun + 30 march) with (28.42) cm. Early planting allows for establishment before harsh

conditions, potentially increasing Plant height. Experimentation across varied planting dates and monitoring Plant height can provide insights into the optimal timing for jamun plantation, aiding in maximizing Plant height and overall crop success.

The maximum Number of leaves was observed in T₃ (Goma priyanka + 1 march) with (20.83) cm followed by T₂ (Goma priyanka + 15 February) with (20.19) which were significantly superior over T₁₀ (Ra jamun + 30 march) with (16.76) . Planting during optimal conditions, such as the onset of the rainy season, typically results in higher Number of leaves due to favorable moisture levels and reduced stress.

The maximum Plant spread was observed in T₃ (Goma priyanka + 1 march) with (21.73) cm followed by T₂ (Goma priyanka + 15 February) with (21.32) cm which were significantly superior over T₁₀ (Ra jamun + 30 march) with (18.08) cm . Planting during optimal conditions, such as the onset of the rainy season, typically results in higher Number of leaves due to favorable moisture levels and reduced stress. The increase in number of leaves simultaneously increases the plant sp

The maximum Leaf area was observed in T₃ (Goma priyanka + 1 march) with (12..57) cm² followed by T₂ (Goma priyanka + 15 February) with (12.35) cm²

which were significantly superior over T₁₀ (Ra jamun + 30 march) with (9.29) cm². Increase in somatic cell division of leaf increases the leaf area of the plant which also helps in the increasing he plant spread.

The minimum Days to Bud break was observed in T₃ (Goma priyanka + 1 march) with (3.96) days respectively, followed by T₂ (Goma priyanka + 15 February) with (4.25) days which were significantly superior over T₁₀ (Ra jamun + 30 march) with (9.89).

The maximum Number of branches was observed in T₃ (Goma priyanka + 1 march) with (10.76) respectively, followed by T₂ (Goma priyanka + 15 February) with (9.54) which were significantly superior over T₁₀ (Ra jamun + 30 march) with (4.23). . Increase in somatic cell division of branches increases the bud of the plant which also helps in the initiation of the

The maximum Leaf area index was observed in T₃ (Goma priyanka + 1 march) with (2.89) respectively, followed by T₂ (Goma priyanka + 15 February) with (2.76) which were significantly superior over T₁₀ (Ra jamun + 30 march) with (1.67).

The maximum chlorophyll content was observed in T₃ (Goma priyanka + 1 march) with (65.87)

respectively, followed by T₂ (Goma priyanka + 15 February) with (61.45) which were significantly superior over T₁₀ (Ra jamun + 30 march) with (45.62). Increasing chlorophyll content in plants enhances photosynthesis, the process crucial for converting light energy into chemical energy. This leads to improved growth, higher yields, and enhanced overall plant vigor. Additionally, higher chlorophyll levels can contribute to better stress tolerance and improved resistance against environmental challenges.

Conclusion

On the basis of our experimental finding it can be concluded that the best date of planting for jamun was 15th of march. The best variety for prayagrajagro climatic condition was found to be GomaPriyanaka. Ultimately on the basis of our overall finding it can be concluded that when Goma Priyanka was planted at 15th march gives the best result in terms of vegetative growth of the jamun plant.

Table 1 Influence of different planting date on Survivality%, Mortality %, Plant height, Number of leaves and Days to break of Jamun.

Symbol	Survivality %	Mortality %	Plant height (cm)					Number of Leaves					Days to bud break
			Initial	30 DAT	60 DAT	90 DAT	120 DAT	Initial	30 DAT	60 DAT	90 DAT	120 DAT	
T ₁	70.67	29.33	6.98	11.85	17.52	24.41	30.28	5.01	8.77	13.31	16.43	20.08	4.87
T ₂	77.54	22.46	7.12	11.99	17.66	24.55	30.42	5.12	8.88	13.42	16.54	20.19	4.25
T ₃	78.87	21.13	7.86	12.73	18.40	25.29	31.16	5.76	9.52	14.06	17.18	20.83	3.96
T ₄	76.87	23.13	6.13	11.00	16.67	23.56	29.43	4.98	8.74	13.28	16.4	20.05	6.73
T ₅	74.56	25.44	6.78	11.65	17.32	24.21	30.08	4.67	8.43	12.97	16.09	19.74	6.89
T ₆	73.87	26.13	6.43	11.30	16.97	23.86	29.73	4.43	8.19	12.73	15.85	19.5	7.43
T ₇	72.79	27.21	5.67	10.54	16.21	23.10	28.97	4.21	7.97	12.51	15.63	19.28	7.56
T ₈	71.32	28.68	5.98	10.85	16.52	23.41	29.28	4.1	7.86	12.4	15.52	19.17	8.01
T ₉	69.57	30.43	5.36	10.23	15.90	22.79	28.66	3.98	7.74	12.28	15.4	19.05	8.56
T ₁₀	68.23	31.77	5.12	9.99	15.66	22.55	28.42	3.76	5.45	9.99	13.11	16.76	9.89
F Test	S	S	S	S	S	S	S	S	S	S	S	S	S
CD _{@5%}	2.543	1.685	0.897	0.88	0.975	0.93	0.893	0.453	0.651	0.569	0.468	0.765	0.764
S.Ed. (±)	1.231	0.814	0.416	0.431	0.462	0.451	0.398	0.231	0.301	0.249	0.221	0.351	0.352

Table 2 Influence of different planting date on Survivality %, Mortality %, Plant height, Number of leaves and Days to break of Jamun.

Symbol	Plant spread (cm)					Leaf area (cm ²)					No. of branches	Leaf area Index	Leaf area index
	Initial	30 DAT	60 DAT	90 DAT	120 DAT	Initial	30 DAT	60 DAT	90 DAT	120 DAT			
T ₁	7.12	10.33	13.45	18.01	20.99	3.54	4.88	8.22	10.09	12.13	9.01	2.54	59.65
T ₂	7.45	10.66	13.78	18.34	21.32	3.76	5.10	8.44	10.31	12.35	9.54	2.76	61.45
T ₃	7.86	11.07	14.19	18.75	21.73	3.98	5.32	8.66	10.53	12.57	10.76	2.89	65.87
T ₄	7.01	10.22	13.34	17.9	20.88	3.45	4.79	8.13	10.00	12.04	8.43	2.39	55.39
T ₅	6.98	10.19	13.31	17.87	20.85	3.39	4.73	8.07	9.94	11.98	7.98	2.21	53.36
T ₆	6.12	9.33	12.45	17.01	19.99	3.31	4.65	6.21	8.08	10.12	6.78	2.13	48.37
T ₇	5.78	8.99	12.11	16.67	19.65	3.1	4.44	6.00	7.87	9.91	5.87	2.01	46.29
T ₈	5.15	8.36	11.48	16.04	19.02	2.98	4.32	5.88	7.75	9.79	5.07	1.98	47.82
T ₉	4.49	7.7	10.82	15.38	18.36	2.76	4.10	5.66	7.53	9.57	4.65	1.81	49.35
T ₁₀	4.21	7.42	10.54	15.1	18.08	2.48	3.82	5.38	7.25	9.29	4.23	1.67	45.62
F Test	S	S	S	S	S	S	S	S	S	S	S	S	S
CD _{@5%}	0.463	0.352	0.875	0.859	0.981	0.078	0.098	0.324	0.431	0.738	0.984	0.0541	2.315
S.Ed. (±)	0.231	0.176	0.4375	0.4295	0.4905	0.039	0.049	0.162	0.2155	0.369	0.437	0.03	1.856

References

- Ara N, Moniruzzaman M, Rahman KS. Performance of hybrid lines of pointed gourd (*Trichosanthes dioica* Roxb) for yield and yield attributes. *Bangladesh Journal Agricultural Research*. 2018;43(3):383-393.
- Chavan et al. (2015)**, Variability And Character Association Studies In Ridge Gourd (*Luffa Acutangula* Roxb.) With Reference To Yield Attributes (*Journal of Global Biosciences* ISSN 2320-1355)
- Choudhary et al. (2011)**, Genetic divergence in hermaphrodite ridge gourd (*Luffa acutangula*) (Vegetable Science (2011) 38(1): 68-72)
- Gowtham KNP, Kuppastij, Mankani KL. A review on *Luffa acutangula*. *International Journal of Pharma World Research*. 2012;3(1):1-15.
- Haque MM, Uddin MS, Mehraj H, Uddin JAFM. Evaluation of snake gourd (*Trichosanthes anguina* L.) test hybrids comparing with four popular checks. *International Journal of Applied Science Biotechnology*. 2014;2(4):525-528.
- Hussain J, Rehman N, Khan AL, Hamayun M, Hussain SM, Shinwari ZK. Proximate and essential nutrients evaluation of selected vegetables species from Kohat region, Pakistan. *Pakistan Journal of Botany*. 2010;42(4):2847-2855.
- Kandlakunta B, Rajendran A, Thingnganing L. Carotene content of some common (cereals, pulses, vegetables, spices and condiments) and unconventional sources of plant origin. *Food Chemistry*. 2008;106:85-89.
- Narasannavar et al. (2014)**, Heterosis studies in ridge gourd [*Luffa acutangula* (L.) (Heterosis studies in ridge gourd [*Luffa acutangula* (L.)])
- Quamruzzaman AKM, Rahman MM, Akter L. Performance of bottle gourd lines in Bangladesh Condition. *Annals of Biological Sciences*. 2017;5(1):5-7.
- Rabbani et al. (2012)**, Variability, Character Association And Diversity Analysis Of Ridge Gourd (*Luffa Acutangula* Roxb.) Genotypes (SAARC J. Agri., 10(2): 01-10 (2012))
- Ramya B, Kerketta A, Topno SE. Evaluation of different hybrids for growth and yield attributes of bitter gourd (*Momordica charantia* L.) in Prayagraj Region. *International Journal of Current Microbiology and Applied Sciences*. 2020;9(12):1008-1012.
- Rathore JS, Collis JP, Singh G, Singh KR, Jat BL. Studies on genetic variability in ridge gourd (*Luffa acutangula* L. (Roxb.)) Genotypes in Allahabad Agro-Climate Condition. *International Journal of Current Microbiology and Applied Sciences*. 2017;6(2):317-338.
- Saensuket et al. (2022)**, A SNP of *betaine aldehyde dehydrogenase (BADH)* enhances an aroma (2-acetyl-1-pyrroline) in sponge gourd (*Luffa cylindrica*) and ridge gourd (*Luffa acutangula*) (Scientific Reports | (2022) 12:3718)
- Sharmin et al. (2013)**, Hypoglycemic and Hypolipidemic Effects of Cucumber, White

Pumpkin and Ridge Gourd in Alloxan
Induced Diabetic Rats (J. Sci. Res. 5 (1),
161-170 (2013).

UNDER PEER REVIEW