

KNOWLEDGE, ATTITUDE AND USAGE OF FEMALE ORAL CONTRACEPTIVE PILLS AMONG FEMALE HAWKERS IN THE CENTRAL BUSINESS DISTRICT, KUMASI, GHANA

ABSTRACT

Background: The use of oral contraceptive pills (OCPs) can prevent pregnancy and improve menstrual health. Their effectiveness, ease of use, and reversibility make them a preferred choice. Accessible birth control options for all individuals, regardless of age or finances, is crucial. Understanding cultural attitudes and usage patterns of contraceptives is vital for effective reproductive health care, including emergency contraception options.

Aims: This study evaluated the awareness, perceptions, and usage of oral contraceptives among women (female hawkers) in the central business district of the Kumasi Metropolis.

Study design: Cross-sectional study.

Place and Duration of Study: Central Business District (CBD) of Kumasi Metropolis encompassing the following areas: Adum Shopping Centre, Kejetia lorry station and Central market between the periods of 15th December 2023 and 12th April 2024.

Methodology: This study employed a descriptive cross-sectional design to evaluate knowledge, attitude and the use of the common female oral contraceptive pills among female street hawkers in Kumasi's CBD. A semi-structured face-to-face interview technique was utilized. Data was analyzed using Statistical Package for Social Sciences (SPSS) version 20 and Microsoft Excel software (2016).

Results: The study surveyed 114 participants, with (91.67 %) below 40 years of age. Most were single (66%), and 30% were married. Over half (54.7%) had used oral contraceptives, while 41.3% had not. Among users, 79% followed the instructions for use as directed by a healthcare professional, but 21% did not. A third (33.3%) reported contraceptive failure, whereas two-thirds (66.7%) did not. A majority (85.2%) believed in a link between contraceptive use and infertility. Usagetime of OCP varied as 28.8% used before, 5% during, and 65% after intercourse.

Conclusion: Majority of the respondents were found not to choose oral contraception as their means of contraception and those who chose it had limited knowledge about its usage and thus influenced compliance and adherence greatly.

Keywords: Contraception, Female hawkers, Unplanned births, Population fertility, Family planning

1. INTRODUCTION

Unintended pregnancies, which are either mistimed or unwanted, are a key factor in understanding population fertility and the unmet need for family planning and contraception. These pregnancies often result from the nonuse, inconsistent, or incorrect use of effective contraceptive methods. In 2012, it was estimated that globally, around 40% of all pregnancies were unintended, with higher rates observed in developing regions such as sub-Saharan Africa (UNFPA, 2022a; WHO, 2019). Specifically, in Ghana, approximately 37% of all pregnancies fall into this category (Ameyaw et al., 2019; Nsubuga et al., 2016).

The repercussions of unintended pregnancies are profound, leading to thousands of abortions and over 300,000 unplanned births in Ghana alone (Rossen et al., 2023; WHO, 2019). Contributing factors include poverty, societal stigma towards unmarried mothers, cultural preferences for male children, the burden of multiple roles on women, and limited access to family planning services (Ayalew et al., 2022; Ameyaw et al., 2019). These pregnancies can have detrimental effects on both maternal and child health, impacting prenatal care, physical health, labor experiences, and psychological well-being postpartum. Moreover, they can lead to substantial economic burdens for individuals, families, and national healthcare systems (Mudiyanselage et al., 2024).

Comprehensive strategies are essential for preventing unintended pregnancies and reducing their negative consequences. These strategies could include accessible contraceptive services, such as emergency contraception and safe, legal abortion options (Klima, 1998). The main objective is empowering women to make informed choices about their reproductive health, without being limited by age or financial capacity. A deep understanding of the public's perception and use of contraceptives is necessary to achieve this goal (Yalew et al., 2023; Stevens et al., 2016).

Additionally, the exploration of plant-based contraceptives presents an opportunity to develop more affordable and potentially safer alternatives to existing contraceptive methods. Research into the efficacy and safety of traditional herbal contraceptives, as well as their regulatory aspects, is ongoing (Upadhyay, 2024). Studies have highlighted the potential of certain plant-derived compounds to serve as effective contraceptives, offering a more accessible option, particularly for populations in rural areas (Upadhyay, 2024; Pradhan et al., 2013).

The world's population is surging, underscoring the need for effective birth control measures and proper fertility management through contraception. Despite the availability of numerous contraceptive options, the rate of unintended pregnancies, which contribute to higher antenatal and postnatal mortality rates, continues to rise (Bongaarts, 2020). Given these circumstances, it's crucial to investigate how contraceptives are perceived and utilized, particularly in African communities where the rates of unplanned births are on the rise.

Ultimately, tackling the challenge of unintended pregnancies demands a comprehensive strategy. This strategy should encompass educational initiatives, enhanced availability of family planning resources, and the creation of new contraceptive methods that are not only efficient but also align with cultural norms (UNFPA, 2022b).

The study investigated the knowledge perception and usage of female oral contraceptives among female hawkers in the central business district (CBD) of the Kumasi metropolis. The main objectives of this study were to assess the common contraceptives of misuse, reasons for use or misuse and knowledge of adverse effects associated with the use of contraceptives.

2. MATERIAL AND METHODS

2.1 Study Design

This study employed a descriptive cross-sectional design to evaluate the knowledge attitudes and usage of the common female oral contraceptive pills among female street hawkers in the Kumasi metropolis of Ghana.

2.2 Study site

The research was conducted in the Central Business District (CBD) Kumasi Metropolis. This area includes the Central Market, Adum Shopping Centre, and the Kejetia Lorry Park. These locations are all found in Adum, a suburb of Kumasi nestled between Bantama and Nhyiaeso. The Kumasi Central Market, founded in 1925, holds the distinction of being the largest traditional market in West Africa. It boasts a daily footfall estimated to be between 15,000 and 20,000 people (Kanwetuu et al., 2023). The market is composed of 20 % tabletop stores and 80 % brick-and-mortar stores, offering a total storage capacity of 25,000. Kejetia, on the other hand, is primarily a commercial hub with 60% of its area dedicated to vehicle parking facilities. The remaining space is divided between stores (30%), offices (5%), and residential areas (5%) (Lambongang, 2023).

2.3 Sampling technique and target population

The snowball sampling method was the primary selection technique for participants in this research. This method starts with a single respondent who then identifies others with similar traits to aid the researcher (Parbhoo-Ebrahim and Fourie, 2020). Due to the transient nature of female hawkers in Adum and Central Market, locating them for the study posed a challenge. However, snowball sampling was effective in reaching this itinerant group.

The process began with a few female hawkers who consented to join the study and were then asked to help locate additional interviewees. These initial participants were discovered by chance during a visit to the potential study locations. The study ultimately involved interviews with 114 participants. When no new insights were collected after 114 interviews, it was presumed that data saturation had occurred, indicating a theoretically adequate sample size.

Participants were selected through a combination of deliberate and random methods to avoid selection bias, and precautions were taken to guarantee that each participant was included in the study only a single time.

2.4 Data collection and technique

The primary data for this study was sourced from female head porters, both adults and youth, in the central business district of the Kumasi, Metropolis. A semi-structured face-to-face interview technique was utilized, with an interview guide that allowed the researcher to probe participants' responses when necessary (Creswell, 2007). The interview guide included questions about the socio-demographic profiles of potential research participants, their knowledge, perception and usage of female contraceptives.

Participants were asked questions in local languages because they could not speak the English language. An individual translator was therefore hired to assist, as the researchers were not familiar with other languages or local dialects. An audio tape recorder was used to record the conversations with research participants after obtaining oral consent. The interviews were conducted at the convenience of the research participants at scheduled times and venues to avoid interfering with their trade.

Using a semi-structured interview approach, probing and follow-up questions were major characteristics of the interview to elicit rich data. Regarding trustworthiness, which Tracy (2010) expresses as qualitative research credibility, the study location, the participants, and even the interview questions were not based on the convenience and subjective opinions of the researchers. The study location and scope were determined by literature and the need to contribute to knowledge on female hawkers' experiences in Kumasi, Ghana. Respondent validation was done for every interview, whereby the researchers repeated the responses the way they understood it to the participants for the latter to confirm the validity of the information. In cases where respondents disagreed with the researcher's interpretation of the data, the former was made to re-present their response for confirmation.

2.5 Inclusion and exclusion criteria

All female hawkers 18 years and above who consented to participate in the study were included; however, those below 18 years as well as those who did not consent to participate were excluded.

2.6 Ethical considerations

During this research, all relevant ethical standards for conducting studies were strictly adhered. In addition to obtaining informed consent from the study participants, disclosures about the objectives and benefits of the study, as well as the confidentiality of their information, were communicated. All participants were volunteers, and all data collected was handled with utmost discretion.

2.7 Data organization, entry and analysis

A manual thematic framework was utilized to develop themes for this study. Once the raw data was collected from all respondents, it was transcribed verbatim by one of the researchers into a word document. The transcripts were then read and organized using the two research questions as broad organizing categories.

After this organization, the responses under each category were read again, and emerging ideas were represented by simple words/phrases (codes). Similar codes identified across all transcripts were combined to form broad themes, which were primarily data-driven. All the researchers then engaged in drawing meanings, discussion, and presentation of the raw data.

Comment [MG1]: This information is not necessary. If someone does not consent to participate then he/she is not a participant.

Comment [MG2]: What are the two research questions?

Comment [MG3]: What bibliographical reference did you use for this analysis?

3. RESULTS

3.1 Sociodemographic Characteristics of female hawkers

Table 1. displays the sociodemographic details of the participants. The average age of the 114 food vendors is 27.13 years and 91.7% were under the age of 40. A significant portion, 54.9%, are Akan, and 68.75% are unmarried. The majority, 79.2%, identify as Christians, and 20.8% have not received formal education.

Table 1: Sociodemographic characteristics of respondents

Variable	Frequency (n=114)	Percentage (%)
Age		
<40	132	91.67
40+	12	8.33
Educational level		
No formal education	30	20.80
Primary	36	25.00
Junior High school	42	29.20
Senior High school	36	25.00
Tertiary	-	-
Marital status		
Married	45	31.25
Single	99	68.75
Religion		
Christian	114	79.17
Muslim	28	19.44
Traditionalist	2	1.39
Ethnicity		
Akan	79	54.90

Ewe	9	6.30
Mole-Dagbani	56	38.88

3.2 Respondents' knowledge attitude and usage of oral contraceptive pills.

Table 2 outlines the knowledge, attitude and usage of contraceptives among female street hawkers. A significant portion, **77.8 %**, acknowledged being aware of oral contraceptive pills. A large percentage, **45.53 %**, reported that their source of information were friends. Approximately **57 %** agreed to the use of OCP, and 80.49 % confirmed that they had experienced side effects. Additionally, **63.41 %** of the hawkers confirmed the use of OCP after sex as directed by the pharmacist or the pharmaceutical support staff (PSS).

Table 3: A table showing the respondents' usage of oral contraceptive pills.

Variable	Frequency (n)	Percentage (%)
Have you heard of oral contraceptives?		
Yes	112	77.78
No	32	22.22
Source of knowledge		
Friends	51	45.53
TV/Radio advertisement	38	33.93
Salespersons	13	11.61
Others	10	8.93
Have you used oral contraceptives pills		
Yes	82	56.94
No	62	43.06
Reasons for not using OCPs		
Free period method	38	61.29
Virgin	19	30.65
Other methods	5	8.06

Preferred brand of OCPs		
Postinor 2	24	29.27
Lydia	38	46.34
Levon 2	13	15.85
Others	7	8.54
Mode of usage of OCPs		
Before sexual intercourse	25	30.49
During sexual intercourse	4	4.88
After sexual intercourse	52	63.41
others	1	1.22
Was the preferred OCP effective?		
Yes	28	34.15
No	54	65.85
Any side effects encountered?		
Yes	66	80.49
No	16	19.51
Side effects encountered		
Irregular menstruation	21	31.82
Profuse bleeding	15	22.73
Menorrhagia	24	36.36
others	6	9.09
Relationship between contraceptive use and infertility		
Yes	69	84.15
No	13	15.85

Comment [MG4]: On a daily basis? Consistent use?

Comment [MG5]: I didn't understand. Is it literally? What do you mean?

Comment [MG6]: morning-after pill?

4. DISCUSSION

The use of oral contraceptive pills has been the major means of preventing unwanted pregnancy among most women of reproductive age. While oral contraceptive pills offer significant contraceptive benefits that are important to many women, inconsistent use of these pills results in significant failure rate that are at least 20 times higher than failure rates of IUDs and implants (Yeboah et al., 2022). However, if taken correctly, the oral contraceptive pill can prove to be up to 99% effective (Cameron et al., 2020). Even though the oral contraceptive pill is one of the easiest and safest means of contraception, perceptions and discernments surrounding its usage are immeasurable varying from race, religion, culture and various traditions. While the primary usage of oral contraceptive pills is aimed at preventing contraception, it is fascinating to realize that, it has a range of other benefits other than their primary purpose. Contraception reduces pregnancy-related morbidity and mortality, reduces the risk of developing certain reproductive cancers and can be used to treat many menstrual related symptoms and disorders (Steller et al., 2021). The findings revealed that the majority of the participants had heard about OCP. This finding is similar to the study conducted by Yeboah et al., (2022) in Ghana, (Kgosiemang, 2011) in Botswana and Mishore et al., (2019) in Ethiopia. Additionally, Adama Town, Ethiopia (38.7%) (Woldemichael et al., 2023) showed a comparatively low level of OCP awareness among study participants which was contrary to this study's findings. The differences in the level of awareness could be due to differences in the study settings and time variation in relation to the acceleration of reproductive health promotion activities (Tessema and Bayu, 2015).

In this research, friends and media emerged as the primary sources of information regarding oral contraceptive pills (OCP) for most participants, as shown in Figure 1. This contrasts with other studies where health professionals were the main informants (Yeboah et al., 2022). Notably, a study by Mohammed et al., (2019) also found health professionals to be a key source. Concerning OCP, 63.41% of this study's participants were aware of the correct usage timing, a figure that is lower than Tamale's 85% but higher than Botswana's 38.2% Mohammed et al., (2019). However, the study in Nigeria (66.7%) showed similar proportions with the present study (Solanke, 2017). Additionally, 80.49% of participants in this study were knowledgeable about OCP side effects, This finding is similar to that of a study conducted in Nigeria which reported that 72.5% of the participants had correct knowledge on the side effects of OCPs but higher than that of the study in India (8.6%) (Relwani et al., 2012). Menorrhagia was identified as a common side effect by 36.36% of participants, whereas another study highlighted nausea and vomiting as more prevalent Sibanda et al., (2017). Research has indicated that poor publicity of contraceptives and limited availability are possible reasons for low awareness (Gosavi et al., 2016). Thus, the high level of awareness of OCP in the present study could be due to high publicity on social media including TV/radio broadcast.




Figure 1: a chart showing the source of knowledge of respondents on oral contraceptives

Comment [MG7]: Figure 1 should be in results section

Religious and cultural beliefs significantly influence the use of oral contraceptive pills. Healthcare advice that conflicts with a patient's religious beliefs may be rejected. For example, many Christians view procreation as a divine command and consider interference with childbirth, including contraception, as sinful (Ugwuewo and Sopuru, 2023). Consequently, Christian couples often rely on natural family planning, which can be unreliable due to various factors, leading to unintended pregnancies. In this study, out of the 144 participants, 31.3% were married, but only 5 continued to use oral contraceptives, none of whom were affiliated with any religion (Figure 2).

Many traditions and culture relate the use of contraceptive methods to infertility. It is believed by a great number of people that; prolonged and long-term use of oral contraceptives has the tendency of causing infertility or delays conception. This belief has left many women who wish to resort to contraception in a state of decisiveness. This belief has thus influenced the usage of oral contraceptives negatively as most women who wish to bear children in the future find it difficult to use contraceptive methods. However, this belief is scientifically proven to be false. When it comes to birth control and fertility, there can be a lot of confusion. But hormonal contraceptives don't cause infertility, no matter which method used or how long it is used. What they're designed to do, however, is temporarily delay your fertility and prevent pregnancy.

From the results accumulated from this study, out of the 82 respondents who responded to this section, 85.2% believed that, the use of oral contraceptives can cause infertility. This may account for the reasons why the use of oral contraceptives is not patronized in the study area and the entire country in general (Figure 3).




Figure 2. A graph showing the relation between marital status, religion and the use of oral contraceptive pills among respondents

Comment [MG8]: Figures 2 and 3 should be in results section




Figure 3. A chart showing respondents' knowledge on the relationship between oral contraceptives and infertility

5. CONCLUSION

The study concludes that oral contraceptive pills offer numerous benefits, from birth control to other therapeutic advantages. However, their usage is not well-monitored. Many potential users lack the necessary in-depth knowledge to fully benefit from these pills. Socioeconomic factors such as religion and culture greatly influence a person's choice of contraceptive method. Despite this, some individuals disregard their religious beliefs and choose oral contraceptive pills, often facing stigmatization and judgment. These factors have become barriers to the use of oral contraceptive pills among the study's respondents, primarily street hawkers in the Kumasi metropolis.

ETHICAL APPROVAL (WHEREEVER APPLICABLE)

This research received approval from the Committee on Human Research, Publications, and Ethics (CHRPE) at Royal Ann College of Health (RACOH, Kumasi), with the reference number CHRPE/EC/114/3.

UNDER PEER REVIEW

REFERENCES

- Ameyaw, K. E., Budu, E., Sambah, F., Baatiema, L., Appiah, F., Id, A. S. and Ahinkorah, B. O. (2019). Prevalence and Determinants of Unintended Pregnancy in Sub-Saharan Africa: A Multi-country Analysis of Demographic and Health Surveys. *PLoS ONE*, 14(8):1–16.
- Ayalew, H. G., Liyew, A. M., Tessema, Z. T., Worku, M. G., Tesema, G. A., Alamneh, T. S., Teshale, A. B., Yeshaw, Y. and Alem, A. Z. (2022). Prevalence and factors associated with unintended pregnancy among adolescent girls and young women in sub-Saharan Africa, a multilevel analysis. *BMC Women's Health*, 22(1):1–9. Retrieved from <https://doi.org/10.1186/s12905-022-02048-7>
- Bongaarts, J. (2020). Trends in fertility and fertility preferences in sub-Saharan Africa: the roles of education and family planning programs. *Genus*, 76(1).
- Cameron, S. T., Glasier, A., McDaid, L., Radley, A., Baraitser, P., Stephenson, J., Gilson, R., Battison, C., Cowle, K., Forrest, M., Goulao, B., Johnstone, A., Morelli, A., Patterson, S., McDonald, A., Vadiveloo, T. and Norrie, J. (2020). Use of effective contraception following provision of the progestogen-only pill for women presenting to community pharmacies for emergency contraception (Bridge-It): a pragmatic cluster-randomised crossover trial. *The Lancet*, 396(10262):1585–1594. Retrieved from [http://dx.doi.org/10.1016/S0140-6736\(20\)31785-2](http://dx.doi.org/10.1016/S0140-6736(20)31785-2)
- Gosavi, A., Wong, H. and Singh, K. (2016). Knowledge and factors determining choice of contraception among Singaporean women. , 57(11):610–615.
- Kanwetuu, D. V. P., Worae, J. and Acheampong, G. B. (2023). Coping with the challenges of head porterage in Ghana: The case of female head porters (kayayei) in Kumasi. *Journal for Social Thought*, 7(1). Retrieved from <http://ir.lib.uwo.ca/jst/>
- Kgosiemang, B. (2011). Emergency contraceptive knowledge , attitudes and practices among female students at the University of Botswana : A descriptive survey. :1–6.
- Klima, C. S. (1998). Unintended pregnancy consequences and solutions for a worldwide problem. *Journal of Nurse-Midwifery*, 43(6):483–491.
- Lambongang, J. M. (2023). Investigating Challenges of Mobile Money usage in the Central Business District of the Kumasi Metropolitan Assembly, Adum- Ghana. *Texila International Journal of Management*, 9(1):42–57.
- Mishore, K. M., Woldemariam, A. D. and Huluka, S. A. (2019). Emergency Contraceptives : Knowledge and Practice towards Its Use among Ethiopian Female College Graduating Students. , 2019:14–16.
- Mohammed, S., Abdulai, A. and Iddrisu, O. A. (2019). Pre-service knowledge , perception , and use of emergency contraception among future healthcare providers in northern Ghana. , 9:1–7.
- Mudiyansele, S. B., Dona, S. W. A., Angeles, M. R., Majmudar, I., Marembo, M., Tan, E. J., Price, A., Watts, J. J., Gold, L. and Abimanyi-Ochom, J. (2024). The impact of maternal health on child's health outcomes during the first five years of child's life in countries with health systems similar to Australia: A systematic review. *PLoS ONE*, 19(3 March):1–19. Retrieved from <http://dx.doi.org/10.1371/journal.pone.0295295>
- Nsubuga, H., Sekandi, J. N., Sempeera, H. and Makumbi, F. E. (2016). Contraceptive use, knowledge, attitude, perceptions and sexual behavior among female University students in Uganda: A cross-sectional survey. *BMC Women's Health*, 16(1):1–11. Retrieved from <http://dx.doi.org/10.1186/s12905-016-0286-6>
- Parbhoo-Ebrahim, N. and Fourie, I. (2020). Pathways to research participant recruitment in a challenging information behaviour context: South African cold case investigators as exemplar, in: *Pathways to Research*.
- Pradhan, D. K., Mishra, M. R., Mishra, A., Panda, A. K., Behera, R. K., Jha, S. and Choudhury, S. (2013). A comprehensive review of plants used as contraceptives. *International Journal of Pharmaceutical Sciences and Research*, 4(1):148–155.

- Retrieved from
<http://www.embase.com/search/results?subaction=viewrecord&from=export&id=L368388155%5Cnhttp://www.ijpsr.com/V4I1/15 Vol. 4, Issue 1, January 2013, IJPSR, RE 799, Paper 15.pdf%5Cnhttp://vb3lk7eb4t.search.serialssolutions.com?sid=EMBASE&issn=23205148&id=doi>
- Relwani, N., Saoji, A., Kasturwar, N. B., Nayse, J., Junaid, M. and Dhatrak, P. (2012). EMERGENCY CONTRACEPTION : EXPLORING THE KNOWLEDGE , ATTITUDE AND PRACTICES OF ENGINEERING COLLEGE GIRLS IN NAGPUR DISTRICT. , 3(1):3–8.
- Rossen, L. M., Hamilton, B. E., Abma, J. C., Gregory, E. C. W., Beresovsky, V., Resendez, A. V., Chandra, A. and Martin, J. A. (2023). Updated Methodology to Estimate Overall and Unintended Pregnancy Rates in the United States: Data Evaluation and Methods Research.
- Sibanda, E., Chb, M. B., Med, D. F., Obst, D., Sa, F., Titus, M. J., Chb, M. B., Sa, F., Dip, P. G. and Res, I. (2017). Knowledge , attitudes and practices of health professionals in public health institutions on emergency contraception in Pietermaritzburg , KwaZulu-Natal Province , South Africa. , 23(1):7–11.
- Solanke, B. L. (2017). Factors influencing contraceptive use and non-use among women of advanced reproductive age in Nigeria. *Journal of Health, Population and Nutrition*:1–14. Retrieved from <http://dx.doi.org/10.1186/s41043-016-0077-6>
- Stevens, G. A., Finucane, M. M. and Paciorek, C. J. (2016). Levels and Trends in Low Height-for-Age. *Disease Control Priorities, Third Edition (Volume 2): Reproductive, Maternal, Newborn, and Child Health*:85–93.
- Tessema, B. M. and Bayu, H. (2015). Knowledge, Attitude and Practice on Emergency Contraception and Associated Factors among Female Students of Debre-Markos University, Debre-Markos Town, East Gojam Zone, North West Ethiopia, 2013. , 15(1).
- Ugwuewo, C. S. and Sopuru, C. S. (2023). UNDERSTANDING BIRTH CONTROL IN THE CONTEXT OF GENESIS 38 : 8-10. *Global Scientific and Academic Research Journal of Multidisciplinary Studies*, 2(4):06–12.
- UNFPA. (2022a). Nearly half of all pregnancies are unintended—a global crisis, says new UNFPA report. Retrieved May 17, 2024, from <https://www.unfpa.org/press/nearly-half-all-pregnancies-are-unintended-global-crisis-says-new-unfpa-report>
- UNFPA. (2022b). UNFPA Strategy for Family Planning, 2022-2030: Expanding Choices – Ensuring Rights in a Diverse and Changing World. *Laser Focus World*, 39(11):60.
- Upadhyay, R. K. (2024). Plant origin contraceptives: phytochemistry, mechanism of action, and side effects. *International Journal of Green Pharmacy*, 18(1).
- WHO. (2019). Emergency contraception. Retrieved May 17, 2024, from <https://www.who.int/news-room/fact-sheets/detail/emergency-contraception>
- Woldemichael, D., Agero, G., Jima, A. and Woldemichael, B. (2023). Postabortion Family Planning Utilization and Associated Factors among Women Seeking Abortion Services : A Cross-sectional Study Dereje Woldemichael , Gebi Agero , Aman Jima and Bedasa Woldemichael Abstract Ethiopia is a country which suffers with one of t. *Canadian Journal of Family and Youth*, 15(3):157–171.
- Yalew, A. Z., Olayemi, O. O. and Yalew, A. W. (2023). Reasons and prevention strategies of unintended pregnancy in Addis Ababa, Ethiopia: A phenomenological qualitative study. *BMJ Open*, 13(7):1–9.
- Yeboah, D. S., Afranie, M., Id, A. and Kampitib, G. B. (2022). Factors influencing the use of emergency contraceptives among reproductive age women in the Kwadaso Municipality , Ghana. , (Ci):1–17. Retrieved from <http://dx.doi.org/10.1371/journal.pone.0264619>