

**Use of organic extracts and fertilizers forenhancement of productivity in greengram (*Vigna radiata* L.)**

---

**ABSTRACT**

Greengram (*Vigna radiata* L.) being a leguminous crop has high nutritional value and is one of the most important pulse crops grown in India. In India greengram production is affected by a variety of factors, including biotic stresses like diseases, weeds, and pests as well as abiotic stresses such as drought, floods and poor management practices. Among those the use of inorganic fertilizers has resulted in ground water pollution, deteriorating soil quality, high production cost and poor quality production for livelihood which has resulted in poor productivity. Thus the main objective of this paper is to review the use of organic extracts and fertilizers and their benefits for enhanced productivity in greengram. As organic sources are known to considerably effect the growth and development of a crop yielding significant results, as a result utilization of such organic sources like seaweed extract, vermiwash, humic acid, panchagavya and jeevamruth can play a major role in enhancing the productivity in greengram.

*Keywords: Greengram, Organic fertilizer, Productivity*

---

## 1. INTRODUCTION

In India, pulses are consumed more often than any other protein source, highlighting their crucial role in the daily eating habits of the people. But the rising demand for pulses in India has outpaced domestic production, necessitating the need for imports [1]. In recent decade pulses saw a growth rate that was 32% less compared to cereals due to farmer's having major dependence on cultivation of cereal crops over pulses. This is due to pulses having low yielding capacity and poor resistant to diseases which is a major limitation in adoption of pulses by farmer's [2]. In India, the cultivation of pulses covers an area of about 23.55 million hectares, leading to an annual production of approximately 17.15 million tonnes, with an average yield of 728 kilograms per hectare [3]. The productivity of pulses in India is considerably low as compared to other major countries thus improved management practices are in need of hour. [1,3].

Greengram scientifically known as *Vigna radiata* (L.) is a staple food rich in protein. It comprises approximately 25% protein, nearly triple the protein content found in cereals. It is an ancient and well-established leguminous crop in Asia, favored for its successful growth during the kharif season. It can be cultivated either as a sole crop, intercropped, or as a fallow crop. It plays a significant role in Indian agriculture because of its capacity of fixing atmospheric nitrogen via its nodules which helps in improving soil fertility [4]. In India, it covers an area of 64.56 lakh hectares with a production of 4.05 million tones [5]. In India, it accounts for nearly eight percent of the total pulse area and ranks second based on area and production among other pulse crops. The seeds contain 24.7% protein, providing a cost-effective protein source, which is why it is known as "poor man's meat." Green gram is highly digestible and palatable, with its pods commonly used as a green vegetable. Both its whole and split grains are utilized in dishes like dal and curry. Due to its high digestibility, its curry is often recommended for patients [6].

Organic fertilizer plays a crucial role in enhancing soil fertility and increasing crop production but farmers typically overlook its utilization in agricultural practices. The production of greengram in India is considerably low due to factors like poor soil fertility which results from cultivation practices without adequate soil fertility management. This is due to the result of extensive use of synthetic chemical fertilizers which poses a serious threat in maintaining sustainable production. The prolonged and indiscriminate use of chemical fertilizers leads to a serious issue concerning soil health, particularly in terms of reduced efficiency in input utilization, and particularly fertilizer use efficiency [7].

Crop residue and other organic residues are valuable to small-scale farmers because they are cost effective, readily accessible source of nutrients, and environmental friendly as compared to inorganic fertilizers [8]. The use of organic manures, either on their own or in combination with liquid fertilizers, aids in enhancing the physical and chemical properties of the soil, as well as optimizing the effective utilization of applied organic fertilizers to improve seed yield and quality [9]. Enhancing soil quality involves increasing soil organic carbon, cation exchange capacity, soil water content, and promoting beneficial soil microorganisms through the addition of organic matter to the soil. Moreover, organic fertilizer is often the key limiting factor for crop production in major agricultural regions globally, and thus, embracing effective organic fertilizer strategies frequently yields substantial economic benefits for farmers [10].

Organic substances like panchagavya and jeevamrutha are important for boosting plant growth and immunity, as they provide essential nutrients, growth-promoting hormones, and beneficial microorganisms including lactic acid bacteria, yeast, actinomycetes, and photosynthetic bacteria. These microorganisms, coupled with well-known fertilizers like azotobacter, azospirillum, and phosphobacterium, also aid in improving soil health, fostering crop growth, and boost yield [11]. Seaweed extract an organic extract exhibits phyto-stimulatory properties that leads to the enhancement of growth, health, yield, and quality in a wide range of important agricultural and horticultural crops. Application of seaweed extract stimulates plant growth by boosting seedling growth, root development, and the absorption of minerals and nutrients through the activation of physiological and molecular mechanisms [12]. Organic fertilizers contain microorganisms that break down organic matter, releasing nutrients into the soil. These microbes obtain their energy from decomposing plant and animal residues, thus providing the soil with a diverse range of nutrients [13]. Restoring soil fertility necessitates nourishing the soil microorganisms, which can be

accomplished through the addition of organic material, minimizing tillage, and eliminating synthetic fertilizers and chemicals [14].

The utilization of liquid organic formulations, as opposed to synthetic or inorganic fertilizers, is acknowledged for its capacity to enhance plant growth and productivity by delivering essential plant nutrients, preserving soil biodiversity, and mitigating environmental concerns, ultimately fostering soil health [15]. Earlier reports indicate that due to rising fertilizer costs and scarcity, as well as mounting environmental concerns and government support for organic farming, we have been prompted to explore alternative methods of nutrient application such as panchgavya, vermiwash and cow urine [16]. In order to achieve sustainable soil fertility and crop productivity, the utilization of organic manures and other nutrient management practices, including fermented liquid organic fertilizers like Panchagavya, Jeevamrut, Beejamrut, Sasyamrut, Gomutra, Angara, Vermiwash, is essential. These fermented liquid organic fertilizers not only provide nutrients but also contain microbial populations and plant growth-promoting substances (PGPR), which contribute to stimulating plant growth, increasing yield, enhancing metabolic activity, and boosting resistance to pests and diseases [17]. Hence the objective of this paper is to review the use of organic extracts and fertilizers for enhancement of productivity in greengram.

## **2. ECOLOGICAL REQUIREMENT OF GREENGRAM**

Various abiotic and biotic factors, including temperature, soil moisture, soil acidity, salinity, drought, waterlogging, pest, and disease pressure, can reduce greengram productivity by altering physiological processes and plant water relations [18,19,20,21,22,23]. The crop has the ability to thrive and produce in regions experiencing lower ranges of rainfall, although its optimal growth and productivity depend on receiving well-distributed precipitation ranging between 350 and 700 millimeters per year [24]. As a short-day plant with quantitative characteristics, the green gram crop exhibits sensitivity to changes in day length [25]. Short days result in early flowering without adequate vegetative biomass while long days promote delayed flowering. While photoperiod responses may differ among various varieties, the crop tends to thrive optimally when temperatures range between 25°C and 35°C [26]. The green gram crop can grow well in a broad spectrum of soils, including those with red laterite, black cotton, and sandy textures. However, it shows optimal growth in deep, fertile, well-drained loamy to sandy loam soils with a pH level ranging between 6.0 and 7.0.

## **3. SOIL FERTILITY STATUS OF INDIA**

Evaluating soil characteristics to ascertain its fertility status within a particular area or region is essential for advancing sustainable agricultural production, given the significant influence of soil fertility on crop yields. In recent years, intensive agriculture has witnessed a notable decline in the efficiency of chemical fertilizer nutrients, attributed to imbalanced and inadequate fertilizer application as well as the low efficiency of other inputs [27]. The extensive barren stretch of the central plain in Uttar Pradesh poses a significant barrier to agricultural productivity and efforts are aimed at restoring the land [28]. In Kapurthala district of Punjab the majority of farmers engage in three or more crop rotations per year and utilize large quantities of fertilizers, insecticides, and pesticides to maximize their gross returns, typically without considering the soil's fertility status [29]. Farmers in the Pilibhit district of Uttar Pradesh heavily utilize urea chemical fertilizer in their fields, causing an elevation in nitrogen levels within the soil system while maintaining consistent levels of other macronutrients, resulting in a change in soil texture. Moreover, farmers employ various chemicals such as pesticides, insecticides, herbicides, and mixtures like calcium oxide and sodium chloride, toxic substances like Sulfas pills, kerosene oil, and tobacco leaves, to eradicate insect pests and weeds from both their agricultural soil and standing crops [30].

#### 4. GREENGRAM PRODUCTION CONSTRAINT IN INDIA

Among the various factors influencing greengram production and productivity, the utilization of varieties with low yield potential at the farmer level is noteworthy. Farmers encountered challenges associated to the use of improved or recommended green gram seeds, including issues like the non availability of required high-quality seeds, higher seed costs and a lack of information about recommended varieties among the farmers [31]. Biological challenges consist of the indeterminate growth pattern found in local cultivars like Jhain Mung, unproductive plant types, flower and pod shedding, low harvesting efficiency, and shattering tendencies which results in low productivity. The crop is primarily grown as a rain-fed crop in marginal and sub-marginal conditions, often without any nutrient management practices implemented. Typically, farmers refrain from utilizing any nutrient or biofertilizer applications, with YMV (Yellow Mosaic virus) serving as the predominant factor contributing to low yields [32].

#### 5. RESPONSE OF GREENGRAM TO VARIOUS ORGANIC FERTILIZERS

##### 5.1. HUMIC ACID

Humic acid (HA), a crucial ingredient in many organic fertilizers and a highly influential component of soil and organic matter, could potentially mitigate production limitations in pulse crops. Humic acid enhances plant growth and increases yield by improving nutrient absorption and influencing multiple physiological processes, such as cellular respiration, photosynthesis, protein synthesis, and enzymatic activity. It shows potential as a natural resource for substituting fertilizers to enhance crop productivity [33]. Humic acid distinguished by its deep black colour, is produced through the microbial decomposition of plant and animal matter and remains resilient to further weathering [34]. Humic acid comprises 51% to 57% carbon (C), 4% to 6% nitrogen (N), and 0.2% to 1% phosphorus (P), alongside minute quantities of other micronutrients [35]. Humic acid can be applied either to the soil or as a foliar spray. When used as a soil amendment, it promotes the formation of soft, aerated soil with added nutrients, while its application as a foliar spray enhances nutrient absorption, permeability, and photosynthetic rates [36].

##### 5.2. EFFECT OF HUMIC ACID ON CROP PRODUCTION

Humic acid regulates the release of nutrients such as nitrogen and phosphorus, potentially boosting growth and yield measures, thereby reducing the dependence on inorganic fertilizers for plant growth. According to Ali *et al.*, (2019) [37] application of humic acid at the rate of 12 kg ha<sup>-1</sup> followed by 9 kg ha<sup>-1</sup> resulted highest seed yield, number of seeds per pod, maximum pod length and number of pods per plant. The probable explanation for increased number of pods per plant and number of seeds per pod may be due to humic acid led to an enhanced reproductive efficiency in mungbean and the heightened presence of certain nutrients caused an increase in cell division, subsequently leading to higher grain yield. The increased seed weight of mungbean following soil application of HA reflects improvements in soil nutrient availability and its positive impact on plant growth [38]. Humic acid improves the absorption and accumulation of nutrients, particularly nitrogen, by facilitating the permeability of cell membranes [39,38]. Study conducted by Metre *et al.*, (2013) [40] mentioned that application of humic acid as foliar spray through cowdung at the rate of 350ppm resulted in increase seed yield, number of pods per plant and 100 seed weight. The increase in yield among the yield contributing parameters might be due to mineralization, including nutrient input, which resulted in enhanced growth and more efficient allocation of assimilates to different metabolic pathways.

### 5.3 SEAWEED EXTRACT

Seaweeds, the macroscopic marine algae, have multiple applications such as serving as human food, fodder for animals, replacing chemical fertilizers, and providing various useful chemicals. In recent years, natural seaweeds have been utilized as replacements for synthetic fertilizers. Seaweed extracts are promoted as liquid fertilizers and bio-stimulants owing to their inclusion of various growth regulators such as cytokinins, auxins, gibberellins and other macro and micronutrients necessary for crop growth and development [41]. Seaweeds contain the majority of the necessary trace elements and plant growth hormones crucial for facilitating plant growth and development. Studies have found that seaweed-based fertilizers are rich in potassium, nitrogen, and phosphorus. Seaweed liquid fertilizers are beneficial for enhancing agricultural productivity as their extract contains growth-promoting hormones (IAA and IBA), cytokinins, gibberellins, trace elements, vitamins, amino acids, antibiotics, and micronutrients [42]. Various seaweed species are divided into three categories based on the pigments they generate: Brown (Phaeophyta), Red (Rhodophyta), and Green (Chlorophyta) [43]. **Table 1** shows the chemical composition of various seaweed species [44].

**Table 1: Chemical composition of various seaweed species**

<b>Analysis</b>	<b><i>Ascophyllum Nodosum</i></b>	<b><i>Sargassum sp.</i></b>	<b><i>Palmaria palmata</i></b>	<b><i>Ulva sp.</i></b>
Crude Protein (%)	8±2.7(5)	8.5±1.8(10)	19.1±6.1(12)	18.6 ± 7.3 (14)
Crude fibre (%)	5.5(4.1-6.8)	10.1±2.4(9)	1.5	6.9 ± 4.1 (7)
Lignin (%)	13.8(6.2-21.4)	4.4(1.0-7.9)	NA	3.5 ± 2.1 (3)
Ash(%)	22.5±2.1(12)	35.9±12.8(9)	24.5(17.5- 31.5)	23 ± 7.4 (13)
Ca (g/kg)	20	3.8± 2.6(4)	NA	29.2 ± 28.9 (3)
P(g/kg)	1	2.2±1(5)	NA	2.7 ± 2.1 (3)
K(g/kg)	24	46.2(15.9-76.6)	NA	22.1 (15.1 -29.0)
Na(g/kg)	NA	-	NA	20.2 (11.0–29.3)
Mg(g/kg)	8	7.7(7.5-7.9)	NA	16.7 ± 3.2 (3)
Mn(mg/kg)	12±3(8)	214±106(4)	11	101
Zn(mg/kg)	181±114(8)	214±151(5)	143	45 (28–61)
Cu(mg/kg)	28±16(1)	7±4(5)	24	12 (7–17)
Fe(mg/kg)	134±36(8)	7291±6327(5)	153	1246 (1052–1440)

**Source:** Makkar *et al.*(2015)

Seaweed enhances the soil's ability to retain water and fosters a more balanced relationship between air and water within it. Liquid extracts from seaweed have been linked to numerous beneficial outcomes, such as boosted crop productivity, improved plant resistance to frost, increased absorption of soil nutrients, and heightened resilience to various environmental stresses [45]. Seaweeds are underutilized in Indian agriculture for specific reasons. Farming methods employed in coastal areas involve incorporating dried seaweed in powdered or flake form into the soil or using spray extracts derived from ground

seaweed [46]. Limited awareness among indigenous farmers regarding the accessibility of seaweeds contributes to their underutilization in Indian agriculture, while Indian farmers typically favor chemical fertilizers, displaying greater confidence in their efficacy. Furthermore, farmers residing in non-coastal or inland regions of India often lack awareness regarding the availability and significance of seaweeds. The lack of awareness among farmers hinders their purchasing of seaweed extracts from the market, posing a challenge for industries aiming to expand production. This obstacle is also influenced by the techno-economic evaluation of the seaweed industry, its products, and their various applications [43].

#### 5.4 EFFECT OF SEAWEED EXTRACT ON CROP PRODUCTION

Seaweed extracts contain key primary and secondary nutrients, amino acids, vitamins, and growth-promoting compounds like cytokinins, auxins, and abscisic acid. Their application has been observed to stimulate plant growth, increase yield, and enhance resistance to various environmental stresses [47]. As per study conducted by Iswarya *et al.*, [48] seeds soaked in a 0.1% seaweed extract solution for 30 minutes, and then applied as a foliar spray once at 25 days after sowing and at 35 days after sowing recorded highest seed yield ( $1242 \text{ kg ha}^{-1}$ ) over control treatment. It may be due to seaweed-based liquid fertilizers provide a range of essential plant nutrients and hormones that positively affect both the growth and physiological aspects of plants, ultimately leading to increased crop yields. Zodape *et al.*, (2010) [49] reported a similar trend of findings, as they observed a substantial boost in yield of 30.11%, 26.30%, and 16.78% respectively in greengram through foliar application of *Kappaphycus spp* extract, compared to the control group. Another study conducted by Chitra and Sreeja (2013) [50] observed *Gracilaria corticata* seaweed when applied on greengram at the rate of 4% showed superior performance in all aspects of growth, yield, and soil nutrient levels compared to the seaweed liquid fertilizer derived from the green algae *Caulerpa peltata*. This could be attributed to the presence of both macro and micronutrients, leading to an enhanced yield. An experiment conducted by Huda *et al.* (2022) [51] reported that foliar spray of seaweed (*Gracilaria tenuistipitata* var. *Liui*) at the rate of 20% recorded maximum number of branches per plant and stated that the seaweed extract demonstrated a 23.04% higher count of branches per plant when utilizing a 20% concentration in comparison to the recommended fertilizer doses. Whereas its application led to a 50% rise in the number of pods per mungbean plant compared to untreated conditions, 4.36% increase in seeds per pod and seed yield by 93.14% in comparison to control condition. The growth-regulating substances (seaweed oligosaccharides) prompted the synthesis of hormones such as abscisic acid, cytokinin, and auxin in treated plants, which can enhance crop growth whereas the increase in crop yield might also be linked to specific nutritional elements, particularly iron, zinc, and manganese from compost, as well as potassium, calcium, magnesium, sulfur, and iron from the seaweed extracts we employed. This correlation is supported by Zodape *et al.* (2009) [52], who noted a boost in green gram yield with the application of *Gracilaria* extracts. Seaweed extracts (SWEs) offer a superior alternative by minimizing the reliance on synthetic fertilizers and agrochemicals in agricultural systems. Additionally, seaweed extracts (SWEs) improve plant health by affecting the microbial community in the rhizosphere, making them a feasible eco-friendly method for sustainable agriculture and a potential component in organic cultivation of agricultural crops [12,50].

#### 5.5 VERMIWASH

Vermiwash refers to a liquid obtained by filtering water through a bed of earthworms, comprising their excretions, mucus, and micronutrients derived from organic matter in the soil. It serves as an effective foliar spray. Vermiwash are enriched with enzymes and secretions from earthworms has the capacity to stimulate crop growth and enhance yields [53]. It comprises a mixture of different enzymes, including protease, amylase, urease, and phosphatase. Moreover, analysis of vermiwash's microbial content has revealed the presence of nitrogen-fixing bacteria like *Azotobacter sp.*, *Agrobacterium sp.*, and *Rhizobium sp.*, as well as phosphate-solubilizing bacteria [54]. The application of vermiwash as a foliar spray notably enhanced the growth, yield, and quality of the crops [55]. The quality of vermiwash generated by earthworms is impacted by the composition of the raw materials used in vermicompost preparation [56].

Vermiwash consists of 0.01-0.001% Nitrogen, 1.70% Phosphorous, 26ppm Potassium, 8ppm sodium, 3ppm calcium, 0.01 ppm Copper, 0.06 ppm Iron, 160 ppm Magnesium, 0.60 ppm Manganese and 0.02ppm Zinc. When combined with a 10% solution of cow urine or extracts from neem or garlic and sprayed, it acts as a biopesticide[57].The vermiwash obtained from municipal solid waste contains organic materials, plant nutrients, and soluble salts, all of which contribute to increasing soil nutrient levels and moisture retention [58].

## 5.6 EFFECT OF VERMIWASH ON CROP PRODUCTION

Vermiwash assists in effectively managing market wastes for profit and can be utilized for crop production through foliar and soil applications. At present, foliar application is the prevailing method for sustainable crop production, aiming to meet production goals while minimizing the use of fertilizers. According to a study conducted by Rajasooriya and Karunarathna [59] it was observed that application of vermiwash at the rate of 75% at 6<sup>th</sup> and 8<sup>th</sup> weeks after planting recorded highest number of pods per plant, number of seeds per pod, 100 seed weight and total yield per plant in greengram. The utilization of vermiwash led to yield enhancements of 2.3 and 2.5 times in greengram compared to the control treatment. It may be due to increased concentration of vermiwash which increased the number of pods per plant and number of seeds per pod according to Bhalero *et al.* [60]. Application of Basal FYM 750kg/ha + 1% Vermiwash spray @30,45 DAS recorded significantly higher seed yield [61]. These conclusions are consistent with the research conducted by Hatti *et al.* 2010 [62] who found that the use of vermiwash spray resulted in increased growth parameters and total biomass of greengram. Vermiwash, acknowledged as an environmentally safe spray solution by Sinha *et al.* (2010) [63] demonstrated its effectiveness by acting as a source of nutrients and promoting growth, resulting in the cultivation of healthy green gram plants and increased seed production. Vermiwash, produced through the decomposition of organic matter by earthworms, is additionally recognized as a valuable liquid fertilizer, playing a significant role in enhancing crop development and productivity when applied as a foliar spray [64].

## 5.7 PANCHAGAVYA

In Sanskrit, the term "Panchagavya" denotes a mixture comprising of five products obtained from cow. When mixed appropriately and utilized, these components demonstrate extraordinary effects. Panchagavya is employed in diverse ways, such as foliar spraying, incorporating into the soil with irrigation water, and treating it with seeds or seedlings [65]. In organic farming, Panchgavya plays a pivotal role as the most favored organic fertilizer for agricultural fields, ensuring the complete elimination of harmful synthetic fertilizers, pesticides, insecticides, and antibiotics. Panchgavya stands out as the most cost-effective and beneficial fertilizer option. It enhances soil fertility, improves earthworm quality, and boosts crop health as an organic fertilizer [66,67]. Panchagavya consists of growth regulators such as Indole acetic acid, Gibberellic acid, and cytokinin, as well as important plant nutrients and beneficial microorganisms like lactic acid bacteria, yeast, and actinomycetes [68].

## 5.8 EFFECT OF PANCHAGAVYA ON CROP PRODUCTION

Panchgavya aids in producing food without the use of synthetic pesticides, facilitating the shift from conventional to organic farming, while also sustaining and enhancing crop yields, as well as improving the flavour and longevity of fruits, grains, and vegetables. Additionally, it speeds up crop maturation by 15 days and decreases production costs by minimizing expenditures on chemicals, ultimately leading to increased profits [69]. Yadav Prakash and Tripathi (2013) [70] found that utilizing foliar spray of Panchagavya at the concentration of 3 percent without additional requirement of fertilizers was the most cost-effective method for enhancing the seed yield of greengram. As per study conducted by Krishnaprabhu (2015) [71] it was revealed that foliar spray of Panchagavya (5%) resulted in a significant increase in both grain yield and yield parameters of greengram. The considerable boost in yield observed

with application of panchagavya can be mainly attributed to the significant improvement in both the number of pods per plant and the test weight. The application of a 3 percent foliar spray of Panchagavya at intervals of 15, 25, 40, and 50 days after sowing (DAS) without any fertilizers proved to be the most efficient and economical technique for increasing the grain yield of greengram [72]. A field experiment conducted by Sagaret *et al.* (2023) [73] reported that foliar application of combination of Panchagavya + in every 10 days recorded significantly higher higher number of pods per plant, number of seeds per pod, seed yield and stover yield respectively. The increase in yield and its characteristics could be a result of stimulated root growth due to inorganic nutrients, alongside enhanced absorption of water and nutrients attributed to the effects of panchagavya post-fermentation, thereby promoting higher yields. Bhargavi *et al.* (2017) [74] concluded panchagavya applied at the rate of 3 percent with 10 days interval recorded higher yield and yield attributes such as number of pods per plant, number of seeds per pods and seed yield. This could be attributed to the enhanced availability of nearly all vital nutrients to the plants, leading to enhanced vegetative growth, heightened photosynthetic activity, and subsequently, the efficient movement and accumulation of photosynthates in the economically valuable parts of the plant.

## 5.9 JEEVAMRUTH

Jeevamrutam, an organic fertilizer, is composed of a combination of cow dung, cow urine, pulse flour, jaggery, and a small amount of soil, with the intention of promoting microbial populations. It acts as a potent plant growth enhancer, enhances the biological effectiveness of crops while also activating soil, defending plants from diseases, and improving the nutritional value of fruits and vegetables Devakumar *et al.* [75]. Jeevamrut consist of macro nutrients, vital micro-nutrients, vitamins, essential amino acids, growth-promoting factors like IAA and GA, and beneficial microorganisms [76]. Applying jeevamrut to acidic soil raises its pH level, while in alkaline soil, it lowers the pH. This practice leads to sustainable increases in crop yield and eliminates the need for chemical fertilizers, thereby enhancing soil health. Jeevamrut exhibits acidity with a pH level of 4.93 and acts as a valuable reservoir of essential macro and micro nutrients, comprising nitrogen (1.97%), phosphorus (0.172%), potassium (0.29%), manganese (47 ppm), and copper (50 ppm). Jeevamrut consists of high concentration of microbial populations, with peak levels typically occurring between the 8th and 13th days after its preparation. Additionally, it promotes the growth of nitrogen-fixing bacteria in locally available materials such as farmyard manure, compost, and biogas slurry [77].

## 5.10 EFFECT OF JEEVAMRUTH ON CROP PRODUCTION

The combined organic nutrient formulations exhibit a synergistic effect and have demonstrated favorable outcomes in enhancing crop production. According to a study conducted by Kumari *et al.* (2023) [78] it was revealed that application of RDF + Vermicompost + Rhizobium + Jeevamrutam recorded higher number of pods per plant, test weight, grain yield, Straw yield as compared to control treatment. Similar findings were recorded by Verma *et al.* (2022) [79] and Marko *et al.* (2013) [80]. Jeevamrut promotes soil microbial activity, leading to enhanced soil fertility [81]. Jeevamrut is hailed as a versatile remedy for organic agriculture, catering to nutrient requirements for various crops and pest management. It is advised that Jeevamrut be prepared solely from the dung and urine of Indian cows, with the dung and urine from a single cow being adequate for organic cultivation on 12 hectares of land [82]. Many organic farmers rely heavily on jeevamrut for their organic farming practices [83,84]. Jeevamruth reportedly hosts a significant population of microorganisms capable of nitrogen fixation, phosphate solubilization, and siderophore production [85]. The organic liquid fertilizer is formulated for a fermentation process that generates beneficial live soil microorganisms, enhancing plant growth, productivity, and nutrient supply. These fertilizers offer cost-effective and environmentally friendly bioinoculants, with promising potential to boost agricultural production sustainably. Using Jeevamrut can decrease the over-reliance on chemical fertilizers, which leads to diminished soil fertility, making it an optimal substitute for chemical fertilizers [86].

**Table 2: Effect of various organic fertilizers on yield and yield attributes of greengram**

Sl. No.	Treatments	Number of pods per plant	Number of seeds per pod	Seed yield (kg ha <sup>-1</sup> )	Reference
1.	12 kg ha <sup>-1</sup> humic acid	13	10	1312.7	[37]
	9 kg ha <sup>-1</sup> humic acid	11	9	826.7	
2.	Control	12.67	9.87	6.04	[51]
	Seaweed 20%	20	10.30	11.66	
3.	Control 6 <sup>th</sup> WAP	23	9	12.68 ± 0.53	[59]
	Control 8 <sup>th</sup> WAP	21	9	11.58 ± 0.63	
	Vermiwash 75% 6 <sup>th</sup> WAP	35	11	29.02 ± 1.66	
	Vermiwash 75% 8 <sup>th</sup> WAP	35	11	29.02 ± 0.33	
4.	Panchagavya + S1 Every 10 days	28.40	10.85	870	[74]
	Jeevamruth + S1 Every 10 days	27.65	9.22	850	
	Cow urine + S1 Every 10 days	23.70	9.32	840	
5.	Control	19.10	7.5	748.5	[78]
	RDF+ Vermicompost+ Rhizobium+ Jeevamrutam	31.35	12.2	1242.5	

## 5 ADVANTAGES OF USING ORGANIC FERTILIZERS

Organic fertilizers demonstrate effectiveness in fostering environmental sustainability and enhancing plant growth over extended periods [87]. Moreover, the amalgamation of irrigation and fertilization methods has the potential to boost the efficiency of nutrient usage while diminishing the risk of nutrient leaching. Additionally, unique constituents present in liquid organic fertilizers, such as chitin, humic and fulvic acids, and other biopolymers, serve as plant biostimulants which helps in increasing productivity of crops. Several studies have indicated that the use of organic fertilizers on the soil surface can provide a substantial nutrient supply for microorganisms, resulting in a notable enhancement in both the composition and diversity of microbial communities compared to situations where such application is not present [88]. Organic fertilizers meet the biological needs of plants and also help control the populations of plant pests [89]. Organic fertilizers boost the quality and productivity of agricultural crops similarly to inorganic fertilizers, yet they do not contribute to environmental pollution [90]. Organic fertilizers furnish plants with nitrogen in a form that promotes growth without the risk of root burn or harm to beneficial soil

microorganisms. Additionally, by fulfilling the plants' nutritional needs and boosting their tolerance, organic fertilizers help prevent diseases, thereby alleviating a significant source of stress [91].

## 6 CONCLUSION

Organic fertilizer sources such as humic acid, seaweed extract, vermiwash, panchagavya and jeevamruth formulations have significant potential in increasing productivity in greengram and as such products should be encouraged in farming practices. Moreover utilizing organic fertilizers and amendments is environmentally friendly and holds a distinct advantage over synthetic, inorganic fertilizers which can lead to environmental pollution and potentially result in soil accumulation, posing risks to human health. Thus the purpose of this review is to enlighten the useful benefits of organic fertilizers and to incorporate it into the existing farming practices so as to increase the productivity of greengram while maintaining sustainable agriculture production.

## REFERENCES

1. Dash S.R, Rautaray B.K, Dhal A. Perception and Constraints faced by Pulse Growers and yield gap analysis of Green Gram (*Vigna radiata L.*) in East and South East Coastal Plain of Odisha, India. *Int.J.Curr.Microbiol.App.Sci.* 2018; 7(1): 338-346
2. Vilas J, Mallikarjuna SN, Gracy CP. Supply-demand gap analysis and projection for major pulses in India. *Econ Aff.* 2018;63:277–285
3. Available: [www.indiastat.com](http://www.indiastat.com)
4. Teggelli R, BS Ahamed Z, SM Suresh. Productivity enhancement of greengram through improved production technologies in Kalaburagi district of Karnataka. *Journal of Pharmacognosy and Phytochemistry.* 2020; 9(1): 928-930
5. Anonymous, Agricultural statistics at a glance 2022, Department of Agriculture & Farmers Welfare Economics & Statistics Division, Ministry of Agriculture & Farmers Welfare, Government of India. 2022;40-41.
6. Khairnar KY, Pokharkar VG, Kadam SA, Yadav DB. Greengram production technology: An economic analysis. *Journal of Pharmacognosy and Phytochemistry.* 2019; 8(3): 2491-2494
7. Gadi P, Dawson J, M Shankar. Effect Of Different Organic Manures, Inorganic Fertilizers And Growth Regulators On Growth And Yield Of Greengram (*VignaRadiataL.*). *Bull. Env. Pharmacol. Life Sci.* 2017; 6 (1) : 67-75
8. Almaz MG, Halim R.A, Yusoff MM. and Wahid SA. Effect of incorporation of crop residue and inorganic fertilizer on yield and grain quality of maize. *Indian Journal of Agricultural Research.* 2017; 51(6): 574-579
9. Shariff A Fazulla, Sajjan SA, Babalad HB, Nagaraj LB, Palankar GS. Effect of organics on seed yield and quality of green gram (*Vigna radiata L.*). *Legume Research.* 2017;40(2):388-392
10. Khasawneh AR, Othman YA. Organic Farming and Conservation Tillage Influenced Soil Health Component. *Fresen. Environ. Bull.* 2020; 29: 895-902.

11. Bhargavi K, Sumathi V, Reddy G. Krishna, Umamahesh V. Influence of Organic Manures and Organic Sprays on Productivity and Economics of Summer Greengram [*Vigna radiata* (L.) Wilczek]. International Journal of Environment and Climate Change. 2023; 13(10): 3111-3116
12. Kumar K. Arun, Anandakumar S, Lakshmi S. Seaweed Extract: An Eco-Friendly Biological Product for Sustainable Agriculture. In Smart Farming: A way towards Sustainability. 2022 December 1(pp.151-157). Department of Agronomy, College of Agriculture, Balaghat.
13. Devi Anjali JS, Anulekshmi AH, Salini S, Aparna RS, George S. Boronic acid functionalized nitrogen doped carbon dots for fluorescent turn-on detection of dopamine. MicrochimActa. 2017;184(10): 4081–4090
14. Wallenstein M. To restore our soils, feed the microbes. The Conversation. 2017. Available: <https://source.colostate.edu/restore-soils-feed-microbes>
15. O'Connell PF, Sustainable agriculture -A valid alternative. Outlook Agriculture. 1992;34:1527-1536.
16. Kumaravelu G, Kadamban D, Panchgavya and its effect on the growth of the greengram cultivar K-851. International Journal of Plant Sciences. 2009;4(2):409-41
17. Yadav P, Tripathi A.K. Growth and yield of greengram (*Vignaradiata*) under foliar application of panchgavya and leaf extracts of endemic plants. Indian Journal of Agronomy. 2013; 58 (4): 618-621
18. Dreesen FE. De Boeck HJ, Janssens IA, Nijs I. Summer heat and drought extremes trigger unexpected changes in productivity of a temperate annual/biannual plant community. Environ. Exp. Bot. 2012;79: 21–30
19. Bitá CE, Gerats T. Plant tolerance to high temperature in a changing environment: scientific fundamentals and production of heat tolerance crops. Frontiers of Plant Sciences. 2013;4:273–296.
20. Suzuki N, Rivero RM, Shulaev V, Blumwald E, Mittler R. Abiotic and biotic stress combinations. New Phytol. 2014; 203:32–43
21. Kaur R, Bains TS, Bindumadhava H, Nayyar H. Responses of mungbean (*Vigna radiata* L.) genotypes to heat stress. Effects on reproductive biology, leaf function and yield traits. Scientific Horticulture. 2015;197:527–541.
22. Zandalinas SI, Sales C, Beltrán J, Gómez- Cadenas A, Arbona V. Activation of secondary metabolism in citrus plants is associated to sensitivity to combined drought and high temperatures. Frontiers of Plant Sciences. 2017;7:1954.
23. Landi S, Hausman JF, Guerriero G, Esposito S. Poaceae vs. abiotic stress: Focus on drought and salt stress, recent insights and perspectives. Frontiers of Plant Sciences. 2017;8:1214
24. Swaminathan R, Singh K, Nepalia V. Insect Pests of Green Gram *Vigna radiata* (L.) Wilczek and Their Management. Agricultural Science, 2012. Dr. Godwin Aflakpui (Ed.). ISBN: 978-953-51-0567-1, InTech
25. Chauhan YS, Williams R. Physiological and Agronomic Strategies to Increase Mungbean Yield in Climatically Variable Environments of Northern Australia. Agronomy. 2018;8(6):83.

26. Malik A, Fayyaz-ul-Hassan WA, Qadir G, Asghar R. Interactive Effects of Irrigation and Phosphorus on Green Gram (*Vigna radiata L.*). Pakistan Journal of Botany. 2006;38:1119-1126
27. Akpapunam M. Mung bean (*Vigna radiata (L.) Wilczek*). In: Nwokolo, E., Smartt, J. (eds) Food and Feed from Legumes and Oilseeds. Springer, Boston. 1996:P209- 215.
28. Sharma DK, Singh A. Salinity research in India-achievements, challenges and future prospects; 2015
29. Manan Jatinder, Sharma Manoj, Singh Gurmeet and Singh Gobinder. Effect of application of various inputs by the farmers and the yield of spring maize hybrids. J Krishi Vigyan. 2016; 4 (2): 22-27.
30. Prakash O. Excessive use of chemical fertilizers, reduce the fertility power of the soil. International Journal of Engineering Inventions. 2023; 12(8):116-118
31. Kadam RP, Umate SM, Pawar GS, Lad AS. Constraint faced in adoption of recommended package practices of greengram. Agric. Update. 2014; 9(4):528-530
32. Dash SR, Rautaray BK, Dhal A. Perception and Constraints faced by Pulse Growers and yield gap analysis of Green Gram (*Vigna radiata L.*) in East and South East Coastal Plain of Odisha, India. Int.J.Curr.Microbiol.App.Sci.2018;7(1): 338-346
33. Narayan MK, Jaiswal A, Singh A. Effect of Soil and Foliar Application of Humic Acid and Brassinolide on Morpho-physiological and Yield Parameters of Black Gram (*Vigna mungo*). Nature Environment and Pollution Technology.2023;22 (4): 2201-2208
34. Bhatt P, Singh V.K. Effect of humic acid on soil properties and crop production– A review. Indian Journal of Agricultural Sciences.2022; 92 (12): 1423–1430
35. Waqas M, Ahmad B, Arif M, Munsif F, Khan AI, Amin M, Kang SM, Kim Yoo-Ha, Lee In-jung. Evaluation of Humic Acid Application Methods for Yield and Yield Components of Mungbean. Am. J. of Plant Sci.2014; 5:2269-2276
36. Hudda J, Godara A, Kukreja S. Effect of Application of Humic Acid and Vermiwash on the Growth, Quality and Yield of Plants: A Review. Int.J.Curr.Microbiol.App.Sci. 2020; 9(11): 1277-1284
37. Ali I, Kha A A, Imran, Inamullah, Khan A, Asim M, Ali I, Zib B, Khan I, Rab A, Sadiq G, Ahmad N, Iqbal B. Humic Acid and Nitrogen Levels Optimizing Productivity of Green Gram (*Vigna radiata L.*). Russian Agricultural Sciences. 2019; 45(1): 43–47
38. Tahir MM, Khurshid M, Khan MZ, Abbasi MK, Kazmi MH. Lignite-Derived Humic Acid Effect on Growth of Wheat Plants in Different Soils. Pedosphere. 2011; 21: 124-131.
39. Nikbakht A, Kafi M, Babalar M, Xia YP, Luo A, Etemadi N. Effect of Humic Acid on Plant Growth, Nutrient Uptake, and Postharvest Life of Gerbera. Journal of Plant Nutrition. 2008; 31: 2155-2167
40. Metre R N, Deotale RD, Ghewande SS, Arsode VS, Gowathami R. Effect of foliar application of humic acid through cowdungwash on chemical, biochemical and yield and yield contributing parameters of greengram. J. Soils and Crops. 2013; 23(2):369-376
41. Durand N, Brian X, and Meyer C. The effect of marine bioactive substances (N PRO) and endogenous cytokinins on nitrate reductase activity in Arabidopsis thaliana. Physiol. Plant. 2003; 119:489-493

42. Anantharaj M, Venkatesalu V. Effect of seaweed liquid fertilizer on *Vigna catajung*. Seaweed Research and Utilisation. 2001; 23(1&2): 33-39
43. Singh M P, Maity D, Saha S, Dhal K. N. Seaweed utilization and its economy in Indian agriculture. Mater Today Proc. 2022; 65(1):63–69
44. Makkar PS H, Tran G, Heuze V, Reverdin G S, Lessire M, Lebas F, Philippe A. Seaweeds for livestock diets: A review. Animal Feed Science and Technology. 2015, <http://dx.doi.org/10.1016/j.anifeedsci.2015.09.018>
45. Manimaran P, Lakshmi J, Rajasekar P. Influence of Foliar Application of Seaweed Extract and Plant Growth Regulators on Growth and Physiological Attributes of *Jasminum sambac*. Environment and Ecology. 2018; 36 (1A) : 262—264
46. Pal A , Kamthania, M. and Kumar A. Bioactive Compounds and Properties of Seaweeds—A Review. Open Access Library Journal. 2014; 1, 1-17.
47. Zhang X, Ervin EH, Schmidt ER. Plant growth regulators can enhance the recovery of Kentucky bluegrass sod from heat injury. Crop Sci. 2003; 43:952-956.
48. Iswarya S, Latha KR, Srinivasan K. Evaluation of seaweed extract on growth determinants, yield and biochemical parameters of greengram (*Vigna radiata*). J Pharmacogn Phytochem. 2019; 8:1861–1864
49. Zodape ST, Mukhopadhyaya S, Eswaran K, Reddy MP, Chikara J. Enhanced yield and nutritional quality in green gram (*Phaseolus radiata L*) treated with seaweed (*Kappaphycus alvarezii*) extract. Journal of Scientific and Industrial Research. 2010; 69:468-471.
50. Chitra G, Sreeja P.S. A Comparative Study on the Effect of Seaweed Liquid Fertilizers on the Growth and Yield of *Vigna radiata (L.)*. Nature Environment and Pollution Technology. 2013; 12(2): 359-362
51. Huda M.N, Mannan M.A, Bari M.N, Rafiquzzaman S.M, Higuchi H. Red seaweed liquid fertilizer increases growth, chlorophyll and yield of mungbean (*Vigna radiata*). Agronomy Research. 2023; 21(S1): 291–305
52. Zodape, S T Mukherjee S, Reddy MP, & Chaudhary D.R. Effect of *Kappaphycusalvarezii* (Doty) Doty ex silva extracts on grain quality, yield and some yield components of wheat (*Triticum aestivum L.*). International journal of plant production. 2009; 3(7): 97–101
53. Jaybhaye MM, Bhalerao S. Influence of Vermiwash on Germination and Growth Parameters of Seedlings of Green gram (*Vigna radiata L.*) and Black gram (*Vigna mungo L.*). Int.J.Curr.Microbiol.App.Sci.2015; 4(9): 634-643
54. Zambare V P, Padul MV, Yadav AA, and TB. Shete .Vermiwash: Biochemical and Biological approach as eco-friendly soil conditioner. ARPN Journal of Agricultural and Biological Sciences 3. 2008; (4): 28-37
55. Sharma SK, Jain D Choudhary, R Jat, G Jain P, Bhojiya AA, Jain R, Yadav, SK. Microbiological and enzymatic properties of diverse Jaivik Krishi inputs used in organic farming. Indian Journal of Traditional Knowledge. 2021; 20(1): 237-243
56. Rai, N. and Bansiwai. Vermiwash: An excellent source of nutrition for plant growth. Electronic Journal of Environmental Sciences. 2008; 1: 19-21

57. Nayak H, Rai S, Mahto R, Rani P, Yadav S, Prasad S.K, Singh R.K. Vermiwash: A potential tool for sustainable agriculture. *Journal of Pharmacognosy and Phytochemistry*. 2019; SP5: 308-312
58. Sayyad NR. Utilization of vermiwash potential against insect pests of tomato. *International Research Journal of Biological Sciences*. 2017; 6(1):44-46
59. Rajasooriya S AP, Karunarathna B. Application of vermiwash on growth and yield of green gram (*Vigna radiata*) in sandy regosol. *GRIEAST*. 2020; 14 (2):31-42
60. Bhalerao, S. A, Sharma, A. S, Poojari, A. C. Toxicity of nickel in plants. *International Journal of Pure and Applied Bio Science*. 2015; 3(2): 345-355
61. Devi S SV, Krishnan R, Somasundaram E, Ganesan K. Studies on organic cultivation practices for Greengram (*Vigna radiata L.*). *International Journal of Chemical Studies* . 2019; 7(3): 1951-1953
62. Hatti SS, Londonkar RL, Patil SB, Gangawane AK, Patil CS. Effect of *Eisenia foetida* vermiwash on the growth of plants. *Journal of crop science*. 2010; 1:6-10
63. Sinha RK, Valani D, Chauhan K, Sunita Agarwal. Embarking on a second green revolution for sustainable agriculture by vermiculture biotechnology using earthworms: Reviving the dreams of Sir Charles Darwin. *Journal of Agricultural Biotechnology and Sustainable Development*. 2010; 2(7):113-128
64. Sudhanshu V, Singh A, Pradhan S.S, Singh R.K, Singh J.P. Bio-efficacy of Organic Formulations on Crop Production-A Review. *Int.J.Curr.Microbiol.App.Sci* .2017; 6(5): 648-665
65. Ali M.N, Ghatak S, Ragul T. Biochemical analysis of Panchagavya and Sanjibani and their effect in crop yield and soil health. *Journal of Crop and Weed*. 2011; 7(2): 84-86
66. Tharmaraj K, Ganesh P, Kumar RS, Anandan A, Kolanjinathan K. A critical review on Panchagavya-a boon plant growth. *Int J Pharm Biol Arch*. 2011; 2: 1611-4
67. Pathak RK, Ram RA. Bio-enhancers: a potential tool to improve soil fertility, plant health in organic production of horticultural crops. *Progress Horticulture*. 2013; 45:237-54.
68. Venkatalakshmi K, Balasubramanian A, Sankaran N. Influence of seed treatment and foliar spray of panchagavya on growth, yield attributes and yield of *Amaranthus viride*. *Madras Agricultural Journal*. 2009; 96(1-6):135-138
69. Bajaj K, Chavhan V, Raut NA, Gurav S. *Panchgavya*: A precious gift to humankind. *J Ayurveda Integr Med*. 2022; 13(2): 100525.
70. Yadav Prakash and Tripathi AK. Growth and yield of greengram (*Vigna radiata*) under foliar application of panchagavya and leaf extracts of endemic plants. *Indian Journal of Agronomy*. 2013; 58: 618- 621.
71. Krishnaprabu S. Studies on Foliar Spray of Nutrients for Enhancing Productivity of Greengram. *International Journal of Innovative Research in Science, Engineering and Technology*. 2015; 4(9): 9457-9460
72. Somasundaram E, Sankaran N, Meena S, Thiyagarajan TM, Chandaragiri K and Panneerselvam S. Response of greengram to varied levels of Panchagavya (organic nutrition) foliar spray. *Madras Agric. J*. 2007; 90: 169-172.

73. Sagar V, Singh V, George SG. Influence of Cow Based Liquid Manures and Spraying Frequency on Growth and Yield of Green Gram (*Vigna radiata* L.) Under Natural Farming. International Journal of Environment and Climate Change. 2023; 13(9): 2973-2977
74. Bhargavi K, Sumathi V, Reddy GK, Umamahesh V, Bhargavi H. Productivity and economics of summer greengram [*Vigna radiata*] as influenced by different organic manures and organic sprays. Bull. Env. Pharmacol. Life Sci. 2018; 7 [SPL 1] : 147-151
75. Devakumar N, Shubha S, Gowder SB, Rao GG. Microbial analytical studies of traditional organic preparations beejamrutha and jeevamrutha. Building Organic Bridges. 2014;2:639-642.
76. Natarajan K, Panchagavya for plant. Proc.Nation. Conf. Glory Gomatha. S. V.Veterinary Univ, Tirupati. 2007;72-75
77. Kumar A, Avasthe,R, Babu, S, Singh R, Verma G, Gudade B, Bhupenchandra I. and Devi E. Jeevamrut: A low cost organic liquid manure in organic farming for sustainable crop production.Kerala Karshakan E Journal. 2021; 9: 32–34
78. Kumari S, Kumar R, Chouhan S, Chaudhary PL. Influence of Various Organic Amendments on Growth and Yield Attributes of Mung Bean (*Vigna radiata* L.). International Journal of Plant & Soil Science. 2023; 35(12): 124-130
79. Verma R, Singh M, Dawson J, Muddassir P, Khan I. Effect of bio-fertilizers and organic manures on growth and yield of greengram (*Vigna radiata* L.). The Pharma Innovation Journal. 2022;11(4):1599-602
80. Marko GS, Kushwaha HS, Singh S, Namdeo KN, Sharma RD. Effect of sulphur and biofertilizers on growth, yield and quality of blackgram (*Phaseolus mungo* L.). Crop Research. 2013;45:175-178.
81. Joshi, M. 2012. New Vistas of Organic Farming. Scientific Publishers, New Delhi. pp140.
82. Palekar, S. 2009. How to Practice Natural Farming? All IndiaPingalwara Charitable Society. pp 22–27
83. Joshi, M. 2008. Studies on organic farming practices in Karnataka.Project Report. University of Agricultural Sciences, Banga-lore, India, pp 12–14
84. Singh, J. 2009. Going green. The Tribune, Feb. 7, 2009.
85. Patel DM, Patel IM, Patel BT, Singh NK, Patel CK. Effect of Panchgavya and jivamrut on yield, chemical and biological properties of soil and nutrients uptake by kharif groundnut (*Arachis hypogaea* L.). International Journal of Chemical Studies 2018;3:804-809
86. Kumar A, Avasthe R K, Babu S, Singh R, Verma G, Dhaka AK, Kumar S, Saha S, Devi EL, Bhupenchandra I, Gudade B A, Hasanain M. Jeevamrut a low cost organic liquid manure inorganic farming for sustainable crop production. Kerala Karshakan E-Journal. 2021; 2: 32-34.
87. Sun J, Zhang Q, Zhou J, Wei Q.P. Pyrosequencing technology reveals the impact of different manure doses on the bacterial community in apple rhizosphere soil. Appl. Soil Ecol. 2014;78: 28–36.

88. Dianco M, Montemurro F. Long term effects of organic amendments on soil fertility. A review. *Sustainable Development*. 2010; 30(2): 401-422
89. Solomon, WGO, RW Ndana, and Y Abdulrahim. The Comparative study of the effect of organic manure cow dung and inorganic fertilizer NPK on the growth rate of maize (*Zea mays L.*). *International Research Journal of Agricultural Science and Soil Science*. 2012; 2:516-519
90. Chowdhury R. Effects of chemical fertilizers on the surrounding environment and the alternative to the chemical fertilizers. *IES- Envis Newsletter*. 2004; (3): 4-5
91. Sharma A, Chetani R. A Review on the Effect of Organic and Chemical Fertilizers on Plants. *IJRASET*. 2017; 5(2): 677-680

UNDER PEER REVIEW