

Growth, yield attributes and yield of rice as influenced by paddy residue and nitrogen management options under rice-rice cropping systems

ABSTRACT

A field experiment was conducted to know the impact of paddy residue and nitrogen management options on “Growth, yield attributes and yield of rice as influenced by paddy residue and nitrogen management options under rice-rice cropping systems during 2021-22 and 2022-23 at Regional Agricultural Research Station, Arepally, Warangal district. It is situated in the Central Telangana Zone. The experiment was laid out in factorial RBD design with 18 treatments replicated thrice. The results of the experiment revealed that grain yield, straw yield was significantly affected by both paddy straw and nitrogen management options highest was found in incorporation of crop residue along with 130% Recommended dosage of Nitrogen (RDN). Growth and Yield attributes like plant height, number of tillers per hill, panicle length, Productive tillers per hill, Number of grains per panicle, 1000 seed weight (gms) was also obtained highest in incorporation along with 130 % RDN. Results of the experiment conclude that highest growth, yield and yield attributes was obtained highest in incorporation along with 130 % RDN.

Keywords: Grain yield, straw yield, plant height, panicle length, 1000 seed weight.

Introduction: India is the second largest producer of rice after china. The area, production and productivity of rice in India about 44.79 M ha, 112.41 million tones and 2578 kg ha⁻¹ respectively. In India, major rice growing states are West Bengal, Punjab, Uttar Pradesh, Tamil Nadu, Andhra Pradesh, Telangana, Bihar, Odisha *etc.* The area and production of rice in Telangana is 2 million hectares and 6.6 million tons respectively. The area under rice cultivation in the state has increased to 2.7 Mha (2019-20) as against 1.9 Mha in 2013-14 and also production increased in leaps and bounds from 5.7 mt in 2013-14 to 9.8 mt in 2019-20 (Directorate of Economics and Statistics, Government of India, Socio Economic Outlook, 2020) (1).

Rice is the most residue producing crop in Asia (826 million tons) contributing 84% of total production of the world. Traditionally, rice straw is removed from fields for use as cattle feed and other purposes in South Asia. On average, rice crop residues contain 0.7% N, 0.23% P and 1.75% K. Therefore, the amount of NPK contained in rice crop residues produced is about 22.13×10^6 and 26.26×10^6 t year⁻¹ in Asia and the world, respectively (Goswami *et al.*, 2019) (2).

From the farmers' point of view, burning may be seen as the most suitable method of disposing of rice straw. It is not only a cost-effective method but it acts as an effective pest control procedure (Adam John, 2013) (3).

Burning of crop stubble causes the air pollution and leads to loss of huge biomass, *i.e.* organic carbon, plant nutrients, the entire amount of C, approximately 80–90% N, 25% of P, 20% of K and 50% of S present in crop residues are lost in the form of various gaseous and particulate matters, resulting in atmospheric pollution and global warming (Kaur *et al.*, 2019) (4). Incorporation in the soil poses challenges in intensive systems with two to three cropping rounds per year. This is due to the insufficient time for decomposition, leaving the straw with poor fertilization properties for the soil and hindering crop establishment. As a result, open-field burning of straw has increased dramatically over the last decade, despite being banned in most rice-growing countries because of pollution and the associated health issues. Therefore, it is important to look for sustainable solutions and technologies that can reduce the environmental footprint and add value by increasing the revenues of rice production systems (Hung *et al.*, 2020) (5).

Materials and Methods: A field experiment was conducted during 2021-22 and 2022-23 at Regional Agricultural Research Station, Arepally, Warangal district. It is situated in the Central Telangana Zone. The experiment was conducted in factorial randomized block design. It consists of factor 1 (three paddy straw management) and factor 2 (six nitrogen management options) a total of 18 treatments replicated thrice. The soil was sandy clay loam in texture. The variety chosen for rice is KNM-118 spacing adopted for rice 20 X 15 cm, plot size of 5 X 4 = 20 Sq. Meters.

Treatmental details:

Factor 1 (paddy straw management options) Factor 2 (fertilizer N levels). Among paddy straw management options it has burning (one week before land preparation), incorporation (a week before land preparation consortium is sprayed, week after its spray paddy straw will be incorporated), Cutting and removal of straw (one week before land preparation paddy straw removed and consortium will be sprayed, week after spray paddy straw is incorporated).

Among fertilizer N levels T₁ 100 % RDN (33.3% during transplanting, 33.3% maximum tillering stage (30 DAT) and 33.3 % panicle initiation stage (60 DAT), T₂ 100% RDN (40% during transplanting, 30% maximum tillering stage (30 DAT), 30% panicle initiation stage (60 DAT), T₃ 100% RDN (10% during transplanting, 30% 15 days after transplanting, 30% maximum tillering stage (30 DAT) and 30% panicle initiation stage (60 DAT), T₄ 100% RDN (15% during transplanting, 30% 15 days after transplanting, 30%

maximum tillering stage (30DAT) and 25% panicle initiation stage (60 DAT), T₅ 115% RDN (18 % during transplanting, 36% 15 days after transplanting, 36% maximum tillering stage (30 DAT) and 25% panicle initiation stage (60 DAT), T₆ 130% RDN (20% during transplanting, 42% 15 days after transplanting, 42% maximum tillering (30 DAT) stage and 26% panicle initiation stage (60 DAT). □

Microbial consortium: PJTSAU consortium consists of decomposers belonging to *genera Phanerochaeta, Asperigillus, Trichoderma*. RDF for Rabi rice - 120-60-40 kg N-P-K kg ha⁻¹. Recommended doses of phosphorus, potassium and FYM (10 t ha⁻¹) will be applied in equal quantities in all treatments. Five randomly selected plants from each plot were taken and measured for plant height (cm), number of tillers hill⁻¹, number of effective tillers per hill, number of grains per panicle, panicle length, test weight and average values were calculated.

Grain yield

Plants in the net plot area were harvested and threshed separately. Grains were dried under sun and grain yield adjusted to 12 per cent moisture. Grain yield per plot was recorded after cleaning. From this yield per plot, grain yield per hectare was computed and expressed as kg ha⁻¹.

Straw yield

After threshing the grain, the remaining straw was dried under sun and yield per plot was recorded and the yield per hectare was computed and expressed in kg ha⁻¹.

Plant height (cm)

Plant height (cm) was recorded from the five randomly selected plants, at 30, 60 and at harvest. They were pooled and average number was presented.

Number of tillers per hill

Number of tillers per hill was recorded from the five randomly selected plants, at 30,60 and at harvest. They were pooled and average number of tillers hill⁻¹ was presented.

Number of effective tillers per hill

Total number of fully opened green functional panicles produced was recorded from the five randomly selected plants, at harvest. The average number of fully opened green functional panicles per hill were treated as effective tillers per hill and presented.

Number of grains per panicle

From five randomly selected panicles, the total number of grains was counted and average was obtained.

Panicle length

Panicles were harvested from the five randomly selected plants of each treatment, panicle length was measured and average was calculated and expressed in centimeters.

Test weight

Samples of grains were collected from the produce of each treatment plot and then 1000 seeds were separated by counting from each sample. The weight of these one thousand seeds of each treatment was recorded on electronic balance to record the test weight in grams.

Results and discussion:

Growth parameters

Plant height (cm)

There is no significant effect of paddy straw and nitrogen management options on plant height and there is no interaction between two factors.

The highest plant height was observed in burning 43.34 cm at 30 DAT followed by cutting and removal 41.96 cm and least was reported in incorporation 40.68 cm.

With the factor 2 the highest was obtained in T₂ 43.56 cm followed by T₁ 42.92 cm and least was obtained in T₃ 40.29 cm.

At 60, 90 DAT highest was reported in incorporation 76.51 cm and 101.20 and least was observed in cutting and removal 72.14 cm and 95.62 cm.

With factor 2 highest was obtained in T₆ 79.53 cm, 104.29 cm followed by T₅ 77.49 cm, 102.22 cm and least was reported in T₁ at 60, 90 DAT respectively. Plant height initial decreased in the incorporation treatment at 30 DAT this is because of due to lack of available nitrogen which was immobilised and utilised by the microbes and at 60, after harvest highest was reported in incorporation along with high dose of nitrogen application in order to compensate the loss of nitrogen by immobilisation. Our results are in conformity with (Chaudhary *et al.*, 2013) (6).

Number of tillers hill⁻¹

The highest number of tillers per hill⁻¹ was obtained in burning 11.36 least was observed in 10.86 incorporation at 30 DAT.

With factor 2 highest was reported in T₂ 11.64 followed by 11.45 T₁ and least was reported in T₃.

At 60, 90 DAT highest was obtained in incorporation 12.51, 15.04 and least was reported in cutting and removal 11.94, 13.22.

With factor 2 highest was observed in T₆ 13.24, 15.20 followed by T₅ 12.83, 14.93 and least was observed in T₁ in 60, 90 DAT. There is no significant effect of paddy straw and nitrogen management options on number of tillers hill⁻¹ and there is no interaction

between two factors.

At harvest from the pooled data of two years the highest number of tillers per hill was obtained in 20.26 incorporation followed by burning 19.25 and cutting and removal 18.19 and there is significant effect of paddy straw on number of tillers per hill at harvest. Among the different levels of nitrogen the highest was obtained in T₆ 21.74 which was on par with T₅ 20.59 and the least was obtained in T₁ 16.96 and there is significant effect on number of tillers per hill. There is no significant interaction between the two factors.

Yield attributes in rice

Panicle length

With the paddy straw and nitrogen management options there is no significant effect on panicle length (cm) and there is no significant interaction between the two factors.

At harvest the highest panicle length was observed in 22.06 (cm) incorporation followed by burning 21.25 (cm) and cutting and removal 20.50 (cm).

T₆ recorded the highest with different nitrogen dose at 22.44 cm, whereas T₁ reported the lowest at 20.25 cm. Incorporation along with higher dose of Nitrogen was reported higher panicle length when compared to rest of the treatments. Our results are in conformity with (Dar *et al.*, 2000) (7).

Productive tillers per hill

With the paddy straw and nitrogen management options there is significant effect on Productive tillers per hill and there is no significant interaction between the two factors.

The highest was obtained in incorporation 15.39 followed by burning 14.27 and cutting and removal 13.15 among the paddy straw management options

With different levels of Nitrogen application the highest was obtained in T₆ 16.76 which was on par with T₅ 15.91 and the least was obtained in T₁ 11.91.

1000 seed weight (gms)

Both paddy straw and nitrogen management significantly affected the 1000 seed weight (gms) and there is no significant interaction between the two factors.

At harvest highest 1000 seed weight gms was obtained in 23.72 gms incorporation which was on par with burning 22.78 gms and the least was obtained in cutting and removal 21.35 gms.

Among the nitrogen levels highest was obtained in T₆ 23.84 gms which statistically on par with T₅ 23.28 gms T₄ 22.92 gms and T₃ 22.39 gms and the least was observed in T₁ 21.47 gms. Our results are in confirmity with (Khatri *et al.*, 2019) (8).

Number of grains per panicle

With the factor 1 highest was observed in incorporation 91.73 which was on par with burning 89.11 and least was observed in cutting and removal 85.65.

Among the nitrogen management options highest was observed in T₆ 93.97 and it is on par with T₅ 91.97, T₄ 89.62, T₃ 88.07 and least was reported in T₁ 83.15.

(Banjara *et al.*, 2019) (9) reported that plant height, number of tillers, dry matter accumulation, number of leaves hill⁻¹ and LAI was recorded significantly higher under treatment incorporation of rice straw as compared to burning and normal transplanting. The lowest yield attributes was obtained in T₁ and T₂ this is because of due to the fact that major share of N were applied during the early growth stages, produced lower grain yield. This may be attributed to the failure to synchronize the N supply as per demand of the crop at all the major system of crop growth crucial for higher yields (Chaudhary *et al.*, 2013). Delayed application of N might be helpful in keeping the plant greener for long and thereby facilitating the higher production and translocation of photosynthetic towards economic parts (Dar *et al.*, 2000). Significant difference was observed with application of different nitrogen levels in grain yield and yield attributing characters *viz*; tillers m⁻², panicle length, number of filled grains per panicle and thousands grain weight (Khatri *et al.*, 2019).

Grain and straw yield (kg ha⁻¹)

Under paddy straw management options, of pooled years the highest grain yield and straw yield was obtained in incorporation plots 7702.94 kg ha⁻¹, 9780.65 kg ha⁻¹ followed by burning 7163.11 kg ha⁻¹, 9182.87 kg ha⁻¹ and cutting and removal 6621.66 kg ha⁻¹, 8535.29 kg ha⁻¹.

Among the Nitrogen levels and time of application the highest grain and straw yield was obtained in T₆ 7955.36 kg ha⁻¹, 10056.46 kg ha⁻¹ which was on par with T₅ and the least was obtained in T₁. Grain and straw yield was found in the order 130% RDN T₆ > 115% RDN T₅ > 100% RDN T₄ > 100% RDN T₃ > 100% RDN T₂ and 100% RDN T₁ with the yield.

The increase in yield with straw incorporation is due to returning of large part of nutrients back into soil that can be observed by the plant (Bakht *et al.*, 2009) (10) and straw adds organic matter to the soil (Huang *et al.*, 2021) (11) which improves soil physical, chemical, biological properties results in improved soil health which contributes to increased grain yield Zhang *et al.*, 2016 (12); Wang *et al.*, 2015 (13). Paddy straw with wide C:N ratio showed higher nitrogen immobilization during its initial stage of decomposition, which was later released and potentially improved the available mineral nitrogen during subsequent rice growth stages thereby enhancing growth and

yield (Xu *et al.*, 2010) (14). It seems to be possible that burning of paddy straw led to loss of soil nutrients and hence a possible decrease in soil fertility (Dobermann and Fairhurst, 2002) (15), could be the reason for the lowest yields in burning treatment. Zhang *et al.*, 2010 (16) reported 7.5% yield increases in wheat crop grown with combination of fertilizer-N and crop residue return to soil. It has been ascribed to the fact that the straw returned to soils leads to large decrease in fertilizer-N losses (Grageda-Cabrera *et al.*, 2011) (17). With the increase of nitrogen fertilization grain yield increased. Similar significant results were reported by Sharma *et al.* (2021) (18), Khatri *et al.* (2019) Singh and Sharma (2020) (19) and Tian *et al.* (2018) (20) that grain yield was increased with increasing nitrogen levels.

Fageria (2014) (21) stated that higher nitrogen helps in the metabolism of protein and ultimately the metabolism of carbohydrate in the latter stages of growth which might be the cause for significantly higher production of total above ground biomass and ultimately higher production of straw.

Table 2: Effect of paddy straw and nitrogen management on number of tillers per hill at different stages in rice-rice cropping system.

Treatments	30 DAT			60 DAT			90 DAT			After harvest		
	2021-22	2022-23	pooled data	2021-22	2022-23	pooled data	2021-22	2022-23	pooled data	2021-22	2022-23	pooled data
Burning	11.46	11.26	11.36	12.49	12.17	12.33	14.74	14.37	14.55	18.70	19.80	19.25
Incorporation	10.96	10.76	10.86	12.70	12.32	12.51	15.27	14.82	15.04	19.71	20.82	20.26
Cutting and removal	11.24	10.97	11.10	12.07	11.81	11.94	13.39	12.71	13.22	17.66	18.72	18.19
SE(m)±	0.22	0.22	0.22	0.38	0.36	0.36	0.57	0.64	0.54	0.50	0.51	0.32
CD	NS	NS	NS	NS	NS	NS	NS	NS	NS	1.43	1.47	0.91
Nitrogen levels and time of application												
T1	11.57	11.33	11.45	11.51	11.17	11.34	13.35	12.87	13.11	16.33	17.58	16.96
T2	11.73	11.55	11.64	11.77	11.50	11.63	13.91	13.02	13.80	17.40	18.44	17.92
T3	10.70	10.46	10.58	12.27	11.98	12.12	14.41	13.81	14.11	18.00	19.48	18.74
T4	10.85	10.61	10.73	12.55	12.21	12.38	14.66	14.30	14.48	19.36	19.56	19.46
T5	11.11	10.90	11.01	12.99	12.67	12.83	15.07	14.78	14.93	19.73	21.44	20.59
T6	11.35	11.12	11.23	13.42	13.07	13.24	15.39	15.02	15.20	21.31	22.18	21.74
SE(m)±	0.30	0.31	0.31	0.53	0.50	0.51	0.80	0.91	0.76	0.70	0.72	0.45
CD	NS	NS	NS	NS	NS	NS	NS	NS	NS	2.02	2.07	1.28

Table 3: Yield attributes of rice as influenced by different paddy straw management options and fertilizer N levels.

Treatments	Panicle length (cm)			Number of tillers per hill			Productive tillers per hill			Number of grains per panicle			1000 seed weight (gms)		
	2021-22	2022-23	Pooled data	2021-22	2022-23	poole d data	2021-22	2022-23	poole d data	2021-22	2022-23	Poole d data	2021-22	2022-23	poole d data
Burning	21.06	21.45	21.25	18.70	19.80	19.25	13.77	14.78	14.27	90.15	88.06	89.11	22.60	22.97	22.78
Incorporation	21.86	22.27	22.06	19.71	20.82	20.26	14.94	15.84	15.39	93.05	90.040	91.73	23.51	23.93	23.72
Cutting and removal	20.32	20.69	20.50	17.66	18.72	18.19	12.58	13.73	13.15	86.98	84.31	85.65	21.16	21.53	21.35
SE(m)±	0.51	0.50	0.50	0.50	0.51	0.32	0.36	0.55	0.38	1.67	1.66	1.66	0.40	0.40	0.40
CD	NS	NS	NS	1.43	1.47	0.91	1.04	1.59	1.08	4.81	4.76	4.77	1.15	1.16	1.15
Nitrogen levels and time of application															
T1	20.05	20.45	20.25	16.33	17.58	16.96	11.38	12.44	11.91	84.33	81.97	83.15	21.28	21.66	21.47
T2	20.42	20.83	20.62	17.40	18.44	17.92	12.46	13.44	12.95	87.42	84.93	86.18	21.60	22.01	21.81
T3	20.83	21.17	21.00	18.00	19.48	18.74	13.03	13.89	13.46	89.08	87.06	88.07	22.22	22.56	22.39
T4	21.23	21.64	21.43	19.36	19.56	19.46	14.24	15.06	14.65	90.75	88.50	89.62	22.71	23.12	22.92
T5	21.71	22.08	21.90	19.73	21.44	20.59	15.40	16.42	15.91	93.35	90.59	91.97	23.09	23.46	23.28
T6	22.24	22.64	22.44	21.31	22.18	21.74	16.07	17.44	16.76	95.44	92.50	93.97	23.64	24.04	23.84
SE(m)±	0.72	0.71	0.71	0.70	0.72	0.45	0.51	0.78	0.53	2.37	2.34	2.35	0.57	0.57	0.57
CD	NS	NS	NS	2.02	2.07	1.28	1.47	2.25	1.52	6.80	6.73	6.75	1.63	1.63	1.63

Table 4: Grain yield and straw yield (kg ha⁻¹) of rice as influenced by paddy straw management options and fertilizer N levels

Treatments	Grain Yield			Straw Yield		
	2021-22	2022-23	pooled data	2021-22	2022-23	pooled data
Burning	7081.0	7245.2	7163.1	9082.3	9283.3	9182.8
Incorporation	7630.9	7774.9	7702.9	9687.3	9873.9	9780.6
Cutting and removal	6524.4	6718.9	6621.6	8414.4	8656.1	8535.2
SE(m)±	185.79	177.7	140.84	235.04	243.64	190.54
CD	533.96	510.73	404.77	675.50	700.22	547.63
Nitrogen levels and time of application						
T1	6596.2	6766.5	6681.4	8685.8	8926.4	8806.1
T2	6353.4	6520.3	6436.8	8321.9	8530.0	8425.9
T3	7068.4	7235.5	7151.9	8968.9	9170.5	9069.7
T4	6978.9	7146.2	7062.5	8788.5	9007.7	8898.1
T5	7603.4	7770.9	7687.2	9645.3	9836.7	9741.0
T6	7872.1	8038.5	7955.3	9957.6	10155.2	10056.4
SE(m)±	262.74	251.31	199.17	332.39	344.55	269.47
CD	755.13	722.28	572.43	955.31	990.26	774.46

Conclusions:

From the results of the present study it can be concluded that Paddy straw incorporation along with additional supply of Nitrogen *i.e* 130% RDN, 115% RDN produced higher grain yield, straw yield and other growth and yield attributes in rice-rice cropping system. Thus, farmers are suggested to incorporate crop residues with nitrogen dose of 130% RDN or 115 % RDN into the soil instead of burning; which is environmentally hazardous to human and soil health.

ACKNOWLEDGEMENT: The authors are thankful to the Department of soil science and agricultural chemistry, professor jayashankartelangana state agricultural university, rajendranagar, (Hyderabad) India for providing the necessary facilities to undertake the research.

COMPETING INTERESTS : Authors have declared that no competing interests exist.

References:

- Directorate of Economics and Statistics, Govt. of India; Socio-Economic Outlook 2020, Govt of Telangana (1).
- Goswami SB, Mondal R and Kumar MS. 2019. Crop Residue Management Options in Rice-Rice System: A Review. *Archives of Agronomy and Soil Science*. 66(9): 1218–1234 (2).
- Adam John. 2013. Alternatives to open-field burning on paddy farms, Agricultural and Food Policy Studies Institute, Malaysia. 18:1-5 (3).
- Kaur K, Kaur P and Sharma S. 2019. food security, Nutrition and Sustainable Agriculture - emerging Technologies. 618-620 (4).
- Hung NV, Concepcion MM, Migo MV, Quilloy R, Balingbing C, Chivenge P and Gummert M. 2020. Rice Straw Overview: Availability, Properties, and Management Practices. *Sustainable Rice Straw Management*.1-13 (5).
- Chaudhary SK, Singh Y, Pandey DN, Dharmindar. Nitrogen scheduling, phosphorus management and green maturing for increasing productivity of low land rice. *Oryza*. 2013; 50(3):253-258(6).
- Dar SA, Bali AS and Shah MH. 2000. Effect of different levels and time of nitrogen application on wet-seeded rice in Kashmir Valley. *Oryza* **37**(3):244– 246.(7)
- Khatri N, Rawal N, Chalise D, Bista M, and Pandey BP. 2020. Effect of Crop Residue and Nitrogen Level in Yield and Yield Attributing Traits of Rice under Rice-Wheat Cropping System. *International Journal of Advanced Biological and Biomedical Research*. 8(2): 146-152.(8)
- Banjara GP, Chandrakar DK and Tigga B. 2019. Performance of Rice Straw Management and Nitrogen Scheduling on Yield and Growth Parameters of Summer Rice in Chhattisgarh Plain. *Int.J.Curr.Microbiol.App.Sci*. 8(10): 1079-1083.(9)
- Bakht J, Shafi M, Jan MT and Shah Z. 2009. Influence of crop residue management, cropping system and N fertilizer on soil N and C dynamics and sustainable

- wheat (*Triticumaestivum L.*) production. *Soil and Tillage Research*. 104(2): 233–240. (10)
- Huang W, Wu JF, Pan XH, Tan XM, Zeng YJ, Shi QH, Liu TJ and Zeng YH. 2021. Effects of long-term straw return on soil organic carbon fractions and enzyme activities in a double-cropped rice paddy in South China. *Journal of Integrative Agriculture*. 20(1): 236-247.(11)
- Zhang P, Chen X, Wei T, Yang Z, Jia Z, Yang B, Han Q and Ren X. 2016. Effects of straw incorporation on the soil nutrient contents, enzyme activities, and crop yield in a semiarid region of China. *Soil and Tillage Research*. 160: 65-72.(12)
- Wang W, Lai DYF, Wang C, Pan T and Zeng C. 2015. Effects of rice straw incorporation on active soil organic carbon pools in a subtropical paddy field. *Soil and Tillage Research*. 152: 8-16.(13)
- Xu Y, Nie L, Buresh RJ, Huang J, Cui K, Xu B, Gong W and Peng S. 2010. Agronomic performance of late-season rice under different tillage, straw, and nitrogen management. *Field Crops Research*. 115(1): 79-84.(14)
- Dobermann A and Fairhurst T. 2002. Rice straw management. *Better Crops International*. 16: 7- 11.(15)
- Zhang J, Wen XX and Liao Y. 2010. Effects of different amounts of maize straw returning on soil fertility and yield of wheat winter. *Plant Nutrition Fertilizer Science*. 16(3): 612-619.(16)
- Cabrera GOA, Vera-Núñez JA, Aguilar-Acuña JL, Macías-Rodríguez L, Aguado-Santacruz GA and Peña-Cabriales JJ, 2011. Fertilizer dynamics in different tillage and crop rotation systems in a Vertisol in Central Mexico. *Nutrient Cycling in Agroecosystems*. 89(1): 125-134.(17)
- Sharma AR, Behera UK, Singh G, Kumar A and 2021. Long-term Effects of Tillage, Crop Residue and Crop Rotations on Soil Microbial Parameters under the Wheat (*Triticumaestivum*) Based Cropping Systems in Semi-Arid Northern India. *Indian Journal of Pure Applied Bioscience*. 9(1), 219-235. (18)
- Singh S and Sharma S. 2020. Temporal changes in rhizosphere biological soil quality indicators of wheat in response to nitrogen and straw incorporation. *Tropical Ecology*. 61(3): 328-344. (19)
- Tian Z, Ge Y, Zhu Q, Yu J, Zhou Q, Cai J, Jiang D, Cao W and Dai T. 2018. Soil nitrogen balance and nitrogen utilization of winter wheat affected by straw management and nitrogen application in the Yangtze river basin of China. *Archives of Agronomy and Soil Science*. 65(1): 1-15.(20)
- FageriaNK. 2014. Nitrogen management in crop production. CRC press, 24-25.(21)