

**Review Article**

**A REVIEW ON DEVELOPMENT OF HERBAL NANOGEL FOR  
ARTHRITIS**

**Comment [C1]:** The content covered under this topic is more hydrogel oriented instead of nanogel.  
Remain coherent with the topic.

UNDER PEER REVIEW

## ABSTRACT

In terms of stability and quick reaction to external stimuli, nanogels are highly favored materials in herbal medicine. In order to provide herbal products with greater cellular penetration than currently possible, nanogel is proving to be a novel approach to oral and transdermal medicine delivery for a number of untreated ailments, including. Inflammation of one or more joints causes pain and stiffness that can worsen with age. Different types of arthritis exist, each with different causes including wear and tear, infections, and underlying diseases. Symptoms include pain, swelling, reduced range of motion and stiffness. The ability of nano gel formulations to alter a drug's profile, genetic makeup, amino acids, peptides, a carbohydrate, or immunologic material, along with their capacity to pass through biological hurdles, biological distribution, and drug kinetics, has been signalled as a promising target for drug delivery systems. This may enhance effectiveness, safety, and compliance among patients, a number of reasons, including their small size, huge surface area, and their ability for covering ability to cover both hydrophilic and hydrophobic molecules, hydrophobic nanogels that are represent a promising drug delivery technique. As an innovative medication delivery method, nanogels can reduce toxicity and enhance the incorporated drug's bioavailability.

**Keywords:** Nanogel, Natural product, Arthritis

**Comment [C2]:** Summarize the abstract and mention clear objective of the study

**Comment [C3]:** Spelling mistake. It should be Highly favored

**Comment [C4]:** Replace this with 'To' for better fluency

**Comment [C5]:** Replace this with Several for better fluency

**Comment [C6]:** 'a' should be removed

**Comment [C7]:** Spelling mistake. It should be 'signaled'

**Comment [C8]:** Replace with several for better fluency

**Comment [C9]:** Grammar mistake. It should be 'include'

**Comment [C10]:** 'Their ability for covering' should be replaced with 'ability to cover'

**Comment [C11]:** Should be removed

## 1. INTRODUCTION

One of the many aspects of nanomedicine, the convergence among medication delivery and release, has been shown by nanogels in recent years of substances, medicine, and nanotechnology. Crosslinked polymer networks, or nanogels, are able to absorb large volumes.(Rane et al., 2021) Hydrogels that are less than a nanometer have been referred to as nanogels. A hydrogel is a polymer-based gel produced by combining chains of monomers to create a network of biomolecules. There are multiple methods for generating hydrogels, but all of them include the formation of polymeric monomers, which must subsequently be polymerized with functional cross-linker molecules to generate a polymer structure that looks a net.(A. Patil&Kontamwar, 2021)It is conceivable to load medications into the openings and subsequently release these later through stimulation might come from an outside source or from the body's immediate surroundings, relying on the release mechanism utilized, frequently to as the volume transitions from phase-to-phase temperature, the most typical internal–external components that result in a physical change are a specific pH and a temperature increase within a certain volume.(S. S. Patil et al., 2023)

Positive results have been reported in the bioavailability of several chemical and herbal bioactive substances utilized as nanomedicines. Considered as a nanomedicine product, nanogels exhibit exceptional stability, pharmaceutical loading competence, biologic consistency, powerful penetration ability, and several sectors, which include transferring genes, organ targeting, chemotherapeutic administration, drug diagnostics, and herbal medicine, considered nanogels as important players.(Gugulothu et al., n.d.)Nanogels have been employed in biotechnology just recently, mostly in the fields of genetics, protein synthesis, and enzyme immobilization.(S. Khan et al., 2024a) They are a source of information for advanced medical treatment plans. likewise, amphiphilic nanogels that are used because to their high drug-loading capacity, which is achieved through collection and sedimentation. Aspect of new generation nanogels,(Jadhao&Harsulakar, 2022)

## 2. USING NANOCARRIERS TO DELIVER DRUGS:

Any type of nanomaterial utilized for carrying another molecule, such a drug, is known as a nanocarrier. As nanocarriers, archetypes like liposomes, carbon-based materials, systems of polymers, and the micelles are frequently employed.(Sindhu et al.,

**Comment [C12]:** Change the wording. Replace are able to absorb with 'can absorb'

**Comment [C13]:** Change the wording. Instead of To create, it should be 'To develop'

**Comment [C14]:** Incomplete. Grammatically incorrect. It should be 'looks like a net'

**Comment [C15]:** Rewrite for clarity

**Comment [C16]:** Sentence should be start with capital letter

**Comment [C17]:** Rewrite for clarity

**Comment [C18]:** Wrong use of preposition. Grammatical mistake. It should be 'because of'

**Comment [C19]:** Incomplete sentence

**Comment [C20]:** Explain more about how nanocarriers are used to deliver drug

**Comment [C21]:** Change the wording

**Comment [C22]:** Can be rewrite in a better way

2022) On an outcome of their distinct features, nanocarriers are right now being investigated for use in the delivery of medications, and treatment may benefit from them. (Chitare et al., 2023) Upon accessing the organs, nanostructures can enter the body by a number of different methods, such as the skin, the respiratory system, the digestive tract, and medication injections. Once there, these can have serious physiological effects such oxidative stress, DNA damage, and inflammatory reactions. Because NPs may load and carry an incredible array of therapeutics actions to all body parts. (A. Khan et al., n.d.)

**Comment [C23]:** Rewrite for clarity

**Comment [C24]:** Replace with 'several'

**Comment [C25]:** It should be 'such as'

Nanogel's characteristics transportation focus because of their surface reliance and attention to local conditions, nanogel carriers are often administered at specific places via binding to them or by other "passive" strategies including retention inside physiological regions. (Hadi et al., 2019)

### 3. **LOWEST DEGREE OF THREAT:**

**Comment [C26]:** Heading should be more explained

The nanogels require to be non-toxic, environmentally friendly, and perishable that release readily expelled non-toxic breakdown products. To guarantee that each remedy is delivered successfully and with the lowest potential side effects, drug delivery needs to take occurred at the intended place of delivery. Drug loading requires to be high in order to achieve goals. Integrity of The Encapsulation; It is essential that drug molecules entrapped in nanogels are not withdrawn or liberated.

**Comment [C27]:** Change the wording

**Comment [C28]:** Rewrite for clarity

**Comment [C29]:** Change the wording

**Comment [C30]:** Replace this with 'to achieve goals'

**Comment [C31]:** Incomplete sentence

**4. TYPES OF HYDROGELS:** Hydrogels can be classified as natural or synthetic based on the type of polymer they include. Natural or synthetic polymers that have undergone hydrogenation are regarded as raw materials for medical purposes. Hydrogels are made from natural and synthetic polymers that are biocompatible, biodegradable, and, in certain cases, compatible with blood in situations where the hydrogel comes into contact with blood. (Sakthiganapathi et al., 2023)

**Comment [C32]:** Make it brief

4.1. Natural Hydrogels: Gels using polymers derived from natural sources are known as natural hydrogels. Hydrogels made from natural polymers have non-toxic, biodegradable, and biocompatibility properties. The goal of using biomaterials determines whether or not natural polymers are used in the production of hydrogels. For instance, hydrogels that are utilized to discharge compounds under control need to be biocompatible, biodegradable, and non-formal. Polysaccharides and associated proteins are examples of natural polymers that are frequently employed as carriers for the release

**Comment [C33]:** Replace with 'biocompatible properties'

**Comment [C34]:** Change the wording

of chemicals. The outcomes of in-body testing demonstrated the biocompatibility of these polymers; polysaccharides are the most appropriate due to their high durability, enzymatic degradation, biocompatibility, and lack of toxicity. Alginate, collagen, gelatin, and fibrin are examples of natural hydrogels that are utilized in clinical applications. For instance, alginate has been utilized. (Samaiya & Kumar Sharma, 2023)

**Comment [C35]:** Change the wording

4.1. **Synthetic Hydrogels:** Polyamides and polyethylene glycol are examples of synthetic polymers from which synthetic hydrogels are made. In hydrogel manufacturing, synthetic polymers have replaced natural polymers more recently because of advantages including longer life, stronger gel, and improved water absorption. Hydrogels made from synthetic polymers have a variety of medical uses. In terms of chemical composition and mechanical structure, synthetic polymers outperform natural polymers and are hydrophobic. These polymers include PEG, polyvinyl alcohol, and polyacrylamide and its derivatives. One of the most widely utilized polymers for synthetic hydrogenation in medicine is PEG, which is employed in wound dressings, tissue engineering, medication release, and bone prostheses. This polymer's qualities make it useful for a number of medicinal applications, including resistance to protein adsorption, immune system non-stimulation, and biocompatibility. PEG has the capacity to create insoluble network structures on its own. Nevertheless, the crosslinking within the hydrogen network's structure is enhanced by the addition of factor groups. (Samaiya & Kumar Sharma, 2023)

**Comment [C36]:** Complete the sentence for clarity

**Comment [C37]:** Make the content under this heading more brief and to the point.

**Comment [C38]:** Remove the phrase

**Comment [C39]:** Rewrite for clarity

**Comment [C40]:** Change the wording

**Comment [C41]:** Remove the phrase

**Comment [C42]:** Replace with 'for several'

**Comment [C43]:** This phrase should be removed and is replaced with 'PEG can develop'

## 5. **THE EVOLUTION OF NANOGELS:**

Hydrophilic, hydrophobic, or amphiphilic polymers which have been cross-linked chemically or physically give the framework of nanogel. Furthermore, proteins and polysaccharides that are chosen for their breakdown and reduced immunogenic responses could be included in the components which generate the nanogel. These have been engineered to be particularly efficient at handling the inclination of all other nanocarriers that to leak when filled with bioactive and boosting the therapeutic payload in the defined region. (Narayana Murthy, 2023) Nanogels having diameter that range from 1 to 1000 nm combined the features of nanomaterials and hydrogels. Two main criteria should be included in a distinctive drug transporter: the drug must travel at the required pace, and the medication should be effective. Two essential features should be incorporated into an exceptional drug transporter: the drug should travel at the specified pace and the medication should be effectively delivered to the appropri-

**Comment [C44]:** The content of this heading should be more focused on evolution of nanogel rather than application of nanogel. The content should remain coherent with the heading

**Comment [C45]:** Grammatical error. It should be 'That have'

**Comment [C46]:** Separate the two sentences with full stop

**Comment [C47]:** Rewrite for clarity

**Comment [C48]:** Change the wording

**Comment [C49]:** Grammatical error. It should be 'combine'

**Comment [C50]:** Rewrite for clarity

ate location. Given this, nanogels contain many of distinct features that can help create novel delivery methods. Better pharmaceutical diffusion into the biological membrane and delayed drug release form two of nanogels have primary benefits. Unlike to traditional transdermal application agents (oils, creams, and lotions), nanogels provide a more stable controlled drug carrier for topical applications. (Akotkhane et al., 2023)Because of their natural ability to swell as a result of chemical alter, with enabling the medication to be provided in the proper dose form, nanogels are employed to achieve regional and widespread drug action. Ionic delivery of medication, biological sensors, and transdermal patches may each be made using nanogels.(Jamadar & Husen Shaikh, 2017) By regulating their release, nanotechnology attempts to boost the bioavailability of weak water-soluble medicines made from herbs. The lipid and polymer makeup of these structures can be utilized for distinguishing them. Through enhancing lectins, drug delivery through carbohydrate-based nanogels, which may also be improved.(Hadi et al., 2019)

The most often used polymers for the fabrication of nanogels that are consist of chitosan, alginate, polyvinyl alcohol, and carbomers. A synthetic polymer called carbomer/Carbopol gels when its acidic pH is adjusted. The carbomer partially uncoils after dissolving in a liquid solvent, nonetheless, it is initially highly coiled. (Jiménez-Rosado & Romero, 2022)Complete uncoiling with thickening is brought about by the salt subsequent generations when the pH rises to 7.0 using sodium hydroxide/potassium hydroxide, leading to the development of a nanogel. When trying to deliver macromolecules such oligonucleotides, genes, peptides, proteins, and antigens, chitosan hydrogel nanoparticles are frequently utilized.

In addition, these nanogels contain a capacity to release the medicines after activation. For the procedure to liberate charged in the opposite direction proteins from Poly acrylic acid and Polyethylene glycol cross-linked hydrogels, The pH of the medium must either be lowered between 7.4 to 5.5 or calcium ions having an affinity to carboxylic groups must be introduced. Medicines in is enclosed in a triple-layered nanogel, which is being shown to dissolve down and release the antibiotic wheneverlipase enzyme is introduced REWRITE FOR CLARITY.(Bagde et al.,)The layers, a lipase-responsive polycaprolactone crosslinked with the particles, an ester-linked with the drug's phosphate group, and a Polyethylene glycol shell compose the nanogel. The

**Comment [C51]:** Sentence lacks coherence. Rewrite for clarity

**Comment [C52]:** Remove the phrase

**Comment [C53]:** Replace with 'develop'

**Comment [C54]:** Correct the spelling

**Comment [C55]:** Rewrite for clarity

**Comment [C56]:** Remove the phrase

**Comment [C57]:** Incorrect .Should be 'chemical alteration'

**Comment [C58]:** Change the wording

**Comment [C59]:** Rewrite for clarity

**Comment [C60]:** Rewrite for clarity

**Comment [C61]:** Rewrite for clarity

**Comment [C62]:** Grammar mistake. Remove 'are'

**Comment [C63]:** Grammar mistake. It should be 'such as'

**Comment [C64]:** Change the wording

**Comment [C65]:** Replace with drug. It would be more appropriate

**Comment [C66]:** Replace with 'charges'

**Comment [C67]:** Rewrite for clarity

PCL (polycaprolactone) is intact, which preserves the antibiotic in the core in an ordinary environment. The cellular microbial lipase dissolves the PCL.(Bagde et al.,)

**6. SYNTHESIS OF NANOGEL:** The polymerization technique and nanometre scale can be used to categorize the methods used to create nanogels. (P. D. Patil et al., 2018) This section includes an exhaustive overview of the techniques used of producing gels.

**Comment [C68]:** Remove the phrase

**Comment [C69]:** Replace with 'to develop'

**Comment [C70]:** Wrong use of preposition. It should be 'used in'

**6.1 Manual methods;** Some common physical methods to produce nanogels are inverse nanoprecipitation, microfluidics, and the Mini emulsion technique. The Mini emulsion approach, involving minuscule particles of oil-soluble surfactants present in a continuous organic phase, produces an oil-in-water emulsion. In the microfluidic method, glass chambers or a capillary tube composed of silica which simulates polymers are used for producing droplets. The most methodical technique for creating aqueous nanogels is the last system, referred to as inverse nanoprecipitation, which involves just mixing an aqueous polymeric solution with another miscible non-solvent.(P. D. Patil et al., 2018)

**Comment [C71]:** Rewrite for clarity

**Comment [C72]:** Change the wording. Write 'developing'

**Comment [C73]:** Remove the phrase

**6.2 Covalent connection;** Physical crosslinking nanogels, such as weak van der Waal forces, hydrophilic and hydrophobic contacts, and so forth, are created via non-covalent binding. These forms are less stable, and the temperature, crosslinking agent, and composition all have a significant impact on sensitivity of the gel. It has been demonstrated that the use of nanogels, which produces micelles. ~~Physical crosslinking nanogels, such as weak van der Waal forces, hydrophilic and hydrophobic contacts, and so forth, are created via non-covalent binding~~ REPEATED LINE. These variations are less consistent.(P. D. Patil et al., 2018)

**Comment [C74]:** Change the wording with 'developed'

**Comment [C75]:** Rewrite for clarity

**Comment [C76]:** Repeated sentence.

**7. ADVANTAGES OF NANOGELS:**(S. Khan et al., 2024)

**Comment [C77]:** Can be improved with proper sentence structure

1. One benefit of nanogel is that less medication is needed.

**Comment [C78]:** Can be rewrite in a better way

2. Offer defense against the drug molecule's internal biodegradation within the body system.

3. The nanogel's size can be changed based on the delivery molecule.

4. Lessen the toxicity of medications.

5. Nanogels can pass through both the blood-brain barrier and the skin's physiological barrier.

**Comment [C79]:** Rewrite for clarity

6. Drug-loaded nanogels can be applied topically and absorbed into the body without causing any negative side effects.

Comment [C80]: Remove the phrase

7. A comfortable and simple to use formulation process.

8. Suitable for a wide range of bioactive substances, including peptides, proteins, and antibodies.

9. Increase the capacity for permeation.

10. Better access to regions after intravenous delivery that hydrogel cannot reach.

### 8. LIMITATION OF NANOGELS; (Jadhao & Harsulakar, 2022)

1. One drawback of nanogel is the high cost of solvent and surfactant removal at the conclusion of preparation.

Comment [C81]: Remove the phrase and replace with 'after'

2. If any traces of surfactants or polymers are still present in the body, negative consequences could happen.

Comment [C82]: Rewrite for clarity

3. Limited ability to load drugs and inadequate release control.

4. The drug-polymer interaction may cause a structural collapse, which would improve the hydrophilicity of the nano-gel matrix and permanently trap the drug molecules.

### 9. CHARACTERISTICS OF HYDROGEL MATERIALS NECESSARY FOR ARTHRITIS;

Persistent synovitis is the primary characteristic of rheumatoid arthritis. Histologically, it shows hyperplasia of the cells, angiogenesis, an influx of leukocytes that are inflammatory and alterations in the expression of proteins, numerous cytokines, cell-surface adhesion molecules, and proteinase inhibitors. Simultaneously, the evolution of rheumatoid arthritis causes several synovial alterations. (Wang et al., 2019) The unique structure and inflammatory milieu present a problem for rheumatoid arthritis medication and cell treatments. To meet these obstacles, hydrogels for the treatment of rheumatoid arthritis should possess great comprehensive characteristics. Certain characteristics of hydrogels for rheumatoid arthritis therapy. (Wang et al., 2019) First and foremost, the most crucial and fundamental characteristic of all hydrogel types is high biocompatibility. Second, hydrogels need to possess strong mechanical characteristics. Mechanical properties such as swelling ratio, toughness, and so forth. Thirdly, hydrogels used to treat rheumatoid arthritis have crucial features such as

Comment [C83]: Replace with 'Inflammatory leukocytes'

Comment [C84]: Incomplete sentence

Comment [C85]: Rewrite for clarity

wear resistance and tribology. The fourth factor is biodegradability and the capacity to bind and release medication or cells. Finally, the transdermal hydrogels capacity to penetrate the epidermal barrier is critical.(Wang et al., 2019)

**Comment [C86]:** Low fluency .Rephrase it

## 10. CHALLENGES

Currently, rheumatoid arthritis treatment hydrogel materials' characteristics are always being enhanced. Higher requirements are placed on the applied hydrogels due to the complicated physiological environment in the articular cavity of rheumatoid arthritis.(Bagde et al., n.d.-b) These hydrogels must have a variety of qualities in order to effectively treat rheumatoid arthritis. Strong mechanical strength, robust drug or cell loading capacity, good biocompatibility, and the ability to release drugs gradually and steadily are some of these qualities.(Sakthiganapathi et al., 2023)Therefore, improving the overall characteristics of hydrogels through the fusing of diverse materials is an urgent topic that needs to be tackled right now. We can shorten experiment periods and simulate the fusion of different materials with the help of artificial intelligence. Ultimately, composite hydrogel composites with improved extensive qualities that are appropriate for rheumatoid arthritis treatment. (Bagde et al., n.d.-b)Patients with rheumatoid arthritis have considerable challenges in their treatment due to the complexity of the disease's genesis and management.(Samaiya & Kumar Sharma, 2023) The rheumatoid arthritis joints' ongoing inflammatory environment is the primary cause. Significant angiogenesis, cell proliferation, alterations in the expression of cell-surface adhesion molecules, proteases, protease inhibitors, and several cytokines, as well as ongoing bone tissue degradation until the patient loses joint function, are all consequences of persistent inflammation. Unfortunately, the majority of hydrogels used to treat rheumatoid arthritis only deliver one kind of medication which may be constrained by the hydrogel's characteristics and lead to a lesser degree of therapeutic impact than when many medications are applied. Thus, future research should concentrate on enhancing the comprehensive characteristics of drug-loaded hybrid delivery systems to co-deliver therapeutic compounds through various routes for a more synergistic therapeutic effect.(Malpure et al., 2018)

**Comment [C87]:** Remove the phrase and replace with 'to'

**Comment [C88]:** Rewrite for clarity

**Comment [C89]:** Rewrite for clarity

**Comment [C90]:** Remove the phrase

**Comment [C91]:** Rewrite for clarity

**Comment [C92]:** Grammatical error. Should be replaced with 'Lesser'

**Comment [C93]:** Change the wording

**11.FUTURE PROSPECTIVE:** Drawing from the various studies covered in this review, it can be inferred that nanogel has emerged as the most promising choice for topical drug administration during the development process of innovative drug delivery systems. Nanogel has been developed to improve pharmacodynamics and phar-

macokinetics and to address the low bioavailability of numerous medications. Many lipophilic medicines with improved therapeutic characteristics across numerous therapeutic areas are currently being developed as nanogels. Within the medical system.

**Comment [C94]:** Choose a synonym

**Comment [C95]:** Incomplete sentence

## 12. CONCLUSION AND SUMMARY:

**Comment [C96]:** Should be topic specific. Incoherency with the topic. Should be nanogel specific

Hydrogel is a three-dimensional network of polymers that is insoluble in water and can absorb bodily fluids in a biological setting. Such a polymer network is created via physical crosslinking, which includes ionic crosslinking, temperature- and pH-dependent processes, and enzyme reactions, as well as chemical crosslinking mechanisms like optical polymerization. Chemical hydrogels are created by covalent forces, whereas physical hydrogels are formed by weak secondary forces. Hydrogels are made from a variety of synthetic and natural polymers. The most significant characteristics of hydrogels are swelling, mechanical characteristics, and biological characteristics, all of which have an impact on the hydrogel's morphology and structure. Hydrogen finds application in numerous medical fields due to its structural resemblance to the extracellular matrix (ECM) and its capacity to absorb water where Hydrogen finds usage in a number of medical applications, including contact lenses, wound dressings, tissue engineering, and therapeutic drug release.

**Comment [C97]:** Remove the phrase

**Comment [C98]:** Replaced with 'developed'

**Comment [C99]:** Remove the phrase

**Comment [C100]:** Rewrite for clarity

**Comment [C101]:** Rewrite in active voice

**Comment [C102]:** Change the wording

**Comment [C103]:** Change the wording

**Comment [C104]:** Inappropriate word. Should be hydrogel

**Comment [C105]:** Inappropriate word

**Comment [C106]:** Change the wording

## REFERENCES :

- Akotkhane, M., Mahaparale, S., Ahire, S., Kanade, R., Wanere, S., & Dhanuka, T. (2023). Development and Evaluation of Herbal Nanogel for the Treatment of Oral Candidiasis. In *JCLMM* (Vol. 2, Issue 11).
- Bagde, S., Sihare, M., & Sandhya Bagde, M. (n.d.-a). Regular Issue 12), 2344-2357 Development And Evaluation Of Herbal Nanogel For Rheumatoid Arthritis Section A-Research paper Eur. *Chem. Bull*, 2022, 2344-2357.
- Bagde, S., Sihare, M., & Sandhya Bagde, M. (n.d.-b). Regular Issue 12), 2344-2357 Development And Evaluation Of Herbal Nanogel For Rheumatoid Arthritis Section A-Research paper Eur. *Chem. Bull*, 2022, 2344-2357.
- Chitare, S. S., Kailas Gosavi, T., & Patil, C. D. (2023). International Journal of Research Publication and Reviews FORMULATION AND EVALUATION OF HERBAL NANOGEL. In *International Journal of Research Publication and Reviews* (Vol. 4, Issue 5).

- Gugulothu, S., Jatoth, M., Paturi, G., Thabitha, J., Shalini, P., & Sridivya, C. (n.d.). *FORMULATION AND DEVELOPMENT OF NANOGEL CONTAINING GREEN TEA EXTRACT: AS TOPICAL DOSAGE FORM.*
- Hadi, H., Arifin, S. F., Alshami, A., Shakirah, S., Omar, S., Azri, M., Jalil, A., & Khalid, K. A. (2019a). Impact of Modern Technology on the Development of Natural-based Products. *Journal of Ayurvedic and Herbal Medicine*, 5(4), 133–142.
- Hadi, H., Arifin, S. F., Alshami, A., Shakirah, S., Omar, S., Azri, M., Jalil, A., & Khalid, K. A. (2019b). Impact of Modern Technology on the Development of Natural-based Products. *Journal of Ayurvedic and Herbal Medicine*, 5(4), 133–142.
- jadhao, P., & Harsulakar, A. A. (2022). Issue 6 www.jetir.org (ISSN-2349-5162). In *JETIR2206409 Journal of Emerging Technologies and Innovative Research* (Vol. 9).
- Jamadar, M. J., & Husen Shaikh, R. (2017). PREPARATION AND EVALUATION OF HERBAL GEL FORMULATION. In *Research Article Journal of Pharmaceutical Research & Education Journal homepage* (Vol. 1, Issue 2).
- Jiménez-Rosado, P. ; & Romero, M. ; (2022). Citation: Sánchez-Cid. *Polymers*, 2022, 3023.
- Khan, A., Parvez, N., Joshi, S. K., & Singh, A. P. (n.d.). Herbal Based Nanogel Formulation For Skin Disease-Optimization And Evaluation Parameters. *Journal of Pharmaceutical Negative Results* , 14
- Khan, S., Agnihotri, J., Sharma, D., & Khan, M. S. (2024a). Promising Approach of Nano Drug Delivery System: Herbal Nanogel. In *Acta Scientific Pharmaceutical Sciences* (Vol. 8).
- Khan, S., Agnihotri, J., Sharma, D., & Khan, M. S. (2024b). Promising Approach of Nano Drug Delivery System: Herbal Nanogel. In *Acta Scientific Pharmaceutical Sciences* (Vol. 8).
- Malpure, P. S., Patil, S. S., More, Y. M., & Nikam, P. P. (2018). A Review On-Hydrogel. *A Review On-Hydrogel. American Journal of PharmTech Research*, 8(3).
- MPharm, V. J., & Narayana Murthy, G. (2023). Formulation and characterization of analgesic and anti-inflammatory transdermal Nanogel containing combi-

nation of herbal extract and synthetic drug. ~ 130 ~ *Journal of Pharmacognosy and Phytochemistry*, 12(1), 130–134.

- Patil, A., & Kontamwar, P. (2021). *FORMULATION AND EVALUATION OF ANTIFUNGAL NANO GEL FOR TOPICAL DRUG DELIVERY SYSTEM*. 14, 2021.
- Patil, P. D., Patil, S. B., Patil, P. S., & Salunkhe, V. R. (2018). DESIGN AND DEVELOPMENT OF NANO GELS CONTAINING CARBAPOL AS POLYMER AND HYDROALCOHOLIC EXTRACT OF PTEROCARPUS MARSUPIUM AS HERBAL MEDICAMENT. *World Journal of Pharmaceutical Research*.
- Patil, S. S., Khulbe, P., Kumar, M., & Nitalikar, M. (2023). Bulletin of Environment, Pharmacology and Life Sciences Preparation and Evaluation Novel Nanogel Formulation Containing Silver Nanoparticles of Bixaorellena Seed Extract. In *Env. Pharmacol. Life Sci* (Vol. 12).
- Rane, B. R., Patil, R. S., & Jain Shri DD, A. S. (2021). FORMULATION DEVELOPMENT AND EVALUATION OF NANO GEL LOADED WITH MONTELUKAST SODIUM NIOSOMES. *International Journal of Pharmaceutical Sciences and Research*, 12(8), 4208.
- Sakthiganapathi, M., Yoganandam, G. P., & Gopal, V. (2023). Formulation, Characterization, and Evaluation of Wound Healing Potency of a Novel Mattantailam Nanogel Based on a Famous Traditional Siddha Formula. *Avicenna Journal of Medical Biotechnology*, 15(1), 38–47.
- Samaiya, S., & Kumar Sharma, V. (2023). HERBAL DRUG LOADED NANO GEL FOR EFFECTIVE TREATMENT OF SKIN DISORDERS. *International Journal of Pharmaceutical Sciences and Research*, 14(4), 1691.
- Sindhu, R. K., Gupta, R., Wadhwa, G., & Kumar, P. (2022). Modern Herbal Nanogels: Formulation, Delivery Methods, and Applications. In *Gels* (Vol. 8, Issue 2). MDPI.
- Wang, K., Hao, Y., Wang, Y., Chen, J., Mao, L., Deng, Y., Chen, J., Yuan, S., Zhang, T., Ren, J., & Liao, W. (2019). Functional Hydrogels and Their Application in Drug Delivery, Biosensors, and Tissue Engineering. In *International Journal of Polymer Science* (Vol. 2019).