

Response of NPK, Vermicompost and FYM on Physico-chemical Properties of Soil Under Cluster bean (*Cyamopsis tetragonoloba* L.)

ABSTRACT

Today, with the advancement of environment pollution and health that is obtained from the improper use of inorganic fertilizer, production and use of organic fertilizer is considered as an important approach in the field of soil science in the world's interest to investors. Research was conducted during *Kharif* season 2023 on central research farm of department of soil science and agricultural chemistry, (NAI) SHUATS, Prayagraj. Field trial was designed on Randomized Block Design with three replications and nine treatments. It may be concluded from the trial that the different level of NPK, Vermicompost and FYM in the experiment gave the greatest value. The best results were resulted the greatest Pore Space, Water Holding Capacity, Electrical Conductivity, Available Nitrogen, Phosphorus and Potassium significantly in T₉ [$@100\%$ NPK + $@$ VC 4 t ha⁻¹ + $@$ FYM 10 t ha⁻¹]. In contrast, the control treatment T₁ [Absolute Control] had the least results in all categories.

Key word : vermicompost, cluster bean, FYM, NPK.

INTRODUCTION

Soil plays a crucial role in determining the sustainable productivity of agro-ecosystems by supplying essential nutrients to growing plants. The uptake of macronutrients by plants is influenced by various factors, including interactions between major nutrients, as noted by [3]. However, soil degradation is becoming increasingly prevalent due to both natural processes and human activities, adversely impacting productivity. With the continuous growth of the human population, there is a greater demand on soil to provide essential nutrients for food and fiber production. Unfortunately, the soil's inherent ability to supply these nutrients has diminished, largely due to increased plant productivity associated with rising food demand [5]. Consequently, a significant challenge today is the development and implementation of soil, crop and nutrient management technologies that improve plant productivity while maintaining the quality of soil, water and air. Assessing soil fertility involves measuring available plant

nutrients and estimating the soil's capacity to sustain a continuous supply of nutrients for crops [4]. Nutrient availability is influenced by factors such as soil type, irrigation methods, pH levels, and organic matter content. [19], the degradation of soil quality concerning productivity or fertility encompasses physical, chemical, and biological processes. Understanding and addressing these degradation processes are essential prerequisites for implementing appropriate conservation activities to monitor and safeguard our natural resource base [13,15,16].

After nitrogen and phosphate, Potassium is the most significant necessary nutrient. It is crucial for plant cell sap, enzymatic activity, photosynthesis, the transportation of sugar, and the synthesis of protein and starch. However, Potassium does not have any chemical bonds with carbon or oxygen. Additionally, it increases the ability of plants to hold off pest and disease attacks and builds tolerance to drought conditions [19]

The application of vermicompost helps to improve and conserve the fertility of soil. Vermicompost imparts a dark colour of the soil and thereby help to maintain the temperature of soil. Vermicompost is one of the manures used by the farmer in growing crops because of early availability and presence of almost all the nutrients required by plants. The composition of vermicompost is 0.6-1.2% N, 0.13-0.22% P and 0.40-0.75. [17]

Farmyard manure refers to the decomposed mixture of dung and urine of farm animals along with litter and left-over material from roughages or fodder fed to the cattle. On an average well decomposed farmyard manure contains 0.5 percent N, 0.25 percent P_2O_5 and 0.5 percent K_2O [7].

The term "guar" derives from the sanskrit word "Gauahar," which means cow fodder or other livestock fodder. An annual legume plant known as the Cluster bean (*Cyamopsis tetragonoloba* L.) ($2n=14$) is cultivated for its edible, fodder, gum, and green fertilizer qualities. An important legume crop, the cluster bean, also known as "guar," is primarily grown under rainfed conditions in arid and semi-arid areas of India during the *Zaid* season. It is a product that tolerates drought very well. Its deeply penetrating roots give the plant the ability to use the rainfall it has access to more effectively, improving the potential for rainfed cropping. The legume can also withstand mild alkalinity and salinity conditions. There is no other legume product that is as resilient and drought-tolerant as the cluster bean [8].

Young cluster bean pods are consumed as vegetables and are an inexpensive source of energy (16 Kcal), moisture (8 g), protein (3.2 g), fat (1.4 g), carbohydrate (10.8 g), vitamin-A (65.3 IU), vitamin-C (49 mg), calcium (57 mg) and iron (4.5 mg) for every 100 g of edible portion [9]. Thus, this study is conducted to determined-aim of the

.....Indicate the contribution and the aims of the study.

2. MATERIALS AND METHODS

The experiment was conducted at research Farm of Soil Science and Agricultural Chemistry at Sam Higginbottom University of Agriculture Technology and Sciences Prayagraj the area is situated on the south of Prayagraj on the right side of the river Yamuna on the South of Rewa Road at a distance of about 6 km from Prayagraj city. It is situated at $25^{\circ} 57'$ N latitude, $81^{\circ} 59'$ E longitude and at the altitude of 98 meter above the sea level. The area of Prayagraj district comes under subtropical belt in the South east of Uttar Pradesh, which experience extremely hot summer and fairly cold winter. The maximum temperature of the location reaches up to $46^{\circ}\text{C} - 48^{\circ}\text{C}$ and seldom falls as low as $4^{\circ}\text{C} - 5^{\circ}\text{C}$. The relative humidity ranged between 20 to 94 percent. The average rainfall in this area is around 1100 mm annually.

Table 1: Physical parameters

Particulars	Scientist and Year
Textural class	Bouyoucous (1927) [2]
Soil Colour	Albert Henry Munsell (1971) [11]
Bulk density (Mg m^{-3})	Muthuval <i>et al.</i> (1992) [12]
Particle density (Mg m^{-3})	
Pore space (%)	
Water Holding Capacity (%)	

Table 2.: Chemical parameters

Parameters	Scientist and Year
Soil pH (1:2.5)	Jackson (1958) [6]
Soil EC (dS m^{-1})	Wilcox (1950) [25]
Organic Carbon (%)	Walkey and Black (1947) [24]
Available Nitrogen (kg ha^{-1})	Subbiah and Asija (1956) [21]
Available Phosphorus (kg ha^{-1})	Olsen <i>et al.</i> (1954) [26]
Available Potassium (kg ha^{-1})	Toth and Prince (1949) [23]

Table 3.: Treatment combination

Treatment	Treatment Combination	Symbol
-----------	-----------------------	--------

T₁	[Absolute Control]	R₀V₀F₀
T₂	@0% NPK + @VC 2 t ha ⁻¹ + @FYM 5 t ha ⁻¹	R₀V₁F₁
T₃	@0% NPK + @VC 4 t ha ⁻¹ + @FYM 10 t ha ⁻¹	R₀V₂F₂
T₄	@50% NPK + @VC 0 t ha ⁻¹ + @FYM 0 t ha ⁻¹	R₁V₀F₀
T₅	@50% NPK + @VC 2 t ha ⁻¹ + @FYM 5 t ha ⁻¹	R₁V₁F₁
T₆	@50% NPK + @ VC 4 t ha ⁻¹ + @FYM 10 t ha ⁻¹	R₁V₂F₂
T₇	@100% NPK + @VC 0 t ha ⁻¹ + @FYM 0 t ha ⁻¹	R₂V₀F₀
T₈	@100% NPK + @VC 2 t ha ⁻¹ + @FYM 5 t ha ⁻¹	R₂V₁F₁
T₉	@100% NPK + @ VC 4 t ha ⁻¹ + @FYM 10 t ha ⁻¹	R₂V₂F₂

Note: NPK:- 20:40:40, vermicompost:- 4 t ha⁻¹, FYM:- 10 t ha⁻¹

Indicate the methods of how the study was carried out. And the statistical tool used in the study.

3. RESULTS AND DISCUSSION (lack of citation) Please indicate the citation of other studies that can be compared or contrasted based on the results.

3.1 physical and chemical properties

In this finding research Bulk thickness, Particle Density, pH and EC was found non-significant. The critical varieties were seen in the event of pore space (%). The highest (%) pore space of soil was found in T₉ [@ 100% NPK + @ VC 4 t ha⁻¹ + @ FYM 10 t ha⁻¹] and lowest was found in T₁ [Absolute Control] serially. The huge varieties were seen in the event of Water holding capacity (%). The most extreme water holding capacity limit of soil was found in T₉ [@ 100% NPK + @ VC 4 t ha⁻¹ + @ FYM 10 t ha⁻¹] and lowest was found in T₁ [Absolute Control] serially. In the event of soil properties, we see that there was tremendous distinction between Organic carbon (%). The highest Organic carbon was kept in T₉ [@ 100% NPK + @ VC 4 t ha⁻¹ + @ FYM 10 t ha⁻¹] and lowest was found in T₁ [Absolute Control] serially. In the event of soil properties, we see that there was critical difference between Nitrogen (kg ha⁻¹) and Phosphorus (kg ha⁻¹). The highest Nitrogen and Phosphorus was kept in T₉ [@ 100% NPK + @ VC 4 t ha⁻¹ + @ FYM 10 t ha⁻¹] and lowest was found in T₁ [Absolute Control] serially in the event of soil properties, we see that there was massive contrast between Potassium (kg ha⁻¹). The highest Potassium was kept in T₉ [@ 100% NPK + @ VC 4 t ha⁻¹ + @ FYM 10 t ha⁻¹] and least was found in T₁ [Absolute Control] serially [10,17,20,22].

4. CONCLUSION

Conclusion Based on the results, the application of NPK, Vermicompost and FYM was found to improve the soil's health in references to cluster bean. Application of T₉ [@ 100% NPK + @ VC 4 t ha⁻¹ + @ FYM 10 t ha⁻¹] was found optimal for improving Soil Properties like Pore space, Water holding capacity, Electrical conductivity, Organic Carbon and Available Nitrogen, Phosphorus, Potassium.

5. ACKNOWLEDGEMENT

Success is the manifestation of diligence, perseverance, inspiration, motivations and innovations. It is my proud privilege to express a deep sense of gratitude of my Advisor Dr. Tarence Thomas, Professor and HoD, Department of Soil Science and Agricultural Chemistry, SHUATS, Prayagraj, whose generous help, untiring guidance, supervision, critical suggestions and his positive attitude towards my abilities enabled me to complete this work.

Table 4.: Influence of NPK, Vermicompost and FYM on Bulk Density, Particle Density, Pore Space, Water Holding Capacity and pH.

Treatments		Bulk Density (Mg m ⁻³)		Particle Density (Mg m ⁻³)		Pore space (%)		Water holding capacity (%)		pH	
T₁	Absolute Control	1.43	1.46	2.64	2.69	42.15	40.27	40.18	38.03	6.902	7.08
T₂	@0% NPK + @VC 2 t ha ⁻¹ + @FYM 5 t ha ⁻¹	1.39	1.43	2.63	2.68	42.86	40.86	40.76	38.61	6.869	7.047
T₃	@0% NPK + @VC 4 t ha ⁻¹ + @FYM 10 t ha ⁻¹	1.33	1.45	2.61	2.66	43.98	41.98	41.88	39.73	6.842	7.02
T₄	@50% NPK + @VC 0 t ha ⁻¹ + @FYM 0 t ha ⁻¹	1.38	1.43	2.62	2.68	43.32	41.32	41.69	39.54	6.856	7.034
T₅	@50% NPK + @VC 2 t ha ⁻¹ + @FYM 5 t ha ⁻¹	1.32	1.42	2.61	2.67	44.52	42.52	42.37	40.22	6.829	7.007
T₆	@50% NPK + @VC 4 t ha ⁻¹ + @FYM 10 t ha ⁻¹	1.28	1.44	2.6	2.65	45.64	43.64	44.14	41.99	6.796	6.974
T₇	@100% NPK + @VC 0 t ha ⁻¹ + @FYM 0 t ha ⁻¹	1.34	1.42	2.61	2.66	45.15	43.15	42.87	40.72	6.816	6.994
T₈	@100% NPK + @VC 2 t ha ⁻¹ + @FYM 5 t ha ⁻¹	1.30	1.45	2.60	2.64	46.25	44.25	44.81	42.66	6.776	6.954
T₉	@100% NPK + @VC 4 t ha ⁻¹ + @FYM 10 t ha ⁻¹	1.24	1.41	2.59	2.61	47.36	45.67	45.54	43.39	6.671	6.849
F- test		NS	NS	NS	NS	S	S	S	S	NS	NS
S.Em. (±)		-	-	-	-	1.7985	0.6009	1.364	0.5163	-	-
C.D (P=0.05)		-	-	-	-	5.3919	1.8016	4.089	1.5479	-	-




Fig 1.: Influence of NPK, Vermicompost and FYM on bulk Density, Particle Density, Pore Space, Water Holding Capacity and pH.

Table 5.: Influence of NPK, Vermicompost and FYM on Electrical Conductivity, Organic Carbon, Available Nitrogen, Phosphorus and Potassium.

Treatments		Electrical conductivity		Organic carbon		Available Nitrogen		Available Phosphorus		Available Potassium	
		(dS m ⁻³)		(%)		(kg ha ⁻¹)		(kg ha ⁻¹)		(kg ha ⁻¹)	
T₁	Absolute Control	0.253	0.247	0.39	0.33	254.68	235.18	23.62	21.62	189.74	182.7
T₂	@0% NPK + @VC 2 t ha ⁻¹ + @FYM 5 t ha ⁻¹	0.263	0.257	0.40	0.34	258.32	248.82	25.55	23.55	190.91	183.87
T₃	@0% NPK + @VC 4 t ha ⁻¹ + @FYM 10 t ha ⁻¹	0.286	0.280	0.42	0.36	265.35	255.85	27.77	25.77	195.24	188.2
T₄	@50% NPK + @VC 0 t ha ⁻¹ + @FYM 0 t ha ⁻¹	0.280	0.273	0.41	0.35	263.87	254.37	24.66	22.66	192.93	185.89
T₅	@50% NPK + @VC 2 t ha ⁻¹ + @FYM 5 t ha ⁻¹	0.296	0.290	0.44	0.36	268.59	259.09	26.58	24.58	197.13	190.09
T₆	@50% NPK + @ VC 4 t ha ⁻¹ + @FYM 10 t ha ⁻¹	0.303	0.297	0.43	0.34	275.64	266.14	28.75	26.75	200.15	193.11
T₇	@100% NPK + @VC 0 t ha ⁻¹ +@FYM 0 t ha ⁻¹	0.276	0.270	0.41	0.36	273.87	264.37	26.52	24.24	197.78	190.74
T₈	@100% NPK + @VC 2 t ha ⁻¹ + @FYM 5 t ha ⁻¹	0.306	0.300	0.44	0.35	278.65	269.15	29.64	27.64	202.25	195.21
T₉	@100% NPK + @ VC 4 t ha ⁻¹ + @FYM 10 t ha ⁻¹	0.316	0.310	0.45	0.36	283.48	273.98	31.42	29.05	206.27	199.23
F- test		S	S	S	S	S	S	S	S	S	S
S.Em. (±)		0.007	0.008	0.008	0.011	6.19	5.70	1.20	0.97	0.52	0.54
C.D (P=0.05)		0.022	0.024	0.024	0.033	18.56	17.10	3.60	2.93	1.56	1.63




Fig 2.: Influence of NPK, Vermicompost and FYM on Electrical Conductivity and Organic Carbon.




Fig 3: Influence of NPK, Vermicompost and FYM on Available Nitrogen, Phosphorus and Potassium.

REFERENCES

1. Black, C.A. (1965) Methods of soil analysis 2, Am. Soc, Agron. madison, Wisconsin, U.S.A.
2. Bouyoucos, G. J. (1927) The hydrometer as a new method for the mechanical analysis of soils. *Soil Science*, 23(6): 343-353.
3. Fageria NK, Gheyi HR, Carvalho MC, Moreira A. Root growth, nutrient uptake and use efficiency by roots of tropical legume cover crops as influenced by phosphorus fertilization. *Journal of Plant Nutrition*. 2016 May 11;39(6):781-92.
4. FAO. Crop production manual A guide to fruit and vegetable production in the Federated States of Micronesia FAO Subregional Office for the Pacific Food and Agriculture Organization of the United Nations Apia, 2020.
5. Food and Agricultural Organisation. The State of Food Security and Nutrition in the World. Rome: FAO, IFAD, UNICEF, WFP AND WHO; 2017.
6. Jackson, M. L. (1958). Soil chemical analysis, Second edition Indian Reprint, prentice hall of India, New Delhi. Pp-498.
7. Jangir, R. P., Gurjar, R. K., Beniwal, R. L., and Kaswan, S. (2014). Effect of integrated nutrient management on yield and economics of cluster bean (*Cyamopsis tetragonoloba*) under arid conditions. *Indian Journal of Agricultural Sciences*, 84(3), 359-62.
8. Kherawat, B.S., Munna Lal, Agarwal, M., Yadav, H.K. And Kumar, S. (2013). Effect of applied potassium and manganese on yield and uptake of nutrients by cluster bean (*Cyamopsis tetragonoloba*). *J. Agric. Physics*, 13(1): 22-26.
9. Kumar, V. and Singh, A. (2002). Nutrient composition of cluster bean [*Cyamopsis tetragonoloba* (L.) Taub.] green pods. *Journal of Food Science and Technology*, 39(5), 526-528.
10. Kumar, J. (2008) Physio-chemical properties of the soil, under the two-forest plantation stands around Varanasi (U.P.), India. influenced by integrated nutrient management under alley cropping system. *Indian Journal of Agricultural Sciences*; 89 (11): 1876–80.
11. Munsell, A. H. (1971). A Color Notation. Baltimore, MD: Munsell Color Company. ed.; 1(2):65.
12. Muthuval, P., Udaysoorian, C., Natesan, R., Ramaswami, P. P. (1992) Introduction to Soil analysis, Tamil Nadu Agricultural University, Coimbatore,

641002.

13. Osman KT. Soils: Principles, properties and management. Springer Science & Business Media; 2012 Dec 4.
14. Page AL, Miller RH, Keeny DR. Methods of soil and plant analysis. American Society of Agronomy, Madison; 1982.
15. Panda SC. Soil management and organic farming. Agrobios;2006.
16. Piper CS. Soil and plant analysis. Scientific Publishers; 2019.
17. Rolaniya, M. K., Thomas, T. and Singh, A. K. (2023) Response of Different Levels of NPK and FYM on Growth and Yield of Cluster Bean (*Cyamopsis tetragonoloba* L.) var. Neelam-61, *International Journal of Environment and Climate Change*, 13(8), 487-491, 2023; ISSN: 2581-8627.
18. Singh YV, Singh SK, Sahi SK, Verma SK, Yadav RN, Singh PK. Evaluation of soil fertility status from Milkipur village, Arajiline block, Varanasi, district, Uttar Pradesh, in relation to soil characteristics. *Journal of Pure and Applied Microbiology*. 2016 Jun 1;10(2):1455-61.
19. Singh, Y. V., Singh, S. K., Sahi, S. K., Verma, S. K., Yadav, R. N. and Singh, P. K. (2016) Evaluation of soil fertility status from Milkipur village, Arajiline block, Varanasi, district, Uttar Pradesh, in relation to soil characteristics. *Journal of Pure and Applied Microbiology*;10(2):1455-61.
20. Sharma, D. K., Hasan, A., Thomas, T., David, A. A., Choudhary, A. and Sharma, A. (2022) Implication of integrated nutrient management on Soil properties of Cluster bean, *The Pharma Innovation Journal*, 11(6): 151-154.
21. Subbiah, B. V. and Asija, G. L. (1956) A rapid procedure for the estimate of *Available nitrogen in soil current sciences*, 2(5): 259-260.
22. Takase, M., Sam-Amoah, Owusu, L. K. and Sekyere, J. D. (2011) The Effects Of Four Sources Of Irrigation Water On Soil Chemical And Physical Properties. *Asian Journal of Plant Sciences* **10(1)**: 92-96.
23. Toth, S. J., Prince, A. L. (1949) Estimation of Cation Exchange Capacity and exchangeable Ca, K and Na Content of Soil by Flame Photometer technique. *Soil Sci.*, 6 (7): 439- 445.
24. Walkley, A. and Black, I. A. (1947) Critical examination of rapid method for determining organic carbon in soils, effect of variance in digestion conditions and of inorganic soil

constituents. *Soil sci* pp. 632-251.

25. Wilcox, L. V. (1950) Electrical conductivity Am. water works Assoc. J 42, 775-776.

26. Olsen, S. R., Cole, C. V., Watanabe, F. S. and Dean, L. A. (1954) Estimation of available Phosphorus in soils by extraction with sodium bicarbonate (NaHCO_3), *U.S.D.A. Circular*. 939: 1-19.

UNDER PEER REVIEW