

# **Discriminant Analysis for Sex and Stature Estimation using Arm and Forearm-Hand Length; A Study among Mgbidi population of Imo State Nigeria**

## **Abstract**

**Background:** The key aspect of forensic anthropology is the estimation of biological profile characteristics such as sex, age, stature, etc. The study aims to estimate sex and stature using arm and forearm-hand length among the Mgbidi population of Imo State, Nigeria

**Method:** The study adopted a descriptive cross-sectional study design and a total of three hundred (150 males and 150 females) subjects were recruited for the study using multistage random sampling techniques. Data were obtained via direct measurements using mega-size calipers and a stadiometer. The obtained data was analyzed using IBM SPSS (version 25).

**Results:** The study shows that males are significantly higher in anthropometric values ( $p < 0.05$ ) compared to the females values. The multivariate regression among the sexes to estimate Stature using the right asymmetry shows  $R=0.62$  and  $SEE= 6.45$  and on the left asymmetry  $R=0.61$  and  $SEE=6.49$ . the stature predictive power of females was ( $R=0.52$ , 52.0%  $SEE= 5.35$ ) and males ( $R=0.46$ , 46.0%,  $SEE=5.51$ ) on the right asymmetry and the left asymmetry shows females ( $R=0.49$ , 49.0%,  $SEE=5.42$ ) and males was ( $R=0.45$ , 45.0%,  $SEE=5.72$ ). the estimation for sex shows overall significance ( $X^2=72.78$ ,  $\lambda= 0.78$ ,  $R=0.46$ ,  $p < 0.05$ ) from the right asymmetry and the left asymmetry also shows significance in sexual estimation ( $X^2= 71.84$ ,  $\lambda=0.79$ ,  $R=0.46$ ,  $p < 0.05$ ).

**Conclusion:** The study reveals males and females have different arm and forearm-hand lengths, with females having higher predictive power for stature in both asymmetries and forearm-hand length being a greater predictor for sex.

**Keywords:** Estimation, arm length, forearm-hand length, Mgbidi population, Sex and Stature

## **Introduction**

One of the key aspects of forensic anthropology is the estimation of biological profile characteristics, such as sex, age, ancestry, and stature, from skeletal remains. Sex and stature estimation are fundamental components of biological profiling, as they can provide valuable information for narrowing down the pool of potential missing persons, aiding in criminal investigations, and reconstructing the biological profile of unidentified individuals [1, 2, 3]. This field of study is essential in forensic anthropology, forensic medicine, and criminal investigations [2, 4]. Several studies have shown that there exist significant relationships between upper limb anthropometry and the overall stature of an individual by establishing robust regression equations that can accurately predict stature based on these upper limb measurements in various racial populations [4, 5, 6, 7]. Also, it has been postulated that sex-specific regression models are better used for stature estimation from limb measurements due to inherent anatomical variations between both male and female genders.

Poorhassan et al. [8] conducted a study on stature estimation from forearm length in Iranian medical students and found a correlation between forearm length and stature, suggesting that forearm measurements can be used as a reliable indicator for estimating stature in the Iranian population. Another study investigated the determination of stature from upper arm length in medical students revealing a correlation between upper arm length and stature, with regression equations developed for stature estimation in the Iranian population [9]. The findings from a previous study by Ahmed [6] having explored the estimation of sex from upper limb measurements in Sudanese adults indicated that upper limb measurements could be used to predict sex with a standard error of estimation (SEE) ranging from  $\pm 3.54$  to  $\pm 5.85$  thereby highlighting the importance of sex-specific models in forensic stature estimation. Kushwah et al [10] conducted a study to show a relationship between the combined length of the forearm and hand with stature using an Indian cadaveric male population. Their study showed that there were significant, positive correlations between the combined length of the forearm and hand and stature with a correlation coefficient of 0.668. Another study revealed that males had higher correlation coefficients for upper arm length and forearm length while females had higher correlation coefficients for hand length in association with stature [11].

The application of discriminant function analysis utilizes body morphological differences to develop mathematical models or formulas that can predict the sex and stature of an individual

based on specific body anthropometric measurements [4, 12]. By analyzing these morphological measurements from known male and female individuals of different heights, discriminant functions can be developed to classify unknown individuals into male or female categories and estimate their stature based on their characteristics. These functions essentially act as decision rules that assign a probability of sex and stature classification to an unknown individual based on their skeletal measurements [4, 13, 14]. There is a huge gap in literature within any Nigerian population concerning the application of discriminant function analysis in classifying sex and stature using upper limb anthropometric measurements – hence the study aims at estimating sex and stature using arm and forearm-hand length among the Mgbidi population of Imo State, Nigeria.

## **Materials and Methods**

### **Study design**

This study used a cross-sectional descriptive research design to gather data on the anthropometry of the hand of Mgbidi people of Imo State, Nigeria. The study lasted for six months (January 2024 to April 2024), and the study population comprised three hundred subjects (150 males and 150 females) within the age interval of 18-35 years. Mgbidi town was used as the study frame, and a multi-stage random proportionate sampling technique was employed to select the subjects without bias. The minimum sample size was calculated using the Taro Yamane formula for Quantitative research.

### **Selection criteria**

The study only recruited respondents whose grandparents and parents are of Mgbidi Origin and had not undergone any surgery or sustained any trauma that could alter the stature or hand morphology. Subjects whose ages were between 18-35 years and who had consented to participate in the study were also recruited in the study. The study excluded subjects who are not of Mgbidi origin or did not meet the age specification of the study as well as subjects who had surgery or sustained deformities that could affect standing height or hand morphology.

### **Anthropometric landmarks**

The study employed some anthropometric variable measures (arms length, forearm-hand length, and standing height), and these variables are defined as follows;

- **Arm length:** measurement from the lateral tip acromion down to the distal part of the arm (lateral and medial epicondyle)
- **Forearm-hand length;** this measurement runs from the proximal head of the radius to the most distal limit of the third finger.
- **Standing height;** this is a vertical measurement of the human body that runs from the vertex of the head to the sole when the subject is standing upright in an anatomical position



**Figure 1, showing arm length and forearm-hand length**

### **Method of data collection**

A semi-constructive descriptive questionnaire and a personal interview were used to gather the sociodemographic data for the Mgbidi people of Imo state Nigeria. This ensured that the respondent subjects met the inclusion criteria and were fit to participate in the study. The arm and forearm-hand lengths were measured using a mega-size caliper, adopting the appropriate anatomical landmarks. The standing height was measured from the vertex of the head to the sole of the foot in an upright position using a ZT-160 Goodcare stadiometer. Data readings were recorded and preserved by the authors.

## Method of Data Analysis

Data obtained were subjected to statistical analysis using the International Business Machine of Statistical Package for Social Science (IBM version 25) and results obtained were presented in the table as mean  $\pm$  standard deviation. T-test was used as an inferential statistic to evaluate sexual and asymmetry differences. Discriminant and multivariate regression were used to estimate sex and stature. A probability less than 0.05 ( $p < 0.05$ ) was considered statistically significant and 95% was denoted as confidence level.

## Results

The present study comprised three hundred subjects of Mgbidi origin who were 18-37 years of age. The comparison between the males and females has shown that males had a standing height of  $176.46 \pm 6.36$ cm, right arm length of  $34.58 \pm 3.76$ cm, right forearm-hand of  $50.91 \pm 3.27$ cm, left arm length of  $34.50 \pm 3.50$ cm and left forearm-hand length of  $50.91 \pm 3.24$  while the females had  $105.97 \pm 6.21$ cm,  $33.61 \pm 3.68$ cm,  $47.21 \pm 3.75$ cm,  $33.50 \pm 3.68$ cm and  $47.23 \pm 3.77$ cm for standing height, right arm length, right forearm-hand length, left arm length and left forearm-hand length respectively. The inference has shown that there are sexual differences in standing height, right arm length, right forearm-hand length, left arm length, and left forearm-hand length (table 2).

Table 3 shows no statistical differences between the left-side anthropometry compared to the right-side anthropometry of the Mgbidi population. Multivariate regression model between the males and females and the right side asymmetry and left asymmetry has shown that right side asymmetry has a correlation coefficient ( $R=0.62$ ) with a standard error of estimate 6.45 while for the left side, asymmetry shows a correlation coefficient ( $R=0.61$ ) with standard error of estimate 6.49. Comparing the males and the females using the right-side asymmetry anthropometry to stature (standing height) shows that females have a higher predictive power ( $R=0.52$ ) with a standard error of estimate of 5.35 while the males have a correlation coefficient ( $R=0.46$ ) with a standard error of estimate of 5.51 which in turn has shown a lower predictive power compared to the males. Using the left asymmetry anthropometry to stature estimation shows that males have a correlation coefficient (0.45) with a 5.72 standard error of estimate to stature and the females display correlation coefficient ( $R=0.49$ ) with a 5.42 standard error of estimate (Table 4). The stature model shows the Right asymmetry for males,  $S = 126.88 + AL(0.39) + FHL(0.70)$  and

females  $S = 122.19 + AL(0.32) + FHL(0.70)$  while left asymmetry for males shows  $S = 126.43 + AL(0.43) + FHL(0.69)$  and females  $S = 124.09 + AL(0.21) + FHL(0.73)$  (Table 4).

Table 5 shows the discriminant analysis using the right asymmetry to classify sex using the arm length and forearm-hand length. The overall chi-square test shows a significant value ( $X^2=72.78$ ,  $\lambda= 0.78$ , canonical correlation = 0.46, Df= 2, and  $p<0.05$ ). The discriminant functions extracted accounted for nearly (21.16%) of the variance of sexual differences and the sexual centroid displayed that male was 0.53 while female was -0.53. The discriminant model shows that  $Sex = -14.03 + R.AL(0.01) + R.FHL(0.28)$  while for specificity and validity, the discriminant classification coefficient model for males shows  $male = -123.75 + R.AL(1.69) + R.FHL(3.68)$  while  $female = -109.01 + R.AL(1.68) + R.FHL(3.38)$  (Table 5).

Discriminant function analysis was used to classify sex and using the left asymmetry of the upper extremity the test showed significance ( $X^2= 71.84$ ,  $\lambda=0.79$ , canonical correlation= 0.46, Df=2,  $p<0.05$ ). The discriminant functions accounted for nearly 21.16% of the variants of sexual differences and the sexual centroid shows that male was 0.52 and female was -0.52. A model was deduced to be;  $sex = -14.29 + l.AL(0.02) + l.FHL(0.27)$  but for more specificity, the discriminant classification coefficient shows that  $male = -127.13 + l.AL(1.88) + l.FHL(3.69)$  while  $female = -112.23 + l.AL(1.86) + l.FHL(3.41)$  (Table 6).

**Table 1. Descriptive Statistics of Stature, Arm, and Forearm-hand Length in Mgbidi Subjects**

	<b>N</b>	<b>Min</b>	<b>Max</b>	<b>Mean</b>	<b>Std. Deviation</b>
AGE (years)	300	18	37	24.57	3.52
S (cm)	300	150.5	191.0	171.21	8.18
R.AL (cm)	300	25.0	44.0	34.09	3.75
R.FHL (cm)	300	34.1	70.0	49.06	3.98
L.AL (cm)	300	25.0	44.0	34.00	3.67
L.FHL (cm)	300	34.1	70.0	49.07	3.97

*S=Standing height, R.AL= right arm length, R.FHL= right forearm-hand length, L.AL=left arm length, L.FHL= left forearm-hand length*

**Table 2. Sexual differences of the hand anthropometry of the Mgbidi subjects**

Parameters	Male	Female	T-test	p-value	Inference
S (cm)	176.46±6.36	105.97±6.21	14.43	0.000	S*
R.AL (cm)	34.58±3.76	33.61±3.68	2.27	0.024	S*
R.FHL (cm)	50.91±3.27	47.21±3.75	9.09	0.000	S*
L.AL (cm)	34.50±3.50	33.50±3.68	2.39	0.018	S*
L.FHL (cm)	50.91±3.24	47.23±3.77	9.01	0.000	S*

*S=Standing height, R.AL= right arm length, R.FHL= right forearm-hand length, L.AL=left arm length, L.FHL= left forearm-hand length and S\*= significant ( $p<0.05$ )*

**Table 3. Difference between the Right and Left-hand anthropometry**

Parameters	Right	Left	T-test	p-value	Inference
AL (cm)	34.09±3.74	34.00±3.66	3.32	0.19	NS
FHL (cm)	49.06±3.97	49.07±3.97	-0.411	0.68	NS

*AL= arm length, FHL= forearm-hand length and NS= non-significant ( $p>0.05$ )*

**Table 4. Multivariate regression of Stature estimation using Arm and forearm-hand length**

Subjects	Multivariate model	R	R squar e	Standard error of Estimate	Sig F Change
ALL	S= 102.81+ R.AL (0.37) +R.FHL (1.14)	0.62	0.34	6.45	0.00
	S= 103.32 + L.AL (0.34) + L.FHL (1.14)	0.61	0.37	6.49	0.00
<b>RIGHT</b>					
Male	S= 126.88 + AL (0.39) + FHL (0.70)	0.46	0.21	5.51	0.00
Female	S= 122.19 + AL (0.32) + FHL (0.70)	0.52	0.27	5.35	0.00

**LEFT**

Male	$S=126.43 + \mathbf{AL} (0.43) + \mathbf{FHL} (0.69)$	0.45	0.21	5.72	0.00
Female	$S= 124.09 + \mathbf{AL} (0.21) + \mathbf{FHL} (0.73)$	0.49	0.24	5.42	0.00

*S=Standing height, R.AL= right arm length, R.FHL= right forearm-hand length, L.AL=left arm length, L.FHL= left forearm-hand length and AL= arm length, FHL= forearm-hand length*

**Table 5. Sex discriminant function of right-hand anthropometry of Mgbidi subjects**

<b>Eigenvalue</b>	<b>Canonical correlation</b>	<b>Wilks' Lambda</b>	<b>Chi-square</b>	<b>Df</b>	<b>Inference</b>
0.28	0.46	0.78	72.78	2	<b>Significant</b>
<b>Sex centroid</b>					
Male	0.53	<b>Sex= -14.03 + R.AL (0.01) + R.FHL (0.28)</b>			
Female	-0.53				
<b>Classification coefficient</b>					
Male = -123.75 + <b>R.AL</b> (1.69) + <b>R.FHL</b> (3.68)					
Female = -109.01 + <b>R.AL</b> (1.68) + <b>R.FHL</b> (3.38)					
<i>R.AL= right arm length, R.FHL= right forearm-hand length,</i>					

**Table 6. Sex discriminant function of left-hand anthropometry of Mgbidi subjects**

<b>Eigenvalue</b>	<b>Canonical correlation</b>	<b>Wilks' Lambda</b>	<b>Chi-square</b>	<b>Df</b>	<b>Inference</b>
0.27	0.46	0.79	71.84	2	<b>Significant</b>
<b>Sex centroid</b>					
Male	0.52	<b>Sex= -14.29 + L.AL (0.02) + L.FHL (0.27)</b>			
Female	-0.52				
<b>Classification coefficient</b>					
Male = -127.13 + <b>L.AL</b> (1.88) + <b>L.FHL</b> (3.69)					
Female = -112.23 + <b>L.AL</b> (1.86) + <b>L.FHL</b> (3.41)					
<i>L.AL=left arm length, L.FHL= left forearm-hand length</i>					

## DISCUSSION

The fundamental difficulty in anthropological study remains the classification of humans and animals. The present study evaluates sexual dimorphism using the arm and forearm-hand length among the Mgbidi population of Imo State, Nigeria. The findings present that Males had a standing height of  $176.46 \pm 6.36$ cm, right arm length of  $34.58 \pm 3.76$ cm, right forearm-hand length of  $50.91 \pm 3.27$ cm, left arm length of  $34.50 \pm 3.50$ cm, and left forearm-hand length of  $50.91 \pm 3.24$ cm, while females had  $105.97 \pm 6.21$ cm,  $33.61 \pm 3.68$ cm,  $47.21 \pm 3.75$ cm,  $33.50 \pm 3.68$ cm, and  $47.23 \pm 3.77$ cm for standing height, right arm length, and right forearm-hand length. Our findings have shown that males have higher mean values compared to females which directly showed gender-based variation in standing height, right arm and forearm-hand lengths, and left arm and forearm-hand lengths. The study's findings are consistent with the fact that males have higher anthropometry values than females [15] due to factors like hormonal surge during puberty. During puberty, males typically experience a significant increase in testosterone production, which promotes growth spurts and contributes to the development of longer bones and larger body size compared to females. The growth plates located at the ends of our long bones are the key players in making us taller as we grow up. Interestingly, these growth plates close later in males than in females, hinting that males might keep growing taller for a bit longer compared to females. This prolonged growth period in males could lead to bigger differences in anthropometric measurements between adult males and females. To mention but a few, social and cultural factors among the Mgbidi population could also be attributed to the variations observed because males from Mgbidi to be precise, are handymen and well known for their creativity and industrialization so the males tend to have more muscle mass compared to females. However, Diet intake can also be attributed to the significant differences observed, because due to the nature of the daily activities males consumed more carbohydrates “Okpa, Ede, ji, Akpu, Abacha, Achicha, Garri” protein and fat to acquire enough energy for works, directly and indirectly promoting growth compared to the females whom by culture are treated like a princess “Ada”. The sexual-based variation observed in this present study is consistent with previous works among different populations, that revealed that males differ significantly ( $p < 0.05$ ) compared to females [16, 17, 18, 19, 20, 21, 22, 23].

The study also explores the bilateral symmetry of the Upper extremity anthropometry (arm and forearm-hand length) to evaluate if there are possible differences in length. The findings present no statistical differences between the left-side anthropometry and the right-side anthropometry of the Mgbidi population. However, variations in the bilateral structure could arise from different factors such as genetic mutation or developmental anomalies, environmental influences, or compensatory mechanisms arising from injuries. But, subjects with such factors were excluded from the study, the study only employed non-pathological samples. The findings of this study agree with Howley et al., [22] that bilateral symmetry of the arm and forearm length shows no statistical differences.

Stature construction from the upper limb extremity (arm and forearm-hand length) has been explored greatly in this study and the findings present that the sex of the right side asymmetry has a correlation coefficient ( $R=0.62$ ,  $SEE= 6.45$ ) while the left asymmetry has a correlation ( $R=0.61$ ,  $SEE=6.49$ ). The findings showed that the left and right asymmetry have a similar predictive power for stature with no statistical differences. However, when compared with sexes using the right asymmetry, the findings show that the females have a higher predictive power ( $R=0.52$ ,  $SEE= 5.35$ ) compared to the males ( $R=0.46$ ,  $SEE= 5.51$ ) and this could be attributed to some factors influencing estimation such as bone length, muscle mass and fat distribution but since females typically have more adipose tissue which is less dense than muscle tissue, their arm and forearm lengths may be slightly longer relative to their overall stature compared to males. This can result in a more accurate prediction of stature in females as the SEE is smaller in females than in males. On the other hand, considering the left asymmetry in comparison with sex, the findings have shown similarity with the right asymmetry. The findings of this study agree with the findings of Navid et al. [16] among medical students at Tehran University of Medical Sciences, Tehran, Iran that using arm length, females had more predictive power compared to the males. In 2008, Kanchan et al. [24] also reported that females have a higher power of prediction of stature compared to males and our findings are also consistent with Akhlaghi et al., [17]. On the contrary, our findings disagreed with Shakya et al. [25], who reported that males have a higher predictive power for stature using upper limb anthropometry compared to females but Ilayperuma et al. [12] findings were consistent with ours. Our study showed the stature model for the Right asymmetry for males,  $S = 126.88 + AL(0.39) + FHL(0.70)$  and females  $S = 122.19 + AL(0.32) + FHL(0.70)$  while left asymmetry for males

shows  $S = 126.43 + AL(0.43) + FHL(0.69)$  and females  $S = 124.09 + AL(0.21) + FHL(0.73)$ .

Sexual discrimination was also explored to classify sex (male and female) using the arm and forearm-hand length from the bilateral asymmetry. The findings from the right asymmetry present that the estimation using arm and forearm-hand length is significant ( $X^2=72.78$ ,  $\lambda=0.78$ ,  $p<0.05$ ). 21.16% accounted for the variances in sexual differences, where the sex centroid shows that applying the discriminant model  $Sex = -14.03 + R.AL(0.01) + R.FHL(0.28)$ , with the study parameters, a positive resultant value from 0-0.53 predicts males and from 0 - -0.53 predicts females. The findings have shown that forearm-hand length has 99.5% accuracy in predicting sex and arm length has 2% accuracy. Our work disagrees with the findings of Shah et al. [21] that arm length is a strong predictor of sex among communities in the Ahmedabad district of Gujarat, India. The works of Shakya et al. [25] are consistent with our study that arm lengths are weak predictors of sex. Table 5, further shows the specificity of the model in classifying males and females by applying the measures to the  $male = -123.75 + R.AL(1.69) + R.FHL(3.68)$  while  $female = -109.01 + R.AL(1.68) + R.FHL(3.38)$ . considering the left asymmetry the discriminant function analysis displayed overall significance ( $X^2=71.84$ ,  $\lambda=0.79$ ,  $p<0.05$ ) which is consistent with [21, 25, 26,]. The sex centroid explains that the resultant application of the model will give a positive value predicting males and a negative value predicting females, but the specificity classification coefficient was employed to either predict males or females accurately. However, the finding shows that arm length has a predictive accuracy of 26.4% and forearm-hand 99.8% using the left asymmetry. This contradicts findings by Shah et al. [21] and Charisi et al. [27].

The study has shown some similarities and differences in sex and stature estimation using arm and forearm-hand length among various populations and these differences could be attributed to various factors, like environmental, race, and body composition.

## **Conclusion**

The study evaluated the sexual dimorphism, and bilateral asymmetry difference of the arm and forearm-hand length. It thereafter, estimated sex and stature using anthropometric standards. The findings show that males differ in the anthropometric value of their arm and forearm-hand

length. There was no statistical difference observed in the bilateral asymmetry of both male and female populations of Mgbidi. Using arm and forearm-hand length for the estimation of stature among sexes, our findings show that females have higher predictive power compared to the males and forearm-hand length serves as a greater predictor for sex in both asymmetries compared to the arm length.

### **Consent**

According to international standards or university Standards, a written consent form was issued to every research participant underlining the aim, purpose, method, and confidentiality of the data. The filled consent forms were retrieved and preserved by the authors.

### **Ethical Approval**

The study was approved by the research and ethics committee of the University of Port Harcourt, Port Harcourt, Nigeria.

## REFERENCES

1. Spradley MK. Metric methods for the biological profile in forensic anthropology: sex, ancestry, and stature. *Academic forensic pathology*. 2016 Sep;6(3):391-9.
2. Austin D, King RE. The biological profile of unidentified human remains in a forensic context. *Academic forensic pathology*. 2016 Sep;6(3):370-90.
3. Krüger GC, Liebenberg L, Myburgh J, Meyer A, Oetlé AC, Botha D, Brits DM, Kenyhercz MW, Stull KE, Sutherland C, L'Abbé EN. Forensic anthropology and the biological profile in South Africa. *New Perspectives in Forensic Human Skeletal Identification* 2018 Jan 1 (pp. 313-321). Academic Press.
4. Krishan K, Chatterjee PM, Kanchan T, Kaur S, Baryah N, Singh RK. A review of sex estimation techniques during examination of skeletal remains in forensic anthropology casework. *Forensic science international*. 2016 Apr 1;261:165-e1.
5. Akhlaghi M, Hajibeygi M, Zamani N, Moradi B. Estimation of stature from upper limb anthropometry in Iranian population. *Journal of forensic and legal medicine*. 2012 Jul 1;19(5):280-4.
6. Ahmed AA. Estimation of sex from the upper limb measurements of Sudanese adults. *Journal of forensic and legal medicine*. 2013 Nov 1;20(8):1041-7.
7. Howley D, Howley P, Oxenham MF. Estimation of sex and stature using anthropometry of the upper extremity in an Australian population. *Forensic science international*. 2018 Jun 1;287:220-e1.
8. Poorhassan M, Mokhtari T, Navid S, Rezaei M, Sheikhezadi A, Mojaverrostami S, Hassanzadeh G. Stature estimation from forearm length: an anthropological study in Iranian medical students. *Journal of Contemporary Medical Sciences*. 2017 Jun 1;3(11):270-.
9. Navid S, Mokhtari T, Alizamir T, Arabkheradmand A, Hassanzadeh G. Determination of stature from upper arm length in medical students. *Anatomical Sciences Journal*. 2014 Aug 10;11(3):135-40.

10. Kushwah N, Thakur PS, Rastogi AK, Dadu SK, Singh BK. An autopsy-based correlation study about developing standards for estimation of stature from anthropometry of combined length of forearm and hand in male population of central India region (Indore, MP). *Indian Congress of Forensic Medicine & Toxicology*. 2018 Mar;16(1):8-13.
11. El AA, Din SM, Elkholy S, Yousef MI. Prediction of stature based on upper limb measurements among Egyptian population. *Eur J Forensic Sci*. 2016;3(2):1.
12. Papaioannou VA, Kranioti EF, Joveneaux P, Nathena D, Michalodimitrakis M. Sexual dimorphism of the scapula and the clavicle in a contemporary Greek population: applications in forensic identification. *Forensic Science International*. 2012 Apr 10;217(1-3):231-e1.
13. Ousley SD, Jantz RL. Fordisc 3 and statistical methods for estimating sex and ancestry. A companion to forensic anthropology. 2012 Apr 20;311-29.
14. Chatterjee PM, Krishan K, Singh RK, Kanchan T. Sex estimation from the femur using discriminant function analysis in a Central Indian population. *Medicine, Science and the Law*. 2020 Apr;60(2):112-21.
15. Fewehinmi HB, Okoh PD, Oghenemavwe LE, Amadi MA, Ebieto CE, Bobbo KA, Asiwe N. Facial Analysis of the Igbo Ethnic Group of Nigeria for the Evaluation of Sexual Dimorphism. *Asian Journal of Medicine and Health*. 2023 Dec 20;21(12):67-75.
16. Navid S, Mokhtari T, Alizamir T, Arabkheradmand A, Hassanzadeh G. Determination of stature from upper arm length in medical students. *Anatomical Sciences Journal*. 2014 Aug 10;11(3):135-40.
17. Akhlaghi M, Hajibeygi M, Zamani N, Moradi B. Estimation of stature from upper limb anthropometry in Iranian population. *Journal of forensic and legal medicine*. 2012 Jul 1;19(5):280-4.
18. Uzun ÖZ, Yegİnoğlu G, Ertemoğlu Öksüz CA, Kalkışım ŞA, Zihni Nİ. Estimation of stature from upper extremity anthropometric measurements. *Journal of Clinical and Diagnostic Research*. 2019;13.

19. Singh B, Kaur M, Kaur J, Singh M, Batra A. Estimation of stature from forearm length in north Indians—an anthropometric study. *International Journal of Basic and Applied Medical Sciences*. 2013;3(1):201-4.
20. Gh M, Shiasi M, Hassanzadeh G, Alaghebandha N, Dehbashipour A, Zeidi H, Barbarestani M. Anthropometric characteristics of upper limb in Iranian and Pakistani subjects. *Journal of Gorgan University of Medical Sciences*. 2014 Sep 10;16(3):80-5.
21. Shah T, Patel M, Nath S, Menon SK. A model for construction of height and sex from shoulder width, arm length and foot length by regression method. *Journal of Forensic Science & Criminology*. 2015 Feb 27;3(1):102.
22. Howley D, Howley P, Oxenham MF. Estimation of sex and stature using anthropometry of the upper extremity in an Australian population. *Forensic science international*. 2018 Jun 1;287:220-e1.
23. Özaslan A, Koç S, Özaslan İ, Tuğcu H. Estimation of stature from upper extremity. *Military medicine*. 2006 Apr 1;171(4):288-91.
24. Kanchan T, Menezes RG, Moudgil R, Kaur R, Kotian MS, Garg RK. Stature estimation from foot dimensions. *Forensic Science International*. 2008 Aug 6;179(2-3):241-e1.
25. Shakya T, Mishra D, Pandey P. Estimation of stature from upper arm length. *Int J Health Sci Res*. 2021;11(5):23-9.
26. Ilayperuma I, Nanayakkara G, Palahepitiya N. A model for the estimation of personal stature from the length of forearm. *Int J Morphol*. 2010 Jan 1;28(4):1081-6.
27. Charisi D, Eliopoulos C, Vanna V, Koilias CG, Manolis SK. Sexual dimorphism of the arm bones in a modern Greek population. *Journal of forensic sciences*. 2011 Jan;56(1):10-8.