

# Computation of Heritability for Production and Reproduction Traits in Frieswal Cattle under Field Progeny Testing

## ABSTRACT

**Aims:** To study the genetic parameter (heritability) for production and reproduction traits in Frieswal cattle under field progeny testing programme.

**Place and Duration of Study:** G B Pant University of Agriculture and Technology, Pantnagar, Uttarakhand, India, between March, 2023 and March, 2024.

**Methodology:** The present study was conducted on 1163 Frieswal cattle comprised of five different sets maintained at six field units over a period from 2013-2021. The traits considered were age at sexual maturity (ASM), age at first calving (AFC), gestation period (GP), test day peak yield (TDPY), 305 days milk yield (305D-MY), fat percentage (FP), lactation length (LL), calving interval (CI), service period (SP), number of services per conception (NSPC), dry period (DP). The data were analyzed for the estimation of heritability by the software WOMBAT.

**Results:** The heritability estimates of the traits namely; ASM, AFC, GP, TDPY, 305D-MY, FP, LL, CI, SP, NSPC and DP were  $0.469\pm 0.082$ ,  $0.467\pm 0.082$ ,  $0.259\pm 0.067$ ,  $0.248\pm 0.059$ ,  $0.354\pm 0.013$ ,  $0.413\pm 0.069$ ,  $0.389\pm 0.071$ ,  $0.394\pm 0.071$ ,  $0.452\pm 0.072$ ,  $0.072\pm 0.046$  and  $0.08\pm 0.06$  respectively.

**Conclusion:** Therefore, the heritability estimates were found to be low to high magnitude for different reproduction and production traits and further it has been concluded that, this study will aid breeders in selecting high-performing animals for future generations, taking into account the genetic factors that influence milk production and reproductive traits during early lactation.

*Keywords: Frieswal, Heritability, Production traits, Reproduction traits*

## 1. INTRODUCTION

Frieswal is a synthetic strain of cattle developed by crossing Holstein Friesian (62.5%) and Sahiwal (37.5%) at military dairy farms with an objective of milk yield of 4000, 4500 and 5000 kg milk yield in first, second and third lactation, respectively, with 4% fat in a standard lactation length of 305 days [4]. The progeny-testing initiative commenced during the Third Five-Year Plan at the cattle breeding farm in Hissar, focusing initially on Haryana cows. This programme aimed to guarantee the production and recognition of high-quality bulls with known genetic merit. Rather than relying solely on the yield of their dams, the programme evaluated bulls based on the actual performance of their progeny [12]. The genetic makeup of a population can be examined by assessing the comparative significance of hereditary influences and environmental factors on the performance of individuals within that population [19]. To evaluate the performance and variation concerning both genetic and non-genetic influences, it's essential to obtain estimates of phenotypic and genetic parameters. This enables the utilization of genetically determined variations for enhancing traits and exploring potential associations between different characteristics [52]. Assessing the extent of additive genetic variability in economically important traits provides insight into the potential for genetic enhancement of those traits through selective breeding [5]. The present investigation was conducted to compute the heritability among different production and reproduction traits in Frieswal cattle under field progeny testing.

## 2. MATERIAL AND METHODS

### 2.1 Source of data and data collection

Data of 1163 first lactation records of Frieswal cows sired by 69 sires spread ~~over a period of for~~ 9 years (2013-2021) maintained at Animal Genetics and Breeding Division, GBPUA&T, Pantnagar, under the pre-existing All India Coordinated Research Project (AICRP) on progeny testing were used for present investigation. The Field Progeny Testing (FPT) programme for Frieswal cattle in the Udham Singh Nagar district of Uttarakhand was launched by ICAR-CIRC, Meerut (Uttar Pradesh). This district is situated in the Tarai region of the Kumaon division, positioned approximately between 29° 1' N latitude and 79° 31' E longitude, with an average elevation of approximately 521 meters.

## 2.2 Data ~~editing~~Editing

Records of animals with known pedigree and normal lactation were taken into account. Any instances of culling, mid-lactation disposal, abortion, stillbirth, or pathological conditions impacting lactation yield were regarded as abnormalities and thus excluded from the analysis. Animals producing less than 1500 kg of milk and having a lactation length shorter than 200 days were also not included in the study.

## 2.3 Traits studied

A total of eleven traits viz., 305-days milk yield (305D-MY), Test day peak yield (TDPY), Fat percentage (FP), Lactation length (LL), Age at sexual maturity (ASM), Age at first calving (AFC), Gestation period (GP), Calving interval (CI), Service period (SP), Dry period (DP) and Number of services per conception (NSPC) were studied.

## 2.4 Statistical analysis

~~Paternal~~The paternal half-sib correlation (intra-sire correlation among daughters) method as described by [8] was used to estimate the heritabilities of different traits. The following statistical model was used:

$$Y_{ij} = \mu + S_i + e_{ij}$$

Where,

$Y_{ij}$  = Adjusted value of  $j^{\text{th}}$  progeny of  $i^{\text{th}}$  sire,  $\mu$  = Overall population mean,  $S_i$  = Effect of  $i^{\text{th}}$  sire,  $e_{ij}$  = Random error, assumed to be normally and independently distributed with mean zero and constant variance i.e. NID (0,  $\sigma_e^2$ )

The standard error of heritability is determined by following the procedure outlined by [56]:

$$SE = \frac{\sqrt{2(n-1)(1-t^2)(1+(k-1)t^2)}}{\sqrt{K^2(N-S)(S-1)}}$$

Where,

$t$  = intra-class correlation,  $S$  = number of sire,  $N$  = total number of observation

$K$  = average number of progeny per sire

## 3. RESULT AND DISCUSSION

~~The~~ table 1 shows the heritability estimates for various production and reproduction traits in Frieswal cattle under ~~the~~ field progeny testing programme.

### 3.1 Age at Sexual Maturity (ASM)

This trait was observed to have moderately high heritability, indicating that a considerable portion of the variation in age at sexual maturity among Frieswal cattle is due to genetic factors. Selective breeding can be employed to target early sexual maturity, which can lead to earlier breeding age and increased lifetime productivity in the herd. There ~~are~~ ~~is~~ ~~little~~ ~~few~~ or scanty literature available on ~~the~~ estimation of this genetic parameter of age at sexual maturity. [58] in Brahman cattle, [54] and [38] in crossbred cattle have reported almost similar estimates. [37] found even ~~high~~ ~~er~~ ~~estimates~~ than the ones observed in the present study. However, [1] reported ~~a~~ lower estimate in crossbred cattle than found in the present study.

### 3.2 Age at First Calving (AFC)

Similar to age at sexual maturity, age at first calving also exhibited a moderately high heritability, suggesting that genetic factors significantly influenced the timing of the first calving. Early calving age is

desirable for efficient dairy production, as it allows for more lactation cycles over the lifetime of the cow. [6] reported almost similar estimates in Frieswal cattle. [11], [26] and [43] reported lower estimates in crossbred cattle, and [62] in Holstein Friesian cattle. [10] and [31] reported a low estimate in Frieswal cattle. [7] reported a very low estimate in Sahiwal cattle than in the present study.

### 3.3 Gestation Period (GP)

Heritability estimate for the gestation period was found to be relatively low compared to production traits, indicating that genetic factors play a smaller role in determining the length of pregnancy. Management practices such as nutrition and health care may have a more substantial impact on the gestation period compared to genetic selection. The heritability estimate of the gestation period in the present study was in agreement with the reports of [2], [13] and [27] in Holstein Friesian and crossbred cattle, respectively. [47] also reported a similar estimate in Sahiwal cattle. [50], [36] and [48] reported a lower estimate in Sahiwal cattle than the present study. [42] and [18] reported a higher estimate in Holstein Friesian cattle.

### 3.4 Test Day Peak Yield (TDPY)

Test day peak yield also exhibited moderate heritability, suggesting that genetic factors contribute to a significant extent to the maximum milk yield recorded on a single test day during lactation. Selective breeding for higher peak yields can lead to increased overall milk production in the herd. [49] and [21] in crossbred cattle and [41] in Holstein Friesian cattle reported similar estimates. [21] in Sahiwal cattle and [15] in Frieswal cattle reported higher heritability estimates than found in the present study. [45] and [48] reported lower estimates in Holstein Friesian and Sahiwal cattle, respectively. [34] and [32] reported a lower estimate in Frieswal.

### 3.5 305 Days Milk Yield (305D-MY)

Relatively higher heritability of 305 days of milk yield indicated that genetic factors played a substantial role in determining the total milk production over a standardized lactation period. Selective breeding for higher milk yield can result in improved productivity in Frieswal cattle. [3], [21] and [26] reported similar estimates in Phule Triveni and crossbred cattle, respectively. [53] reported a similar estimate in Karan Fries cattle. [16], [44], [48] and [14] reported lower estimates in Karan Fries and Sahiwal cattle, respectively. [18] reported a similar estimate in Holstein Friesian cattle. [33] and [52], however, reported a high estimate in crossbred and Karan Fries cattle, respectively.

**Table 1. Heritability for Production and Reproduction Traits in Frieswal Cattle (N=1163)**

S.N.	Traits	$h^2 \pm S.E.$	Abbreviation
1	Age at Sexual Maturity	0.469±0.082	ASM
2	Age at First calving	0.467±0.082	AFC
3	Gestation Period	0.259±0.067	GP
4	Test Day Peak Yield	0.248±0.059	TDPY
5	305 Days Milk Yield	0.354±0.013	305D-MY
6	Fat Percentage	0.413±0.069	FP
7	Lactation Length	0.389±0.071	LL
8	Calving Interval	0.394±0.071	CI
9	Service Period	0.452±0.072	SP
10	Number of Services Per Conception	0.072±0.046	NSPC
11	Dry Period	0.08±0.06	DP

### 3.6 Fat Percentage (FP)

Fat percentage in milk was observed to have moderately high heritability, suggesting that genetic factors influence the proportion of fat content in the milk produced by Frieswal cattle. Selective breeding can be employed to enhance milk composition, targeting higher fat content if desired by the market. [55] reported a similar estimate in Holstein Friesian cattle. [24] in Holstein Friesian cattle and [23] in crossbred cattle

reported higher estimates of heritability than found in the present study.[61]and [57] have reported lower heritability estimates in crossbred cattle.

### 3.7 Lactation Length (LL)

Lactation length had a moderately high heritability estimate (Table 1), which indicated that genetic factors significantly influenced the duration of milk production during a lactation cycle. Selective breeding for longer lactation lengths can lead to increased milk production efficiency in the herd. [11], [59], [30] and [26] reported lower estimates in crossbred cattle. [18] reported higher estimates in Holstein Friesian cattle. [46] and [28] reported a higher estimate in Sahiwal cattle.

### 3.8 Calving Interval (CI)

The calving interval exhibited a moderately high heritability ( $0.394 \pm 0.071$ ). Similar findings have also been reported by [11] in crossbred cattle. Moderately high heritability for calving interval suggested that genetic factors play a significant role in determining the time between consecutive calvings. Selective breeding for shorter calving intervals can improve reproductive efficiency and overall herd productivity. [29] and [40] reported low estimates in Frieswal cattle. [18] and [26] also reported lower estimates in Holstein Friesian and crossbred cattle. [17] and [39] reported a high heritability estimate of calving interval in crossbred cattle.

### 3.9 Service Period (SP)

The heritability value of the service period was observed to be  $0.452 \pm 0.072$  which was moderately high. [20] and [35] also found similar estimates in Frieswal and crossbred cattle, respectively. The service period is a reproductive trait which is lowly heritable, however, the high estimate in the present study indicated a greater influence of genetic factors on this trait in Frieswal cattle. [40] in Frieswal cattle [26] in crossbred cattle, [18] in Holstein Friesian and [48] and [14] in Sahiwal cattle reported low estimates.

### 3.10 Number of Services per Conception (NSPC)

This trait being reproductive in nature, was found to have low heritability ( $0.072 \pm 0.046$ ), which suggested that genetic factors had a minor role in determining the successful conception. [22], [9] and [62] have reported similar heritability estimates in Holstein Friesian cattle. [47] reported a higher estimate in Sahiwal and [36] reported a slightly higher estimate in Sahiwal cattle than the present study. [62] reported a lower estimate in Holstein Friesian cattle than in the present study. [60] reported a low estimate and [32] reported a high estimate in crossbred cattle. Now-a-days, the artificial insemination technique is adopted for inseminating cattle, therefore, several number numbers of services depends upon the skill of AI workers in recognition of appropriate heat and deposition of semen at an appropriate site in the female reproductive tract. In addition to this, other managerial practices such as reproductive health may also have a more significant impact on this trait compared to genetic selection.

### 3.11 Dry Period (DP)

The dry period had an extremely low heritability (Table 1). [25] in Sahiwal cattle and [48] in Sahiwal cattle reported similar estimates of heritability. [26] reported higher estimates in crossbred and [46] reported high estimates in Sahiwal cattle. The heritability estimate in the present study suggested that genetic factors had negligible influence on the length of the dry period. Management practices such as nutrition and udder health management are crucial for determining the optimal dry period length in dairy cows.

## 4. CONCLUSION

In summary, understanding the heritability of production and reproduction traits is essential for designing effective breeding programmes aimed at improving overall productivity, and reproductive efficiency. By focusing on traits with higher heritability, such as milk yield, fat percentage, and reproductive performance, breeders can make more informed decisions to genetically enhance the traits of interest in the population.

## REFERENCES

1. Ali I, Muhammad SS, Shafiq M. Heritability estimates and genetic correlations of various production and reproductive traits of different grades of dairy cattle reared under subtropical conditions. *Reproduction in Domestic Animals*. 2019; 54(7):1026-1033.
2. Al-Samarai FR. Relationships between milk yield and fertility in [holstein-Holstein](#) cattle. *Life Sciences International Journal*. 2008; 2(4):819-824.
3. Ambhore GS, Singh A, Deokar DK, Singh M, Sahoo SK. Phenotypic, genetic and environmental trends of production traits in Phule Triveni synthetic cow. *Indian Journal of Animal Sciences*, 2017; 87(6):736–41.
4. Annual Report, 2019. ICAR-Central Institute for Research on Cattle, Meerut Cantt. – 250 001, Uttar Pradesh, India.
5. Balasubramaniam S, Singh M, Gowane GR, Kumar S. Estimate of genetic and non-genetic parameters and trends for age at first calving in Sahiwal cows. *The Indian Journal of Animal Sciences*. 2013; 83(9).
6. Banerjee S. 1996. Genetic studies on reproductive and productive traits in HF X S crosses of military dairy farms. Thesis, MSc. (Ag.) G. B. Pant University of Agriculture and Technology, Pantnagar.
7. Bansal SK, Bhawe, KG, Shirsath TV, Joshi SA, Marimuthu S, Khadse JR. Genetic analysis of age at first calving in graded Sahiwal cows in the field conditions. *Indian Journal of Veterinary Research*. 2018; 27(1):23-26.
8. Becker WA. 1975. *Manual of Quantitative Genetics*. 4th Edn., Washington State University, Pullman. Washington. [The USA](#).
9. Ben Zaabza H, Ben Gara A, Hammami H, Jemmali B, Ferchichi MA and Rekik B. Genetic parameters of reproductive traits in Tunisian Holsteins. *Archives Animal Breeding*. 2016; 59:209-213.
10. Bhaduria SS and Kathpatal BG. Factors affecting age at first calving in Friesian x Sahiwal crosses. *Indian Journal of Animal Research*. 2003; 30:70-72.
11. Chamwazi FM, Nguluma A, Nziku ZC. Effect of Genetic and Non-Genetic Factors on Productive and Reproductive Performances of Dairy Cattle at TALIRI–Tanga. *Asian Journal of Research in Animal and Veterinary Sciences*, 2024; 7(1):17-28.
12. Das AK, Kumar R, Singh U, Kumar S, Raja TV, Sirohi AS. Field progeny testing of Frieswal bulls-Retrospective and perspective. *The Indian Journal of Animal Sciences*. 2022; 92(8):968-975.
13. Das PK, Ali SZ, Islam ABMM, Roy BK. A Comparative Study of Productive and Reproductive Performance and Estimates of Heritability for Economic Traits in Different Genetic Groups of Cattle Available at Baghabarighat Milk Pocket Area of Bangladesh. *Online Journal of Biological Science*. 2003; 3:726–740.
14. Dash S, Parveen K., Singh M, Bharadwaj A, Singh KP, Rahim A. Performance evaluation, genetic parameter and genetic trend estimation of production and reproduction traits of Sahiwal cattle in Chhattisgarh. *Indian Journal of Animal Research*. 2023; 57(1):1-6.
15. Deshpande KS and Bonde HS. Genetic studies on peak yield in Holstein Friesian x Sahiwal cross-bred cattle. *The journal of agricultural sciences*. 2009; 97(3):707-711.
16. Divya P, Singh A, Gandhi RS, Singh RK. 2014. Estimation of breeding values of first lactation 305 days milk yield from single and [multi-multi](#)-trait animal models in Karan Fries cattle. *Indian Journal of Animal Science*. 2014; 84(10):1085-1089.

17. Dubey PP and Singh CV. Estimates of genetic and phenotypic parameters considering first lactation and lifetime performance traits in Sahiwal and crossbred cattle. *The Indian Journal of Animal Sciences*. 2005; 75(11):1289-1294.
18. Easa A, Rashad AM, Saleh AA. Evaluation of some economic traits in Holstein dairy cattle under Egyptian conditions. *Journal of Advanced Veterinary Research*. 2024; 14(1):198-203.
19. Falconer DS and Mackay TF. *Introduction to Quantitative geneticsGenetics*. 4<sup>th</sup> edition. Pearson Education. London: Longman; 2009.
20. Gaur GK. Estimating breeding values of Frieswal bulls for the performance traits. *Indian journal Journal of Animal ScienceScience*. 2003; 73(1):79-82.
21. Girimal DG, Kumar D, Shahi BN, Ghosh AK, Kumar S. Genetic evaluation of Sahiwal and crossbred cattle for some economic traits. *Pantnagar Journal of researchResearch*. 2020; 18(2):153-157.
22. Guo G, Guo X, Wang Y, Zhang X, Zhang S, Li X, Liu L, S Wanhai, Usman T, Wang X, Du L, Zhang, Q. Estimation of genetic parameters of fertility traits in Chinese Holstein cattle. *Canadian Journal of Animal Science*. 2014; 94(2):281-285.
23. Jayawardana JMDR, Lopez-Villalobos N, McNaughton LR, Hickson RE. (2023). Genomic regions associated with milk composition and fertility traits in spring-calved dairy cows in New Zealand. *Genes*. 2023; 14(4):860.
24. Junior GO, Schenkel FS, Alcantara L, Houlahan K, Lynch C, Baes CF. Estimated genetic parameters for all genetically evaluated traits in Canadian Holsteins. *Journal of dairy science*. 2021; 104(8):9002-9015.
25. Kannan D. 2002. Lifetime performance evaluation of Sahiwal cattle. MSc. Thesis, NDRI, Deemed University, Karnal, Haryana, India.
26. Kaur S, Ghosh AK, Shahi B. Assessment of genetic parameters for first lactation production traits in crossbred cattle in India. *Indian Journal of Animal Research*. 2023; 13(1):33-41.
27. Kumar A, Mandal A, Gupta AK, Ratwan P. Genetic and environmental causes of variation in gestation length of Jersey crossbred cattle. *Veterinary World*. 2016; 9(4):351.
28. Kumar A, Singh CV, Kumar D. 2010. Genetic studies of economic traits in Sahiwal cattle. *Indian Journal of Animal Production and Management*. 2010; 26(1-2): 89-92.
29. Kumar A. 2003. Genetic analysis of economic traits of Sahiwal cattle using animal models. Thesis, Ph.D-D, G.B. Pant University of Agriculture and technologyTechnology, Pantnagar.
30. Kumar BS, Gangaraju G, Suresh J, Kumari BP, Vinod U. (2023). Effect of Non-Genetic Factors on Productive Traits in Jersey-Sahiwal Crossbred Cows in Andhra Pradesh. *Indian Journal of Animal Research*. 2023; 13(4):557-563.
31. Kumar P. 2015. Genetic evaluation of Frieswal sires for test day milk records and first lactation. MVSc. Thesis, LUVAS, Hisar, India.
32. Kumar R, Das AK, Raja TV, Rathee SK, Dubey PP, Prakash B. (2017). Performance of crossbred cattle (HFx Sahiwal) under tropical farming conditions of Punjab. *Indian Journal of Animal Sciences*, 2017; 87(11):1402-1405.
33. Kumari S. (2015). Studies on genetic and non-genetic factors affecting the prediction of first lactation milk yield from part records in crossbred cattle (Doctoral dissertation, Guru Angad Dev Veterinary and Animal Sciences University, Ludhiana).
34. Lakshmi BS, Gupta BR, Sudhakar K, Prakash MG, Sharma S. (2009). Genetic analysis of production performance of Holstein Friesianx Sahiwal cows. *Tamilnadu Journal of Veterinary and Animal Sciences*. 2009; 5(4):143-148.
35. Lodhi G, Singh CV, Barwal RS, Shahi BN. Genetic and phenotypic parameters of first lactation and lifetime traits in crossbred cattle. *International Journal of Agricultural Policy and Research*. 2016; 4(8):143-148.
36. Lomole, M. A. (1989). Reproductive traits and milk yield relationship of the Kenya sahiwal cattle. Doctoral dissertation, University of Nairobi, Nairobi, Kenya.
37. MacNeil MD, Cundiff LV, Dinkel CA, Koch RM. Genetics correlations among sex-limited traits in beef cattle. *Journal of Animal Science*. 1984; 58(5):1171-1180.
38. Martínez-Velázquez G, Gregory KE, Bennett GL, Van Vleck LD. Genetic relationships between scrotal circumference and female reproductive traits. *Journal of animal science*. 2003; 81(2):395-401.

39. Moges TG. 2008. Genetic studies on first lactation traits and evaluation of sires using different multitraits sire evaluation in crossbred cattle. M.Sc. (Ag.) Thesis, G.B. Pant University of Agriculture and Technology, Pantnagar.
40. Mukharjee S. (2005). Genetic evaluation of Frieswal cattle. Ph.D. Thesis. NDRI. Deemed University. Karnal, India.
41. Nawaz A, Nizamani AH, Marghazani IB, Nasrullah N, Fatih A. Influence of genetic and environmental factors on lactation performance of Holstein Friesian cattle in Balochistan. *Journal Animal and Plant Science*. 2013; 23:17-19.
42. Olson KM, Cassell BG, McAllister AJ, Washburn SP. Dystocia, stillbirth, gestation length, and birth weight in Holstein, Jersey, and reciprocal crosses from a planned experiment. *Journal of dairy science*. 2009; 92(12):6167-6175.
43. Pardo AM, Elzo MA, Gama LT, Melucci LM. Genetic parameters for growth and cow productivity traits in Angus, Hereford and crossbred cattle. *Livestock Science*. 2020; 233:103952.
44. Parveen K, Gupta AK, Gandhi RS, Chakravarty AK, Mumtaz, S. (2018). Genetic analysis of trends in production traits of Sahiwal cows over years using BLUP animal model. *Indian Journal of Dairy Science*. 2018; 71(4):396-403.
45. Radwan HAA, Abo Elfadl EA, Fardos A. (2015). Estimates of population parameters for some economic traits in Holstein Friesian cows by using a statistical program. *Global Veterinaria*. 2015; 14(1):129-135.
46. Raja KN. (2004). Genetic and phenotypic trends for production and reproduction performance of Sahiwal cattle. Thesis, MVSc. National Dairy Research Institute, Karnal, India. 94p.
47. Ratwan P, Chakravarty AK, Kumar M, Gupta AK. Genetic analysis of reproductive traits of Sahiwal cattle. *Indian Journal of Animal Sciences*. 2019; 89(9):961-965.
48. Ratwan P, Kumar M, Chakravarty AK. (2024). Bayesian approach for assessment of co-variances and genetic parameters of economically important traits in Sahiwal cattle. *Research Square*. 2024; DOI: <https://doi.org/10.21203/rs.3.rs-3905504/v1>
49. Ratwan P, Kumar M, Kumar A, Chakravarty AK, Mandal A. (2018). Impact of additive direct and maternal heritability on production efficiency traits in Jersey crossbred cattle. *Indian Journal of Animal Sciences*. 2018; 88(7):848-852.
50. Selvan AS, Tantia MS, Kumaresan A, Kumar A, Kumar DR, Karuthadurai T, Upadhyay A. Phenotypic and genetic parameters estimation for birth weight in Zebu and crossbred calves born under organized farm conditions in India. *International Journal of Livestock Research*, 2018; 8(06):48-58.
51. Shahi BN and Kumar D. Sire evaluation using first lactation traits and univariate animal models in Sahiwal and its crosses. *Indian Journal of Animal Science*. 2006; 76(10):853-854.
52. Singh MK and Gurnani M. Performance evaluation of Karan Fries and Karan Swiss cattle under closed breeding system. *Asian-Australasian journal of animal sciences*. 2004; 17(1):1-6.
53. Singh RK. 2013. Genetic evaluation of Karan Fries sires using multitrait models. Thesis, Ph.D. NDRI (Deemed University), Karnal, Haryana, India.
54. Smith GM, Fitzhugh Jr HA, Cundiff LV, Cartwright TC, Gregory KE. A genetic analysis of maturing patterns in straightbred and crossbred Hereford, Angus and Shorthorn cattle. *Journal of Animal Science*. 1976; 43(2):389-395.
55. Stoop WM, Van Arendonk JAM, Heck JML, Van Valenberg HJF, Bovenhuis H. Genetic parameters for major milk fatty acids and milk production traits of Dutch Holstein-Friesians. *Journal of dairy science*. 2008; 91(1):385-394.
56. Swiger LA, Harvey WR, Eversson DU and Gregory KE. The variance of intraclass correlation involving group with one observation. *Biometrics*. 1964; 20:818-826.
57. Valsalan J, Sadan T, Anilkumar K, Aravindakshan TV. Estimation of co-variance components and genetic parameters of fertility and production traits in crossbred cattle of Kerala. *Theriogenology*. 2022; 181:126-130.
58. Vargas CA, Elzo MA, Chase Jr CC, Chenoweth PJ, Olson TA. Estimation of genetic parameters for scrotal circumference, age at puberty in heifers, and hip height in Brahman cattle. *Journal of animal science*. 1998; 76(10):2536-2541.
59. Verma UM, Kumar S, Yousuf S, Ghosh AK, Aswal APS. Genetic Evaluation of First Lactation Traits in Red Sindhi Cattle. *International Journal of Livestock Research*. 2018; 8(2):210-216.

60. Vinothraj S, Subramaniyan A, Venkataramanan R, Joseph C, Sivaselvam SN. (2016). Genetic evaluation of reproduction performance of Jersey× Red Sindhi crossbred cows. *Veterinary World*. 2016; 9(9):1012.
61. Wongpom B, Koonawootrittriron S, Elzo MA, Suwanasopee T. Milk yield, fat yield and fat percentage associations in a Thai multi-breed dairy population. *Agriculture and Natural Resources*. 2017; 51(3):218-222.
62. Zhu K, Li T, Liu D, Wang S, Wang S, Wang Q, Pan Y, Zan L, Ma, P. Estimation of genetic parameters for fertility traits in Chinese Holstein of south China. *Frontiers in Genetics*. 2024; 14:1288375.

UNDER PEER REVIEW