

**Field efficacy of selected chemicals and biopesticides against brinjal shoot and fruit borer(*Leucinodes orbonalis*) on brinjal (*Solanum melongena*)**

**Abstract :-**

A field experiment was carried out at the Central research farm of Sam Higginbottom University of Agriculture, Technology, and Sciences, located in Prayagraj, Uttar Pradesh, during the *kharif* 2023-24. The experiment was designed following a randomized block design with three replications. It encompassed eight treatments: T<sub>1</sub> - Neem oil 3% @ 30ml/lit, T<sub>2</sub>- *Metarhizium anisopliae*@2.5ml/lit, T<sub>3</sub> -*Bacillus thuringiensis* var. *kurstaki*@ 2gm/lit, T<sub>4</sub>- Spinosad 45SC @ 0.3 ml/lit, T<sub>5</sub> - Emamectin benzoate @ 0.4 gm/lit, T<sub>6</sub>-Chlorpyrifos 20EC@ 2.5ml/lit, T<sub>7</sub>- Chlorantraniliprole 18.5 SC @ 0.4ml/lit and T<sub>8</sub>- untreated control. The objective was to assess the field efficacy of selected chemicals and biopesticides against *Leucinodes orbonalis* (Guenee) infestation on brinjal in Prayagraj. Two rounds of spraying was conducted at 15-days interval. Data on shoot and fruit infestation was recorded after each spraying and picking, including the percentage of shoot infestation. The results showed that the initial population of the pest before the spray exhibited a non-significant distribution. Following the spray, the findings demonstrated that T<sub>5</sub> – Emamectin benzoate @ 0.4 gm/lit exhibited significant effectiveness against shoot and fruit borer, comparable to T<sub>7</sub> – Chlorantraniliprole 18.5 SC, at 3, 7, and 14 days post-spraying. Following closely in efficacy were Spinosad 45SC and Chlorpyrifos 20EC. The highest cost benefit ratio was recorded- T<sub>5</sub>-Emamectin benzoate, followed by T<sub>7</sub>-Chlorantraniliprole 18.5 SC. The highest cost-benefit ratio was recorded in T<sub>5</sub>– Emamectin benzoate @ 0.4gm/lit followed by T<sub>7</sub>- Chlorantraniliprole 18.5SC @ 0.4ml/lit and the lowest monetary return was observed with the control (T<sub>0</sub>) .

**Keywords:-** *Bacillus thuringiensis* var. *kurstaki*, R.B.D, *Leucinodes orbonalis*, brinjal shoot

**Introduction :-**

Brinjal (*Solanum melongena* Linn.), with a chromosome count of 2n=24, stands as one of the most beloved vegetables, also recognized by names like eggplant, aubergine, or guinea squash. Belonging to the nightshade family Solanaceae, it holds the prestigious title of "King of the Vegetables". Renowned for its high yielding capabilities, brinjal can thrive year-round across a

variety of agro-climatic conditions, particularly in tropical and sub-tropical regions. The Indian sub-continent serves as the center of origin for this esteemed vegetable(**Omprakash and Raju 2014**).

In India, brinjal cultivation spans approximately 0.743 million hectares of agricultural land, yielding a production of 12.77 million tonnes annually, and a productivity of nearly 17.17 MT/ha(**Anonymous,2022**).The primary brinjal cultivating states in India include Bihar, Odisha, Karnataka, Andhra Pradesh, Maharashtra, West Bengal, Uttar Pradesh, and states with climatic conditions conducive to tropical and subtropical cultivation. In Karnataka, brinjal cultivation covers an area of 1.58 lakh hectares, contributing to a production of 402.5 MT, accounting for a 3.13% share, with a productivity of 25.4 MT/ha(**Anonymous,2016**). Brinjal has been acknowledged in Ayurveda for its therapeutic potential in managing diabetes. Additionally, it is esteemed for its diverse medicinal properties, acting as a beneficial appetizer, aphrodisiac, cardiac tonic, laxative, and inflammation reliever. Moreover, it serves as an excellent remedy for liver-related health issues(**Lalita and Kashyp,2020**).

The year-round availability of brinjal exposes the crop to a spectrum of biotic and abiotic factors. Among these, insect pests emerge as crucial contributors to yield reduction, as they assail the crop from its nursery stage through harvesting. Brinjal faces attacks from approximately 142 species of insect pests, along with four species of mites and nematodes, across various regions worldwide(**Jat and Shrivastva,2023**). Brinjal crops are susceptible to numerous insect pests, including aphids (*Aphis gossypii* Glover), whiteflies (*Bemisia tabaci* Lind.), jassids (*Amrasca biguttula biguttula* Ishida), spotted leaf beetles (*Epilachna vigintioctopunctata* Fab.), brinjal shoot and fruit borers (*Leucinodes orbonalis* Guenee), brinjal leaf beetles (*Psylliodes bali* Jacoby), and leaf folders (*Eublemma oleracea* Walk.)(**Patra et al,2016**).

Among these pests, the brinjal shoot and fruit borer, *Leucinodes orbonalis* (Guenee) (Lepidoptera: Crambidae), poses a significant and destructive threat to brinjal production(**Singh et al, 2019**). During the early stages of crop growth, adult female moths predominantly lay eggs on the lower side of young leaves near the midrib, occasionally on tender shoots. Upon hatching, young larvae bore into the young leaves near the midrib or tender shoots, sealing the opening with frass and feeding within the shoot or midrib of the leaves. Symptoms such as drooping,

wilting, or withering of shoots are commonly observed during the initial stages of crop growth, indicating shoot damage. As the crop matures and fruits begin to form, larvae typically penetrate through the underside of the calyx, bud, or fruit, sealing the entry point with frass. Infestation of buds often leads to flower drop, further exacerbating the impact of the pest on crop yield. The holes observed on the fruits are actually exit holes of the larvae. Such infested fruits are partially unfit for human consumption and fetch lower prices in the market(Shigaonkar et al,2022). This pest inflicts damage on brinjal crops, leading to yield losses ranging from 60-80 percent, or even causing complete damage if no control measures are implementedThakare et al, (2021).

### **Justification :-**

In recent years, a diverse array of insecticides from different chemical families have been employed to combat pests. However, over-reliance on these chemicals has given rise to several issues including resistance, resurgence of pest populations, environmental contamination, and potential health risks for consumers. Consequently, there is an urgent need to judiciously administer insecticides at optimal dosages to mitigate environmental damage and safeguard human health. To address these challenges and minimize pest infestations, a combination of botanical extracts and insecticides from various chemical groups are recommended for managing the shoot and fruit borer (*Leucinodes orbonalis*). Continuous utilization of chemical pesticides poses significant threats to natural ecosystems, including environmental pollution, the development of pest resistance, and potential health hazards. To address these concerns, exploring the insecticidal properties of plant-derived products against the shoot and fruit borer (*Leucinodes orbonalis*) on brinjal is imperative. This study aims to elucidate the effectiveness of different insecticidal formulations, shedding light on their outcomes. Through rigorous analysis, the most effective treatments can be identified and implemented, offering sustainable solutions for pest management in brinjal cultivation.

### **MATERIALS AND METHODS :-**

The investigation was conducted at the experimental field of Sam Higginbottom university of agriculture, Technology and Sciences, Prayagraj, Uttarpradesh during the *kharif* 2023-24.

### **Experimentalsite:**

The present investigation was conduct at the Central research farm of “Sam Higginbottom

University of Agriculture, Technology and Sciences” Prayagraj, Uttar Pradesh during *Kharif* season 2023-24. The research farm is situated on the right side of Rewa road at 25° 28' 22.9224" North, 81° 52' 42.0852" East longitude and is about 129.2 cm above sea level.

**Sowing of seed on raised bed:**

Theseedsof brinjalvariety**Supriya**wasusedtoraiseseedlinginnursery.Regularwateringand weedingwasdoneupto transplanting of seedling to the main field.

**Application of fertilizers:**

Chemical fertilizers was applied @ NPK 50:25:25 kg/ha and 20 tonnes FYM per hectares. Full P, K, 33.3% N and FYM was applied as basal dose in the drills before sowing the seed and rest of the nitrogen was top dressed in two equal splits at 21 and 41 days after transplanting.

**Intercultural operations:**

Weedingoperationswascarriedouttoconserveoilmoistureandtokeeptheexperimentalfieldfreefrom the weeds.Weeding doneat20 days,40daysand60(DAT).

The data were subsequently converted into percentages of infestation utilizing specific formulas.-

**On Shoot :-**

**Number basis**The total number of shoots and the number of infested shoots from five selected plants in each treatment replication was recorded(**Soulakhe et al, 2021**).

$$\% \text{Shoot infestation} = \frac{\text{No. of shoots infested}}{\text{Total no. of shoots}} \times 100$$

**On Fruit :-**

**Number basis** During each picking, the total number of fruits and the number of infested fruits from five selected plants in each treatment replication was recorded (**Gowrish et al, 2015**).

$$\text{Fruit infestation} = \frac{\text{No. of fruits infested}}{\text{Total no. of fruits}} \times 100$$

**Cost benefit ratio of treatments:**

Gross returns was calculated by multiplying total yield with market price of the produce. Cost of

cultivation and cost of treatments was deducted from the gross returns, to find out returns and cost benefit of ratio by following formula,

Net returns

BCR=-----

Costoftreatment

Where,

BCR=Benefit Cost Ratio

### Statistical Analysis:

The data averaged into respective parameter requisite was subjected to suitable transformation. After analysis, data was accommodated in the table as per the needs of objectives for interpretation of results. The standard procedures in agriculture statistics given by **Gomez and Gomez (1976)** were consulted throughout. The interpretation of data was done by using the critical difference value calculated at 0.05 probability level. The level of significance was expressed at 0.05 probability. The F-test were used to determine the significant difference.

### List 1: Dates of Spray Application:

Sr. No.	Spray application	Date of application
1	<b>Fist spray</b>	25/09/2023
2	<b>Second spray</b>	10/10/2023

### Results and Discussions :-

#### Field efficacy of selected chemicals and biopesticide against brinjal shoot and fruit borer (*Leucinodes orbonalis* Guenee), infesting brinjal.

The data concerning the percentage of infestation by shoot and fruit borers after the first and second spray indicated that all chemical treatments exhibited significant superiority over the control. Among all treatments, the lowest percentage of infestation by shoot and fruit borers was observed in Emamectin benzoate (T<sub>5</sub> - 7.94), followed by Chlorantraniliprole (T<sub>7</sub> - 8.93), Spinosad (T<sub>4</sub> - 10.12), Chlorpyrifos (T<sub>6</sub> - 10.82), *Metarhizium anisopliae* (T<sub>2</sub> - 11.57), Neem oil (T<sub>1</sub> - 12.46), *Bacillus thuringiensis* var. *Kurstaki* (T<sub>3</sub> - 11.57), and the control (T<sub>0</sub> - 26.90).A

comprehensive analysis revealed that all the biopesticides were effective, with the following decreasing order of efficacy: Emamectin benzoate > Chlorantraniliprole 18.5SC > Spinosad 45SC > Chlorpyrifos 20EC > *Metarhizium anisopliae* > Neem oil 3% > *Bacillus thuringiensis* var. *Kurstaki*. Their significant impact on reducing shoot infestation was evident when compared against the untreated control. These findings align closely with previous studies conducted by **Mane and Kumar (2020)**, **Verma et al,(2021)** and **Shyamrao et al,(2018)** which also highlighted the superiority of Emamectin benzoate in reducing the population of shoot and fruit borer.


Table 1: Field efficacy of selected chemicals and bio-pesticides against *Leucinodes orbonalis* (Guenee) on brinjal. : (overall mean, yield, B:C ratio)

TREATMENTS		Dose (gm/ml/L)	%Shootinfestation/5plants								Overall mean	Yield (q/ha)	B:C Ratio	
			1DBS	3DAS	7DAS	14DAS	Mean	3DAS	7DAS	14DAS				Mean
T <sub>1</sub>	Neemoil@ 3%	30ml	20.94	15.69	13.75	14.21	14.55	12.05	9.05	10.04	10.38	12.46	72.3	1:3.2
T <sub>2</sub>	<i>Metarhizium anisopliae</i>	2.5ml	20.38	14.88	13.24	13.76	13.96	10.72	7.89	8.95	9.18	11.57	75	1:5.0
T <sub>3</sub>	<i>Bacillus thuringiensis</i> var. <i>kurstaki</i>	2gm	21.29	16.35	14.37	15.24	15.32	13.50	10.01	10.79	11.43	13.37	69.5	1:4.7
T <sub>4</sub>	Spinosad45S C	0.3ml	21.35	13.55	11.82	12.56	12.64	9.15	6.61	7.01	7.60	10.12	85.12	1:5.3
T <sub>5</sub>	Emamectin benzoate5%	0.4gm	21.75	11.21	8.98	9.73	9.97	7.01	5.14	5.59	5.91	7.94	90	1:6.0
T <sub>6</sub>	Chlopyrifos 20EC	2.5ml	21.13	14.28	12.76	13.13	13.39	9.81	7.38	7.61	8.26	10.82	79.4	1:5.3
T <sub>7</sub>	Chlorantraniliprole18.5SC	0.4ml	20.60	12.74	10.79	11.34	11.62	7.51	5.45	5.77	6.24	8.93	87.5	1:5.5

T <sub>0</sub>	Controls	-	21.58	24.19	25.51	26.02	25.24	27.69	28.30	29.73	28.57	26.90	44	1:3.1
	F-test		<b>NS</b>	<b>S</b>	<b>S</b>	<b>S</b>	<b>S</b>	<b>S</b>	<b>S</b>	<b>S</b>	<b>S</b>	<b>S</b>	-	-
	S. E(d)(±)		-	0.50	0.38	0.78	0.99	0.50	0.39	0.70	0.71	1.17	-	-
	CD (5%)		-	1.29	0.78	1.42	1.14	1.07	0.77	0.85	1.41	2.42	-	-

\*Figures in parentheses are arc sin transformation values, DAS - days after spray, NS - non significant, S - significant, CD 5% level – Critical differences at 5% level of significances

UNDER PEER REVIEW



**Fig1:-Overall mean of brinjal shoot and fruit borer infestation at the 1<sup>st</sup> and 2<sup>nd</sup> spray**

### Yield-

The data revealed that the highest grain yield of 90 q/ha was attained with Emamectin benzoate (T<sub>7</sub>) support by **Prasad et al, (2018)**, followed by Chlorantraniliprole 18.5 SC (T<sub>7</sub>) with 87.5 q/ha support by **Jalali et al, (2012)**, Spinosad 45SC (T<sub>4</sub>) with 85.12 q/ha support by **Chakraborty et al,(2019)**, Chlorpyrifos 20EC (T<sub>6</sub>) with 79.4 q/ha support by **Singh et al, (2015)**, *Metarhizium anisopliae* (T<sub>2</sub>) with 75 q/ha support by **Kumar and Singh (2013)**, Neem oil 3% (T<sub>1</sub>) with 72.3 q/ha support by **Anitha and Chakravarthy (2018)**, and *Bacillus thuringiensis* var. *kurstaki* (T<sub>3</sub>) with 69.5 q/ha support by **Srinivas and Ramachandramurthy (2016)**. The untreated control plot (T<sub>0</sub>) recorded the lowest yield of 44 q/ha.

In this study, Emamectin benzoate (T<sub>5</sub>) exhibited the most promising results among all treatments, followed by Chlorantraniliprole (T<sub>7</sub>), in reducing crop infestation. Spinosad 45SC (T<sub>4</sub>), Chlorpyrifos 20EC (T<sub>6</sub>), *Metarhizium anisopliae* (T<sub>2</sub>), and Neem oil 3% (T<sub>1</sub>) also demonstrated effectiveness against the Brinjal shoot and fruit borer [*Leucinodes orbonalis* (Guenee)].

### Cost Benefit Ratio –

The analysis of the Cost-Benefit Ratio for all treatments revealed that the highest monetary return was achieved with Emamectin benzoate (T<sub>5</sub>) (1:6.0) support by **Singh and Verma, (2018)**, followed by Chlorantraniliprole 18.5SC (T<sub>7</sub>) (1:5.5) support by **Sharma and Yadav, (2019)**, Spinosad 45SC (T<sub>4</sub>) (1:5.3) support by **Thakur and Kumar, (2017)**, Chlorpyrifos 20EC (T<sub>6</sub>) (1:5.3) support by **Gupta and Sharma, (2016)**, *Metarhizium anisopliae* (T<sub>2</sub>) (1:5.0) support by **Singh and Singh, (2018)**, Neem oil 3% (T<sub>1</sub>) (1:3.2) support by **Singh and Singh, (2015)**, and *Bacillus thuringiensis* var. *kurstaki* (T<sub>3</sub>) (1:5.0) support by **Kumar and Jha, (2019)**. The lowest monetary return was observed with the control (T<sub>0</sub>) (1:3.1).

### Conclusion-

The data on the percent infestation of shoot and fruit borer after the first and second sprays revealed that all the chemical treatments were significantly superior to control. Among all the treatments, the lowest percent infestation of shoot and fruit borer was recorded in Emamectin benzoate (7.94) with highest yield 90q/ha and cost benefit ratio of (1:6.0). The highest infestation of in untreated control (26.90) with the lowest cost benefit ratio (1:3.1).

### References :-

1. **Omprakash S and Raju SVS.** A brief review on abundance and management of major insect pests of brinjal (*Solanum melongena* L.). *International journal of applied biology and pharmaceutical technology*, 2014;5(1): 228-238.
2. **Anonymous.** Area, production and productivity of brinjal in Karnataka and India (2021-2022-1st advance estimates), *Indiastat.com*; 2022.
3. **Anonymous.** *National Horticultural Mission* ; 2016.<https://nhb.gov.in/>
4. **Lalita S and Kashyap L.** Biology and mechanisms of resistance to brinjal shoot and fruit borer: A review. *Journal of Entomology and Zoology Studies*, 2020;8(2):2111-2118.
5. **Jat HK and Shrivastva VK.** Cost: Benefit analysis of newer insecticides used in control of brinjal shoot and fruit borer (*Leucinodes orbonalis* Guenee) at Gwalior region

of Madhya Pradesh. *Journal of pharmaceutical innovation*, 2023;12(3): 3091-3093.

6. **Patra S, Thakur NS and Firake DM.** Evaluation of bio-pesticides and insecticides against brinjal shoot and fruit borer (*Leucinodes orbonalis* Guenee) in Meghalaya of North-Eastern India. *International Journal of Bio-resource and Stress Management*, 2016;7(5): 1032-1036.
7. **Singh D, Kumar D, Singh R, and Sharma A.** Management of brinjal shoot and fruit borer (*Leucinodes orbonalis*) through different eco-friendly management practices. *Journal of Entomology and Zoology Studies*, 2019;7(1), 108-112.
8. **Shigaonkar RS, Shinde BD, Shelke SB, Chopkar PS, Durge SM and Choudhari RJ.** To screen some brinjal cultivars against shoot and fruit borer, *Leucinodes orbonalis*. *Journal of Pharmaceutical Innovation*, 2022;11(1):1337-1341.
9. **Thakare VS, Undirwade DB, Kulkarni US and Ghawade SM.** Screening of brinjal genotypes for resistant reaction against brinjal shoot and fruit borer (BSFB) *Leucinodes orbonalis* Guenee. *Journal of Entomology and Zoology Studies*, 2021;9(1):1653-1657.
10. **Soulakhe SD, Bantewad NE and Jayewar NE.** Biorational management of brinjal shoot and fruit borer *Leucinodes orbonalis*. *Pharma Innov. J.*, 2021;10(8):617-623.
11. **Gowrish BR, Ramesha B and Ushakumari R.** Biorational management of major pest of brinjal. *Indian J Entomol*, 2015;77(1):51-55.
12. **Singh M and Sachan SK.** Comparative efficacy of biopesticides against shoot and fruit borer, *Leucinodes orbonalis* Guenee in brinjal. *Plant Arch.* 2015;15(2):805-808.
13. **Sharma JH and Tayde AR.** Evaluation of bio-rational pesticides against brinjal fruit and shoot borer, *Leucinodes orbonalis* Guen. On brinjal at Allahabad Agroclimatic region. *Int. J Curr. Microbiol. Appl. Sci.* 2017;6(6):2049-2054.
14. **Shyamrao NJ, Kumar A, Patil AA and Narode MK.** Efficacy of certain insecticide and bio-pesticide against brinjal shoot and fruit borer, *Leucinodes orbonalis* Guenee in brinjal. *Journal of Entomology and Zoology Studies*, 2018; 6(5): 292-295.
15. **Mane PD and Kumar R.** Bio-efficacy of new chemicals against shoot and fruit borer of brinjal. *Journal of Entomology and Zoology Studies*, 2020; 8(6): 1692-1693.
16. **Verma AP, Chandra U, Shakya A, Singh RP, Singh B and Alam K.** Efficacy of bio-insecticide against brinjal shoot and fruit borer, *Leucinodes orbonalis* Guenee. *The Pharma Innovation*, 2021; SP-10(12): 1071-1077.

- 17. Prasad YG, Ramasubba Reddy PP, and Kadiyala MDM.** Evaluation of newer molecules against shoot and fruit borer, *Leucinodes orbonalis* Guenee (Lepidoptera: Crambidae) infesting brinjal. *Journal of Pharmacognosy and Phytochemistry*,2018; 7(2), 2982–2985.
- 18. Jalali SK, Seal DR, Das SK and Venugopal MS.** Relative toxicity of insecticides to eggplant fruit and shoot borer, *Leucinodes orbonalis* Guenee (Lepidoptera: Pyralidae) and their natural enemies. *Journal of Biopesticides*,2012; 5(1), 26–32.
- 19. Chakraborty S and Mandal B.** Field efficacy of newer insecticides against eggplant fruit and shoot borer, *Leucinodes orbonalis* Guenee in West Bengal. *International Journal of Chemical Studies*, 2019; 7(2), 1818–1821.
- 20. Singh VK and Sahoo S.** Evaluation of newer insecticides against eggplant shoot and fruit borer, *Leucinodes orbonalis* (Guenee) under field condition. *Indian Journal of Entomology*, 2015; 77(4), 344–346.
- 21. Kumar S and Singh D.** Efficacy of different bio-pesticides against eggplant shoot and fruit borer, *Leucinodes orbonalis* Guenee. *Biopesticides International*,2013; 9(1), 1–7.
- 22. Anitha K and Chakravarthy AK.** Evaluation of botanicals against eggplant shoot and fruit borer, *Leucinodes orbonalis* (Guen.). *Journal of Entomology and Zoology Studies*, 2018; 6(2), 56–59.
- 23. Srinivas T and Ramachandramurthy V.** Field evaluation of newer insecticides against eggplant shoot and fruit borer, *Leucinodes orbonalis* Guen. *Journal of Entomology and Zoology Studies*, 2016; 4(1), 232–234.
- 24. Singh RK and VermaA.** Economics of management practices for brinjal shoot and fruit borer, *Leucinodes orbonalis* Guenee. in Bihar. *Journal of Pharmacognosy and Phytochemistry*,2018; 7(1), 1031–1034.
- 25. Sharma S and Yadav R.**Economics of management practices against brinjal shoot and fruit borer (*Leucinodes orbonalis* Guen.). *Indian Journal of Economics and Development*, 2019; 15(4), 676–680.
- 26. Thakur A and Kumar V.** Economics of different management practices against *Leucinodes orbonalis* (Guen.) infesting brinjal. *Progressive Agriculture*, 2017; 17(1), 128–132.

27. **Gupta P and Sharma H.**Economics of brinjal shoot and fruit borer, *Leucinodes orbonalis* Guenee management practices. *Economic Affairs*,2016; 61(1), 141–145.
28. **Singh D and Singh R.** Economics of biopesticides against brinjal shoot and fruit borer, *Leucinodes orbonalis* Guenee in Bihar. *Journal of Pharmacognosy and Phytochemistry*, 2018; 7(5), 330–332.
29. **Singh J and Singh V.** Economics of management practices against brinjal shoot and fruit borer, *Leucinodes orbonalis* Guen. infestation. *International Journal of Agriculture Sciences*, 2015; 7(35), 4022–4024.
30. **Kumar A and Jha MK.**Economic analysis of management practices against brinjal shoot and fruit borer (*Leucinodes orbonalis* Guen.) infestation. *Asian Journal of Agricultural Extension, Economics & Sociology*,2019; 35(1), 1–7.
31. Verma , N. S., Kuldeep , D. K., Chouhan , M., Prajapati , R., & Singh, S. K. (2023). A Review on Eco-Friendly Pesticides and Their Rising Importance in Sustainable Plant Protection Practices. *International Journal of Plant & Soil Science*, 35(22), 200–214. <https://doi.org/10.9734/ijpss/2023/v35i224126>
32. Lal , B., & Dhurve , V. (2024). Bio-efficacy of Different Biopesticides against Jassid, (*Amrasca biguttula biguttula* Ishida) Infesting Okra [*Abelmoschus esculentus* (L.) Moench]. *Journal of Experimental Agriculture International*, 46(1), 37–47. <https://doi.org/10.9734/jeai/2024/v46i12289>
33. Yadav IC, Devi NL, Syed JH, Cheng Z, Li J, Zhang G, Jones KC. Current status of persistent organic pesticides residues in air, water, and soil, and their possible effect on neighboring countries: A comprehensive review of India. *Science of the Total Environment*. 2015 Apr 1;511:123-37.