

Impact of foliar application of boron on yield and yield attributes, assimilate supply, production efficiency and fertilizers productivity of Indian mustard (*Brassica juncea*) in field experiments

ABSTRACT: Field experiments were conducted for three consecutive *rabi* seasons from 2018 to 2020 to assess the impact of foliar application of boron fertilizer on yield and yield attributes, assimilate supply, production efficiency and fertilizers productivity of Indian mustard [*Brassica juncea*(L.) Czern. & Coss.]. The results showed that the number of primary branches/ plant (7.00 to 8.08), number of siliquae/plant (427.2 to 468.8), number of seeds/siliqua (13.03 to 14.01), test weight of seed (3.60 to 3.80 g) and plant dry matter production (38.10 to 41.20 g/plant) were significantly increased with T2 treatment (100% NPKS + foliar application of Boron @ 0.5 % Borax) as compared to farmer's practice T1 (100% NPKS + without boron) during three years of experiments. Significantly higher seed yield (22.5 to 24.7 q/ha) of the mustard crop was also reported with T2 (100% NPKS + foliar application of Boron @ 0.5 % Borax) as compared to farmers' practice T1 (100% NPKS + without boron) (seed yield 19 to 22.5 q/ha) during the studies. Similarly, Treatment T2 resulted in significantly higher production efficiency and assimilates supply of mustard crop than that of T1 (farmer's practice). The fertilizer productivity in the mustard crop that is directly related to seed yield and the partial factor of productivity of N, P, K and S fertilizers (PFP_{NPKS}) were also increased significantly with the T2 compared to farmers practice T1 (farmer's practice), during 3 years of studies. The highest gross return and net return and benefit-cost ratio were obtained with T2 (100% NPKS + foliar application of Boron @ 0.5 % Borax) treatment over T1 (100% NPKS + without boron).

Keywords: Assimilate supply, Economics, Fertilizers productivity, Production efficiency, and Seed yield

1. INTRODUCTION

Rapeseed-mustard is the important oilseed crop in the World, and India is the third largest producer of rapeseed-mustard after Canada and China. Mustard is the second most important oilseed crop in India after peanut with regard to area and production

(Pramanick, *et al.*, 2023). In India, the area of rapeseed-mustard is 8.85 million hectares, and the total production is 12.81 mt in 2022–23. The average productivity of Rapeseed-mustard crop for the year 2022-23 was 1447 kg/ha (GOI, 2022). Boron is an essential micronutrient for vascular plants and necessary for the growth of crop plants (Wimmer *et al.* 2019). Low availability of boron in soils impairs the development of floral organs and the elongation of pollen tubes, which leads to a loss of crop yield (Zhao *et al.* 2021). Boron deficiency is the second most important micronutrient constraint in crop production after zinc (Rehman *et al.* 2022). Rapeseed–mustard is extremely sensitive to boron (B) deficiencies (Zhao *et al.* 2021). Boron deficiency in soil leads to the erratic growth of seedlings and reduced photosynthesis (Safdar *et al.* 2023). Boron is an important component for various biological processes in plants, including the growth and development of pollen tubes, the maintenance of membrane integrity, pollination and seed production (Ahmad *et al.* 2012). The primary roles of boron include the breakdown of nucleic acids, carbohydrates, proteins, indole acetic acid, and phenol, which are involved in the synthesis of plant cell walls and the maintenance of membrane integrity (Goldbach *et al.* 2001). Additionally, boron plays a key role in cellular division and the control of carbohydrate and protein metabolism, and these processes influence the reproductive phase and development of seeds (Miwa *et al.* 2010). Boron is closely associated with plant growth and plays a vital role in cell division as well as synthesis of oil and protein. Flowering and pods formation is restricted due to boron deficiency and rapeseed-mustard crops are very responsive to boron fertilizer (Islam 2005, and Saha *et al.* 2003). Yield reduction due to boron deficiency often occurs even if specific visual symptoms of leaves are not apparent. Keeping in mind the problem of low yield of mustard due to boron deficiency in the soil, an on farm trial was designed on foliar application of boron fertilizer in mustard to reduce the yield loss due to the deficiency of boron and to obtain higher yield and quality of mustard crop.

2. MATERIAL AND METHODS

On -farm trials on foliar application of boron on Indian mustard were carried out at the farmer's fields of Qazipur, Shikarpur and Jhatikara villages of NCT Delhi under

similar agro climatic conditions during the *rabi* season from 2018 to 2020 (Figure 6). The rainfall and average temperature of the experimental area from 2018 to 2021 have been depicted in Figure 3,4,5. The experimental soil was a sandy loam in texture, low in organic carbon (0.31%), available nitrogen (225 Kg/ ha), phosphorous (21 Kg/ ha), potash (195 Kg/ha). The experiments were laid out on five farmer's fields with 5 replications in a 1000 m² plot area each and having 2 treatments *i.e* T1 Farmer's practice (100% NPKS + without boron) and T2 (100% NPKS + foliar application of Boron @ 0.5 % Borax). The recommended dose of fertilizer was 80:60:40: 30 kg N, P, K,S ha⁻¹, respectively, for Indian mustard. Nutrients were supplied through urea, single super phosphate, muriate of potash and bentonite sulphur. Boron was supplied through borax which contains 11% boron through foliar application. Whole of P, K and S and half of the N were supplied as basal dose and the remaining half of the nitrogen was applied 30 days after sowing. Proper weed management, insect- pest management and irrigation management was done in experimental fields. The Number of primary branches and total siliquae/ plant was counted from five randomly selected plants at harvest and the mean value was taken. The numbers of seeds per siliquae is also counted from randomly selected siliquae for the selected plants. Four plants were selected randomly from the observational rows of each plot and dug out with the help of *khurpi* at 60 DAS, 90 DAS and at harvest. These plants were sun-dried for 48 hours. After sun drying, these plants were dried in the oven at 65±5°C temperature for 48-72 hours or till the samples attained a constant weight, and weighed. The dry matter was expressed in g plant⁻¹. The data output was collected from experimental fields as well as farmer's and cost of cultivation, net income, and benefit -cost ratio were also worked out. The yield data were collected from both the experiment and farmers' practice and analyzed by using simple statistical tools. All data were analyzed were using ANOVA, and two sample test values at 5% level of significance were calculated and used to test significance difference between treatment means. Partial factor productivity of nitrogen (PFP_N), phosphorus (PFP_P) and Potash (PFP) fertilizer was calculated as per the following formula (Singh *et al.* 2021).

$$\text{Partial factor of productivity (PFP}_{\text{npk}}) = \frac{\text{Seed yield (kg)}}{\text{Rate of fertilizer applied (kg)}}$$

Assimilate supply (AS) is the ratio of above ground dry matter of individual plant / number of siliquae. Production efficiency (PE), it represents the increase in seed yield on a daily basis and calculated by dividing total mustard seed production ha⁻¹ in a sequence with total duration of the crops in a sequence (Premi *et al.* 2013).

3. RESULTS AND DISCUSSION

3.1 Yield attributes

Foliar application of boron had a significant impact on yield attributes of the mustard crop such as number of primary branches per plant, number of siliquae/plant, and number of seeds/siliqua and 1000-seed weight during 3 years of studies (Table 1). Significantly higher number of primary branches/ plant (7 to 8.08), number of siliquae/plant (427.2 to 468.8), number of seeds/siliqua (13.03 to 14.01) were obtained with T₂ treatment (100% NPKS + foliar application of Boron @ 0.5 % Borax) as compared to farmer's practice T1 (100% NPKS + without boron) during three years of experiments. Similarly, the higher test weight of seed (3.6 to 3.8 g) was reported with the with T₂ treatment (100% NPKS + foliar application of Boron @ 0.5 % Borax) in mustard crop, which was significantly higher over farmer's practice T1 (100% NPKS + without boron) in which test weight of seed was 3.2 to 3.3 g. The partial or complete failure of boron fertilization leads to pollen abortion, seed drop, deformed and shrunken seeds and partially filled pods resulting in lower test weight (Kapila *et al.* 2008) and (Kumararaja *et al.* 2015). Jankowskiet.al (2020) also reported the significantly higher primary branches/ plant, number of siliquae/plant, and number of seeds/siliqua and 1000-seed weight with foliar application of boron in mustard crop. The highest values of yield attributes in treatment T2 might be attributed to the involvement of boron in hormone synthesis and translocation, carbohydrate metabolism and the synthesis of DNA in plants (Verma *et al.* 2020). Identical results for plant growth parameters were reported by Shah *et al.* (2016) and Handiganooret al. (2017).

3.2 Plant dry matter production

Foliar application of boron had a notable impact on dry matter accumulation during 3 years of the studies (table 2). The dry matter production (38.1 to 41.2 g/plant) in mustard crop was significantly higher with T₂ treatment (100% NPKS + foliar application of Boron @ 0.5 % Borax) as compared to the farmer's practice T₁ (100% NPKS + without boron), in which an increase of 36.4 to 38 g/plant dry matter was recorded. The dry matter production was found to be increased could be attributed to the increased number of branches per plant arising out of the better growth and development conditions facilitated by proper supply of photosynthates to the plants parts. Boron is needed in small quantity by plants for division and cell enlargement in meristem tissue, then the meristem generate new cells at the tip of the root or stem that resulted in increased plant growth or length. The increase growth of plant and number of leaf affects the light received by the plant so that it can improve the performance of photosynthesis and produce many photosynthase in forms of starch, lipids, and proteins. Photosynthase is translocated inside the plant which helps to increase the dry matter of the plant (Timotwuet *al.* 2018).

3.3 Seed yield

The seed yield of the mustard crop was influenced by foliar application of boron during 3 years of studies. The significantly higher seed yield (22.50 to 24.70 q/ha) of the mustard crop was reported with T₂ treatment (100% NPKS + foliar application of Boron @ 0.5 % Borax) as compared to the farmer's practice T₁ (100% NPKS + without boron), in which the seed yield was 19.00 to 22.50 q/ha during the experiments (Table 2). These results corroborated well with the studies conducted by Jankowski *et al.* 2020 and Zhao *et al.* (2021) in which they reported higher seed yield with boron in mustard field. Lu *et al.* (2000) and (Randhawa *et al.* 2021) found that boron fertilizer contributed 611 kg/ha (48.5%) yield advantage of the rapeseed. The increase in the seed yield can be ascribed to boron as it is directly linked with the process of fertilization, pollen producing capacity of anther, viability of pollen grains, pollen germination and

pollen tube growth (Padhbhusan and Kumar, 2015) and (Jaiswal *et al.* 2015). Boron's involvement in maintaining the structural integrity of plasma membrane, metabolism of carbohydrates, and synthesis of DNA probably resulted in the additional growth in crops (Handiganoore *et al.* 2017). The reason for the lowest seed yield under farmers practice yield might be the higher pollen infertility and lower seed filling as they play an extremely crucial role in both the processes (Hussain *et al.* 2012). Correlation analysis showed that seed yield had a significant positive correlation with number of siliquae, seeds/ siliqua, test weight, dry matter accumulation, assimilate supply, however, negative correlation was observed with primary branches (table 4).

3.4 Production efficiency

The production efficiency of a crop refers to the daily increase in seed yield. Foliar application of boron exerted significant effects on the production efficiency of mustard crop during 3 years of studies (referencing Table 2). Specifically, as a result of foliar application of boron (T₂ treatment), the production efficiency of the mustard crop became significantly higher (16.67 to 18.31) than that of the farmer's practice T₁ (100% NPKS + without boron), whose production potential was found to be 14.07 to 16.67. Foliar application of boron helps in proper translocation of photosynthates toward the seeds and siliquae formation, which enhances in the seeds and siliquae production in mustard. Whereas, the seeds and siliquae development was poor under farmers practices in which boron fertilizer was not applied. The translocation of food material from source to sink was better in boron spray treatment as compared to farmer's practices.

3.5 Assimilate Supply

Assimilate supply was influenced significantly with the application of boron fertilizer has been depicted in table 2. Assimilate supply refers to a plant's ability to mobilize photosynthates once again and it is the ratio of above ground dry matter of individual plant to the number of siliquae formation. During the 3 years studies, assimilate supply of the mustard crop was increased significantly

(11 to 11.76) with T₂ treatment (100% NPKS + foliar application of Boron @ 0.5 % Borax) over farmer's practice T₁ (100% NPKS + without boron) in which it was increased up from 10.33 to 11.17. At the time of the peak flowering period, the nutrients and carbohydrates begin to remobilize toward seed setting and siliquae formation. Similar findings were also confirmed by Shekhawat *et al.* (2016). Application of boron might be involved in the synthesis of protein, chloroplast pigments, and electron transfer system that led to the increased photosynthetic activity as well as assimilate supply of mustard crop (Srinivasan *et al.* 2019). The positive effect of boron application has been reported on assimilate supply by Haque *et al.* (2000) and Islam (2005).

3.6 Fertilizer productivity

Foliar spray of boron had a significant impact on NPK & S fertilizers productivity during 3 years of studies. Fertilizer productivity was evaluated in terms of partial factor of productivity (PFP) and calculated based on the yield of the mustard crop (Table 3). Over the course of study years, the PFP of N, P, K and S fertilizers (PFP_{NPKS}) were increased significantly (28.13 to 30.9, 37.5 to 41.19, 56.25 to 61.79, 75 to 82.39 kg kg⁻¹) with T₂ treatment (100% NPKS + foliar application of Boron @ 0.5 % Borax) as compared to the farmer's practice T₁ (100% NPKS + No use of boron). It was observed that the PFP of S was found to be higher followed by PFP of K over PFP of N and P. It means boron had a positive interaction with other nutrients like N, P, K and S, and plays an important role to increase the fertilizer productivity of NPK&S. Higher PFP of NPK&S with a foliar application of boron might be ascribed for higher seed yield production and better translocation of food materials toward seeds development than without boron application practice. All these are in accordance with the findings reported by Singh *et al.* (2021) and (Zhao *et al.* 2021).

3.7 Economics

The economics of the treatments was calculated depending on the prevailing market prices of the inputs and outputs for the particular year. During 3 years of

studies, the higher gross return of Rs. 85500 to 111600/ha, net return of Rs.63210 to 72828/ha and B: C ratio 2.83 to 4 were obtained with T₂ treatment (100% NPKS + foliar application of Boron @ 0.5 % Borax) as compared to farmer's practice T₁ (100% NPKS + without boron) has been illustrated in figure 1. The higher benefit- cost ratio with T₂ treatment was found due to the higher yield obtained under T₂ treatment as compared to T₁ (farmer's practice) (figure 2). Hence, higher benefit- cost ratio proved the economic viability of the technology intervention and convinced the farmers about the utility of improved technologies. Similar economic benefits owing to adoption of foliar application of boron in mustard were also reported by Dubey *et. al* (2018).

4. CONCLUSION

Based on the results of on-farm trials conducted in real farming conditions, it can be concluded that the foliar application of boron, specifically at a concentration of 0.5% borax alongside 100% recommended NPKS fertilizers, enhances the yield attributes of the mustard crop. Additionally, the foliar application of boron has been shown to improve the production efficiency and assimilate supply of the mustard crop. Moreover, the fertilizer productivity of N, P, K, and S fertilizers was observed to increase with the foliar application of boron at the mentioned concentration over the course of three years of studies. Consequently, foliar application of boron at 0.5% borax concentration was identified as optimal for achieving higher seed yields in boron-deficient soils. The adoption of foliar application of boron has the potential to significantly augment both the income and livelihoods of farming communities.

REFERENCES

- Ahmad, W., Zia, M.H., Malhi, S.S., Niaz, A. and Ullah, S., 2012. Boron Deficiency in soils and crops: a review. *Crop plant*, 2012, pp.65-97.
- Dubey, S.K., Gautam, U.S., Singh, A.K., Singh, A., Chahal, V.P., Singh, A.K., Singh, C.H.A.N.D.A.N. and Srivastava, A.J.I.T., 2018. Quantifying the yield gap minimization in lentil (*Lens culinaris*) under Cluster Frontline Demonstrations (CFLD) conducted in Uttar Pradesh. *Indian Journal of Agricultural Sciences*, 88(6), pp.851-859.

- GOI (Government of India). 2022. *Agricultural Statistics at a Glance*. Agricultural Statistics Division, Department of Agriculture and Cooperation and Farmers Welfare, Ministry of Agriculture, GOI, New Delhi.
- Goldbach, H.E., Yu, Q., Wingender, R., Schulz, M., Wimmer, M., Findekle, P. and Baluška, F., 2001. Rapid response reactions of roots to boron deprivation. *Journal of Plant Nutrition and Soil Science*, 164(2), pp.173-181.
- Gupta, U. and Solanki, H., 2013. Impact of boron deficiency on plant growth. *International journal of bioassays*, 2(7), pp.1048-1050.
- Handiganoor, M., Patil, S. and Vasudevan, S., 2017. Response of pigeonpea (*Cajanus cajan* L.) to seed polymerization with micronutrients and foliar spray at different growth stages. *British Journal of Environment and Climate Change*, 7(4), pp.205-213.
- Haque, M.A., Jahiruddin, M. and Islam, M.R., 2000. Effect of sulphur and boron on seed yield of mustard (*Brassica napus*). *Bangladesh J. Seed Sci. & Tech*, 4(1&2), pp.7-11.
- Hussain, M., Khan, M.A., Khan, M.B., Farooq, M. and Farooq, S., 2012. Boron application improves growth, yield and net economic return of rice. *Rice Science*, 19(3), pp.259-262.
- Islam, B., 2005. Requirement of boron for mustard, wheat and chickpea based cropping pattern. *Unpublished [Ph. D. Thesis], Department of Soil Science, Bangladesh Agricultural University, Mymensingh*, pp.1-124.
- Jaiswal, A.D., Singh, S.K., Singh, Y.K., Singh, S. and Yadav, S.N., 2015. Effect of sulphur and boron on yield and quality of mustard (*Brassica juncea* L.) grown on Vindhyan red soil. *Journal of the Indian Society of Soil Science*, 63(3), pp.362-364.
- Jankowski, K.J., Załuski, D. and Sokólski, M., 2020. Canola-quality white mustard: Agronomic management and seed yield. *Industrial crops and products*, 145, p.112138.
- Kapila Shekhawat, K.S., Shivay, Y.S. and Dinesh Kumar, D.K., 2008. Productivity and nutrient uptake of spring sunflower (*Helianthus annuus*) is influenced by nitrogen sources, sulphur and boron levels. *Indian Journal of Agricultural Science* 78, 90-94.
- Kumararaja P, Premi O P and Kandpal B K. 2015. Application of boron enhances Indian mustard (*Brassica juncea*) productivity and quality under boron deficient calcareous soil in a semi-arid environment S249-S254.
- Li, C., Xiong, Y., Cui, Z., Huang, Q., Xu, X., Han, W. and Huang, G., 2020. Effect of irrigation and fertilization regimes on grain yield, water and nitrogen productivity of mulching cultivated maize (*Zea mays* L.) in the Hetao Irrigation District of China. *Agricultural Water Management*, 232, p.106065.
- Lu, X., Lu, Y. and Mao, D., 2000. The levels of available boron in red earth in central Zhejiang Province and boron nutrition of rape. *Soils and Fertilizers (Beijing)*, (1), pp.30-34.

- Miwa, K. and Fujiwara, T., 2010. Boron transport in plants: co-ordinated regulation of transporters. *Annals of Botany* 105 (7), pp.1103-1108.
- Padbhushan, R. and Kumar, D., 2015. Yield and nutrient up take of Green gram (*Vigna radiate* L.) as influenced by boron application in boron deficient calcareous soils of Punjab. *Communications in Soil Science and Plant Analysis* 46, 908-923.
- Pramanick, B., Mahapatra, B.S., Datta, D., Dey, P., Singh, S.P., Kumar, A., Paramanik, B. and Awasthi, N., 2023. An innovative approach to improve oil production and quality of mustard (*Brassica juncea* L.) with multi-nutrient-rich polyhalite. *Heliyon* 9(3).
- Premi, O.P., Kandpal, B.K., Rathore, S.S., Shekhawat, K. and Chauhan, J.S., 2013. Green manuring, mustard residue recycling and fertilizer application affects productivity and sustainability of Indian mustard (*Brassica juncea* L.) in Indian semi-arid tropics. *Industrial Crops and Products* 41, pp.423-429
- Randhawa, M.K., Dhaliwal, S.S., Sharma, V., Toor, A.S., Sharma, S., Kaur, M. and Verma, G., 2021. Nutrient use efficiency as a strong indicator of nutritional security and builders of soil nutrient status through integrated nutrient management technology in a rice-wheat system in northwestern India. *Sustainability* 13(8), p.4551.
- Rashid, A. and Ryan, J., 2004. Micronutrient constraints to crop production in soils with Mediterranean-type characteristics: a review. *Journal of Plant Nutrition* 27(6), pp.959-975.
- Rehman, A., Wang, X., Hussain, A., Qamar, R., Khalofah, A. and Hussain, M., 2022. Boron Application in Yermosols Improves Grain Yield and Quality of Chickpea (*Cicer arietinum* L.). *Journal of King Saud University-Science* 34(2), p.101768
- Safdar, M.E., Qamar, R., Javed, A., Nadeem, M.A., Javeed, H.M.R., Farooq, S., Głowacka, A., Michałek, S., Alwahibi, M.S., Elshikh, M.S. and Ahmed, M.A., 2023. Combined application of boron and zinc improves seed and oil yields and oil quality of oilseed rape (*Brassica napus* L.). *Agronomy* 13(8), p.2020.
- Saha, P.K., Saleque, M.A., Zaman, S.K. and Bhuiyan, N.J., 2003. Response of mustard to S, Zn and B in calcareous soil. *Bangladesh Journal of Crop Sciences* 28, 633-636.
- Shah, K.A., Gurjar, R., Parmar, H.C. and Sonani, V.V., 2016. Effect of sulphur and zinc fertilization on yield and quality of pigeonpea in sandy loam soil. *Green Farming* 7 (2), pp.495-497.
- Shekhawat, K., Rathore, S.S., Kandpal, B.K., Premi, O.P., Singh, D. and Chauhan, B.S., 2016. Crop establishment techniques affect productivity, sustainability, and soil health under mustard-based cropping systems of Indian semi-arid regions. *Soil and Tillage Research* 158, pp.137-146.
- Singh, S.P., Mahapatra, B.S., Pramanick, B. and Yadav, V.R., 2021. Effect of irrigation levels, planting methods and mulching on nutrient uptake, yield,

- quality, water and fertilizer productivity of field mustard (*Brassica rapa* L.) under sandy loam soil. *Agricultural Water Management* **244**, p.106539.
- Srinivasan, G., Gobi, R., Balasubramanian, A. and Sathiyamurthi, S., 2019. Influence of nipping and nutrient management practices on growth, yield attributes and yield in pigeon pea. *Plant Archives* **19**, 737–740.
- Timotiwu, P.B., Agustiansyah, A., Pramono, E. and Agustin, S., 2018. Phosphorus and boron application on growth, yield, and quality of soybean seeds (*Glycine max* [L.] Merril). *Journal of Agricultural Science* **10**(6), pp.372-382.
- Verma, A., Shahi, U.P., Chaurasia, P. and Kumawat, L., 2020. Effect of mode of micronutrients application on growth and yield of pigeon pea (*Cajanus cajan* L.) in sandy loam soil. *International Journal of current Microbiology and applied Sciences* **9**, 17–24.
- Wimmer, M.A., Abreu, I., Bell, R.W., Bienert, M.D., Brown, P.H., Dell, B., Fujiwara, T., Goldbach, H.E., Lehto, T., Mock, H.P. and Wirén, N., 2019. Boron: an essential element for vascular plants. *New Phytologist* **226**(5), pp.1232-1237
- Zhao, Z., Wang, Y., Shi, J., Wang, S., White, P.J., Shi, L. and Xu, F., 2021. Effect of balanced application of boron and phosphorus fertilizers on soil bacterial community, seed yield and phosphorus use efficiency of *Brassica napus*. *Science of The Total Environment* **751**, p.141644.

Table:1 Effect of foliar application of boron on number of primary branches, number of siliquae /plant, number of seeds / siliqua, test weight (g) of the mustard crop during 2018 -2020

Treatment	Number of primary branches /plant			Number of siliquae / plant			Number of seeds/ siliqua			Test weight(g)		
	2018	2019	2020	2018	2019	2020	2018	2019	2020	2018	2019	2020
T1- Farmer's Practice (100%NPK+ without boron)	5.00±0.8	5.50±1.1	5.5±1.1	402.5±8.2	398.0±8.6	400.8±12.4	11.75±0.47	10.95±0.72	12.01±0.88	3.2±0.23	3.37±0.30	3.25±0.35
T2- (100% NPKS + foliar application of Boron @ 0.5 % Borax)	7.75±0.5	7.00±0.7	8.08±0.8	468.8±26.9	427.2±26.6	451.3±36.4	13.53±0.76	13.03±0.65	14.01±0.48	3.6±0.28	3.70±0.63	3.85±0.33
P(T<=t)	0.001	0.020	0.016	0.002	0.022	0.024	0.001	0.005	0.001	0.005	0.027	0.028

Table: 2 Effect of foliar application of boron on dry matter production (g/plant) seed yield (q/ha), assimilate supply and production efficiency of the mustard crop during 2018 -2020

Treatment	Dry matter production/plant (g)			Seed yield (q/ha)			Assimilate Supply (g siliqua ⁻¹)			Production efficiency (kg ha ⁻¹ day ⁻¹)		
	2018	2019	2020	2018	2019	2020	2018	2019	2020	2018	2019	2020
T1- Farmer's Practice (100%NPK+ without boron)	38.4±1.8	36.4±4.6	38.9±1.7	22.5±0.9b	19.0±2.0b	20.0±1.7b	10.51±0.79	11.17±0.61	10.33±1.11	16.67±0.70	14.07±1.49	14.81±1.30
T2- (100% NPKS + foliar application of Boron @ 0.5 % Borax)	39.9±1.2	38.1±1.1	41.2±1.4	24.7±0.6a	22.5±0.9a	24.0±0.7a	11.76±1.8	11.22±0.66	11±0.60	18.31±0.51	16.67±0.70	17.78±0.64

P(T<=t)	0.108	0.221	0.044	0.007	0.021	0.001	0.026	0.480	0.203	0.001	0.041	0.001
---------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------

Table: 3 Effect of foliar application of boron on partial factor of productivity of N,P, K and S of the mustard crop during 2018 -2020

Treatment	N Partial factor productivity (kg kg ⁻¹)			P Partial factor productivity (kg kg ⁻¹)			K Partial factor productivity (kg kg ⁻¹)			S Partial factor productivity (kg kg ⁻¹)		
	2018	2019	2020	2018	2019	2020	2018	2019	2020	2018	2019	2020
T1- Farmer's Practice (100%NPK+ without boron)	28.13±1.19	23.75±2.5	25±2.19	37.50±1.5	31.67±3.3	33.3±2.92	56.25±2.39	47.50±5.5	50±4.38	75.00±3.19	63.33±6.73	66.67±5.85
T2- (100% NPKS + foliar application of Boron @ 0.5 % Borax)	30.90±0.87	28.13±1.1	30±1.08	41.19±1.1	37.50±1.4	40.0±1.44	61.79±1.74	56.25±2.40	60±2.16	82.39±2.32	75.00±2.28	80±2.88
P(T<=t)	0.007	0.021	0.001	0.027	0.001	0.005	0.007	0.021	0.001	0.002	0.011	0.005

Table:4 Relationship between yield and yield attributes of mustard as influenced by boron spary (mean data of three years)

X-axis	Y-axis	Correlation coefficient R ²	Regression equation
Grain yield	sehcnaB yramirP	0.57	y = 0.447x - 3.4144
Grain yield	Number of siliquae/ plant	0.77	y = 11.739x + 165.11
Grain yield	Number of seeds/ siliqua	0.75	y = 0.4563x + 2.4576
Grain yield	Test weight (1000 seed weight)	0.39	y = 0.0736x + 1.8658
Grain yield	noitlaumcca rettamyrd	0.60	y = 0.5695x + 26.223
Grain yield	Assimilate supply	0.22	y = 0.1108x + 8.5482




Figure: 1 Effect of foliar application of boron on cost of cultivation, gross return and net return of the mustard crop during 2018 - 2020




Figure: 2 Effect of foliar application of boron on benefit- cost ratio of the mustard crop during 2018 -2020



Figure: 3,4,5 Rainfall distributions (mm) during crop growing *rabi* season 2018 -21



Figure: 6 Site of Krishi Vigyan Kendra in NCT Delhi and its working area