

New host record of invasive thrips, *Thrips parvispinus* (Karny) (Thysanoptera: Thripidae) on mungbean in Gujarat, India

ABSTRACT

The present investigation was carried out for the incidence of thrips on mungbeans in the Northern region of the Indian state of Gujarat has revealed that four thrips species viz., *Thrips parvispinus* (Karny), *Megalurothrips usitatus* (Bagnall), *Megalurothrips* sp., and *Thrips palmi* Karny, were discovered and identified from mungbean flowers in the region during the investigation periods 2021–22 and 2022–23. For the first time in Gujarat and India, *Thrips parvispinus* (Karny), a novel invasive thrips was found infesting pulse crop-mungbean or green gram

Keywords: Invasive thrips, *Thrips parvispinus* and *Vigna radiata*

INTRODUCTION

Pulses hold a significant role in Indian agriculture, being both the largest producer and consumer of these crops globally. They offer a high-quality protein, nearly three times that of cereals (Upadhyay *et al.*, 1999). This makes them a cost-effective protein source, crucial in addressing malnutrition. Mungbean or green gram comprising about 24 per cent protein, contain roughly two thirds of soyabean protein, twice of wheat and thrice the protein found in rice. The protein is abundant in lysine, an essential amino acid which is lacked in cereal grains (Haytowitz and Matthews, 1986).

Thysanoptera is an order of insects that has two suborders: Terebrantia and Tubulifera. Due to the existence of plant viral vectors and agriculturally significant pests, the suborder Terebrantia is the most commercially significant suborder within Thysanoptera (Rachana *et al.* 2020). Thrips are minute insects belonging to the order Thysanoptera. Their pre-pupal stage, which occurs between the larval and pupal stages, fringed wings and a protruding bladder-like structure at the tarsal end are their characteristics. The asymmetrical mouthparts with the remaining right mandible were yet another distinctive feature. This order of insects has a

remarkably diverse range of eating preferences; they can be herbivorous, fungivorous, pollinivorous or predatory (Mound, 2005^a).

The discovery that an invasive insect is present on a host plant that is crucial to agriculture causes serious concern and panic for quarantine officials. *T. parvispinus* gradually spreading across states, as indicated by previous reports of this species on various economically significant host plants in Karnataka and the most recent report on an agriculturally significant host in Tamil Nadu. One of the notorious pest species from South East Asia, *T. parvispinus*, causes a significant threat to a wide range of agriculture and horticulture crops. *T. parvispinus*, a thrips of quarantine relevance belonging to the "Thrips orientalis group" (Mound, 2005^b), has been described as a significant pest on numerous unrelated plant families from Thailand to Australia (Mound and Collins, 2000).

MATERIALS AND METHODS

The investigation on inventory of thrips on pulses in the Northern part of the Indian state of Gujarat includes the districts of Aravali, Banaskantha, Gandhinagar, Mehsana, Patan and Sabarkantha during 2021-22 and 2022-23. The survey was conducted at random and consisted of visiting regions of North Gujarat wherever pulses are cultivated. The thrips were collected from the flowers and tender leaves of mungbean by tapering on white paper. At least 40-50 thrips were collected and preserved in 70 percent ethyl alcohol for identification. When the thrips were being collected from their host, the host name, the location of the collection and the date were registered. For additional identification, the specimens were sent to the National Bureau of Agricultural Insect Resources (NBAIR), Bengaluru, an affiliate of the Indian Council of Agricultural Research (ICAR).

RESULTS AND DISCUSSION

Mungbean or green gram is a significant pulse crop in India. Both the seedling and blooming phases of mungbean are infested with thrips. As the seedlings emerge from the earth during the seedling stage, thrips infest the growth point, inhibiting the seedlings from growing if the infestation is severe. Flowering thrips, which feed on the pedicles and stigma of flowers, inflict severe damage and active during the flowering and pod development stages. Severe infestation results in blossom drops and no pod production (Fig. 1 A and B).

Although history of *T. parvispinus* as a pest has never been reported earlier. The purpose of this paper includes outlining *T. parvispinus* as a mungbean pest in India for first time and to

highlight the key traits that are able to be used for recognizing the species when it is collected from that host.

The identification result obtained from ICAR-NBAIR indicated that the discovery of a new invasive thrips, *Thrips parvispinus* (Karny), infesting mungbean reported from Sardarkrushinagar- Dantiwada, district Banaskantha, Gujarat (Table 1). The other species viz., *Megalurothrips usitatus* (Bagnall), *Megalurothrips* sp., and *Thrips palmi* Karny were also identified indicating the complex thrips population in mungbean from different districts of North Gujarat (Aravali, Banaskantha, Gandhinagar, Mehsana, Patan and Sabarkantha). All the thrips species is also identified at molecular level also.

Despite the fact that *T. parvispinus* already recognised, the species was first identified in India on papaya from Bangalore by Tyagi *et al.* (2015). Rachana *et al.* (2018) then reported the same on *Dahlia rosea* blooms in Puttur, Karnataka. The same was later found by Patel *et al.* (2022) on the chilli flower in Umreth, Gujarat. In a survey (Anonymous (2023) conducted in different states specifically targeting mungbean and uradbean thrips. Many thrips species were morphologically identified and reported like *Megalurothrips usitatus* (Bagnall), *Scirtothrips dorsalis* Hood, *Thrips palmi* Karny and *Thrips* sp. From Panjab; *Megalurothrips* sp., *T. palmi* and *Thrips* sp. from Tamil Nadu; *M. distalis*, *Thrips* sp. and *T. palmi* were observed in Uttarakhand; *M. usitatus*, *M. distalis*, *Scirtothrips* sp., *Thrips* sp. and *S. dorsalis* were discovered from West Bengal; *M. usitatus*, *Megalurothrips* sp. and *T. palmi* were recorded from Sardarkrushinagar, Dantiwada in Gujarat; *Thrips* sp., *T. palmi*, *S. dorsalis* and *Scirtothrips* sp. were recorded in Karnataka and *M. typicus* (Bagnall), *M. usitatus* (Bagnall) and *T. palmi* Karny were found in Andhra Pradesh.

Conclusion

The paper deal with identified invasive, *Thrips parvispinus* (Thripidae) recorded from green gram or mungbean from Sardarkrushinagar, Dantiwada, which is new record in Gujarat and India especially in pulse crops.

DECLARATION

We, the undersigned authors of new host record of invasive thrips, *Thrips parvispinus* (Karny) (Thysanoptera: Thripidae) on mungbean in Gujarat, declare that we have fully participated in the research work and writing of this manuscript. We confirm that the content presented in this paper

is original and has not been published elsewhere. Furthermore, we affirm that there are no conflicts of interest, financial or otherwise, that could influence the objectivity or integrity of our work.

REFERENCES

- Anonymous, 2023. All India Coordinated Research project on *Kharif* Pulses. Annual Report on Mungbean, Urdbean, Cowpea, Cluster bean for Spring/Summer/Rice fallow and Rajmash for *Kharif* cultivation. ICAR-Indian Institute of Pulses Research, Kanpur. pp. 255-258.
- Haytowitz, O. B. and Matthews, R. H. 1986. Composition of foods: legumes and legume products. United States Department of Agriculture. *Agricultural Hand Book*. pp. 8-16.
- Mound, L. A. 2005^a. Thysanoptera: diversity and interactions. *Annu. Rev. Entomol.* **50**: 247-269.
- Mound, L. A. 2005^b. The Thrips *orientalis* group from South-east Asia and Australia: Some species identities and relationships (Thysanoptera; Thripidae). *Aust. J. Entomol.* **44**(4): 420-424.
- Mound, L. A. and Collins, D. W. 2000. A South Asian pest species newly recorded from Europe: *Thrips parvispinus* (Thysanoptera: Thripidae), its confused identity and potential quarantine significance. *Eur. J. Entomol.* **97**: 197-200.
- Murai, T., Watanabe, H., Toriumi, W., Adati, T., Okajima, S. 2009. Damage to vegetable crops by *Thrips parvispinus* Karny (Thysanoptera: Thripidae) and preliminary studies on biology and control. *J. Insect Sci.* **10**: 166.
- Patel, N. B., Bhagora, J. K., Raghunandan, B. L. and Patel, N. M. 2022. First Report of New Invasive Thrips, *Thrips parvispinus* (Karny) (Thripidae: Thysanoptera) in Chilli Fields of Umreth in Anand District of Gujarat State. *Int. J. Environ.* **12**(3): 73-78.

- Rachana, R. R., Rayer, S. G., Giraddi, R. S., Kalappanavar, I. K. and Alagundagi, S. C. 2020. New records of Terebrantia thrips from Karnataka, India. *J. Entomol. Zool. Stud.* **8**(4): 2189-2192.
- Rachana, R.R., Roselin, P. and Varatharajan, R. 2018. Report of invasive thrips species, *Thrips parvispinus* (Thysanoptera: Thripidae) on *Dahlia rosea* (Asteraceae) in Karnataka. *Pest Manage. Hortic. Ecsyst.***24**(2): 175-176.
- Tyagi, K., Kumar, V., Singha, D., Chakraborty, R. 2015. Morphological and DNA barcoding evidence for invasive pest thrips, *Thrips parvispinus* (Thysanoptera: Thripidae), newly recorded from India. *J. Insect Sci.***15**(1): 105.
- Upadhyay, R. G., Sharma, S. and Drawal, N. S. 1999. Effect of Rhizobium inoculation and graded levels of P on the growth and yield of greengram. *Legum. Res.***22**: 277-279.



Figure 1A: General view of mungbean field




Figure 1B: Mungbean flowerinfected by *Thrips parvispinus*

Table 1.: Thrips species reported from North Gujarat in Mungbean

Sr. No.	Location	Scientific name	classification
1.	Location: Latitude: 24.3219 Longitude: 72.3008	<i>Megalurothrips usitatus</i> (Bagnall) <i>Thrips parvispinus</i> (Karny)	Thysanoptera: Terebrantia: Thripidae
2.	Location: Latitude: 24.4611 Longitude: 71.6729	<i>Megalurothrips usitatus</i> (Bagnall)	Thysanoptera: Terebrantia: Thripidae
3.	Location: Latitude: 23.8341 Longitude: 71.5860	<i>Megalurothrips usitatus</i> (Bagnall)	Thysanoptera: Terebrantia: Thripidae
4.	Location: Latitude: 23.6397 Longitude: 72.7043	<i>Megalurothrips usitatus</i> (Bagnall)	Thysanoptera: Terebrantia: Thripidae
5.	Location: Latitude: 23.3755 Longitude: 72.6877	<i>Megalurothrips usitatus</i> (Bagnall)	Thysanoptera: Terebrantia: Thripidae
6.	Location: Latitude: 23.3442 Longitude: 72.6854	<i>Megalurothrips usitatus</i> (Bagnall)	Thysanoptera: Terebrantia: Thripidae
7.	Location: Latitude: 24.3176 Longitude: 72.2752	<i>Megalurothrips usitatus</i> (Bagnall) <i>Megalurothrips</i> sp. <i>Thrips palmi</i> Karny	Thysanoptera: Terebrantia: Thripidae
8.	Location: Latitude: 24.3215 Longitude: 72.3035	<i>Megalurothrips usitatus</i> (Bagnall)	Thysanoptera: Terebrantia: Thripidae
9.	Location: Latitude: 23.1244 Longitude: 73.2099	<i>Megalurothrips usitatus</i> (Bagnall)	Thysanoptera: Terebrantia: Thripidae
10.	Location: Latitude: 24.2292 Longitude: 73.0151	<i>Megalurothrips usitatus</i> (Bagnall)	Thysanoptera: Terebrantia: Thripidae
11.	Location: Latitude: 24.1247 Longitude: 73.0128	<i>Megalurothrips usitatus</i> (Bagnall)	Thysanoptera: Terebrantia: Thripidae