

Effect of Soil Application of cyazypyr 20% SC, New Anthranilic Diamide Insecticide against whitefly, *Bemisia tabaci* Genn. in brinjal

ABSTRACT

Brinjal *Solanum melongena* L. is an herbaceous, tropical perennial plant, belongs to the family Solanaceae which is grown for its edible fruit. Among the different major insect pests infesting brinjal, whitefly, *Bemisia tabaci* (Genn.), is very important under West Bengal condition. The experiment was conducted during the 2013 and 2014 in the University farm at Kalyani, West Bengal state of India. Cv 'Muktakeshi' was grown in plots measuring 5 m×5 m, at spacing of 1m x 0.75m with three replication. The plots were set out in a randomized block design with six treatments including an untreated check. Five doses of cyazypyr 20% SC (4.5 MAT ,6.0 MAT, 7.5MAT, 9.0 MAT and, 12 MAT in both year 2013 and 2014) were sprayed every year for their efficacy, After 50 days of treatment cyazypyr 20% SC @ 9.0 and 7.5 MAT (3.33 and 6.87 whiteflies / 5 leaves, respectively) maintained their superiority in controlling whiteflies, while @ 6.0 and 4.5 MAT (7.93 and 8.40 whiteflies / 5 leaves, respectively) and this treatment failed to show any significant difference from untreated control

INTRODUCTION

Brinjal *Solanum melongena* L. is an herbaceous, tropical perennial plant, belongs to the family Solanaceae which is grown for its edible fruit. Brinjal is known as Bazinga in Egypt, aubergine in France and England, eggplant in the United States, and brinjal in India (Zayed *et al.*, 2015). It is also referred to as guinea squash or King of vegetables and India and Indochina are considered the centers of origin (14). Being the top ten vegetables in the world, it is extensively grown in India, Pakistan, China, Philippines, Bangladesh, Egypt, France, Italy, Middle East, Far East, and the U.S.A. It is rich in nutrients like dietary fiber, ascorbic acid, vitamin K, folate, niacin, vitamin B6, pantothenic acid, potassium, iron, magnesium, manganese, phosphorus, and copper (12). It is native of India and second largest brinjal producing country after China with 27.1 % share. It is an important vegetable grown in all the seasons.

Its fruits are wide range in size, display variation in fruit shape ranging from oval or egg-shaped to long club-shaped and color depending on the varieties, fruits may be

black, purple, purple white, white, yellow or purple. It has two main groups: long (called Bride) and oval or spherical (called Romy) (1). Among the insect pests infesting brinjal, the major ones are epilachna beetle, *Epilachna vigintioctopunctata* (Fab.), shoot and fruit borer, *Leucinodes orbonalis* (Guen.), whitefly, *Bemisia tabaci* (Genn.), leafhopper, *Empoasca flavescens* (Distant), and non insect pest, red spider mite, *Tetranychus macfurlanei*. Among the different major insect pests infesting brinjal, whitefly, *Bemisia tabaci* (Genn.), is very important under West Bengal condition. To avoid the crop loss by this insect, the frequent use of toxic chemical insecticides has been a common practice to the brinjal growers. The new generation of pesticide molecules have been claimed to be effective as well as safer for non-target organisms [13, 3, 8, 12]. The use of insecticides could be more effective depending on selection of chemicals, doses, method and time of application. Hence, keeping the above point in view, present investigation was carried to evaluate the bio-efficacy of cyazypyr 20% SC on whitefly, *Bemisia tabaci* (Genn.), under field condition.

MATERIAL AND METHODS

The experiment was conducted during the 2013 and 2014 in the University farm at Kalyani, West Bengal state of India. Cv 'Muktakeshi' was grown in plots measuring 5 m×5 m, at spacing of 1m x 0.75m with three replication during the period from mid-April to July, two year, following recommended package of practices. The plots were set out in a randomized block design with four doses of cyazypyr 20% SC, 4.5, 6.0, 7.5 and 9.0 MAT (Milligram active ingredient per target) were applied in soil at the base of the plant twice; first at time of planting and after 15 days of planting. There were altogether 5 treatments including untreated control with 3 replications. Control plots were treated with equal amount of water only.

Data on whiteflies (adults) was recorded at 10 days interval starting from 30 days after planting up to 60 days after planting from 5 randomly selected plants / plot following the same method as in the previous experiment.

RESULTS AND DISCUSSION

In the season 2013, whitefly population recorded in different treatments, showed significant reduction in plots treated with cyazypyr 20% SC @ 9.0, 7.5, 6.0 and

4.5 MAT (0.80, 1.33, 2.87 and 4.53 whiteflies / 5 leaves, respectively) as compared to untreated control (10.53 whiteflies / 5 leaves). After 40 days of treatment, cyazypyr 20% SC @ 9.0, 7.5, 6.0 and 4.5 MAT (2.87, 3.27, 4.73 and 7.80 whiteflies / 5 leaves, respectively) as compared to untreated control (11.60 whiteflies / 5 leaves). After 50 days of treatment cyazypyr 20% SC @ 9.0 and 7.5 MAT (3.33 and 6.87 whiteflies / 5 leaves, respectively) maintained their superiority in controlling whiteflies, while @ 6.0 and 4.5 MAT (7.93 and 8.40 whiteflies / 5 leaves, respectively) and this treatment failed to show any significant difference from untreated control (8.33 whiteflies / 5 leaves). After 60 days of treatment cyazypyr 20% SC @ 6.0 and 4.5 MAT showed similar results (9.93 whiteflies / 5 leaves). Cyazypyr 20% SC @ 9.0 and 7.5 MAT (6.80 and 7.53 whiteflies / 5 leaves), these treatments decreased pest population as compared to untreated control (9.93 whiteflies / 5 leaves) Vivek *et. al.* (2016) also found that whitefly mortality in different treatments ranged between 62 and 96% for cyantraniliprole, and 95–100% for combination treatments. No phytotoxicity was observed for any treatment. Christian and Nabil (2023) found that Cyantraniliprole showed better results relative to spiromesifen, spirotetramat, tolfenpyrad, sulfoxaflor, and flupyradifurone treated plots. After the fifth application, cyantraniliprole was able to reduce the whiteflies eggs' populations by 23.69% and 42.47% in greenhouses 1 and 2, respectively; whereas whiteflies nymphs' populations were reduced by 76.25% in greenhouse. Chand *et. al.* (2018) The most effective treatment was cyazypyr 10% OD @ 105g a.i./ha followed by cyazypyr 10% OD @ 90g a.i./ha.

In the season 2014, after 30 days of planting all the doses of cyazypyr 20% SC had significantly lower population of whitefly than untreated control. Cyazypyr 20% SC @ 9.0 and 7.5 MAT showed to be superior significantly lowering the whitefly population than the two lower doses of the same insecticide.

After 40 days of treatment, cyazypyr 20% SC @ 9.0 MAT recorded superiority over rest of the treatments in controlling whitefly population. Result achieved with cyazypyr 20% SC @ 6.0 and 7.5 MAT was statistically homogeneous (2.27 – 7.40 and 2.07 – 6.47 whitefly / 5 leaves).

Table 1. Number of whitefly / 5 leave due to different treatment

Treatment	2013				2014			
	30DAS	40DAS	50DAS	60DAS	30DAS	40DAS	50DAS	60DAS
Cyazypyr 20% @ 4.5 MAT	4.53 (2.12)	7.80 (2.79)	8.40 (2.89)	9.93 (3.15)	1.80 (1.34)	4.40 (2.10)	6.07 (2.46)	7.40 (2.71)
Cyazypyr 20% @ 6.0 MAT	2.87 (1.68)	4.73 (2.16)	7.93 (2.81)	9.93 (3.15)	0.93 (0.96)	2.27 (1.49)	3.13 (1.76)	7.40 (2.71)
Cyazypyr 20% @ 7.5 MAT	1.33 (1.14)	3.27 (1.80)	6.87 (2.62)	7.53 (2.74)	0.40 (0.62)	2.07 (1.43)	2.93 (1.71)	6.47 (2.54)
Cyazypyr 20% @ 9.0 MAT	0.80 (0.89)	2.87 (1.69)	5.33 (2.31)	6.80 (2.61)	0.27 (0.51)	1.00 (1.00)	2.07 (1.44)	4.93 (2.21)
Untreated Control	10.53 (3.25)	11.60 (3.40)	8.33 (2.88)	9.93 (3.15)	16.80 (4.09)	10.47 (3.23)	13.20 (3.63)	7.40 (2.70)
CD	0.36	0.32	0.38	0.32	0.30	0.39	0.29	NS

*Values within parentheses are square root transformed**MAT- milligram ai per target

REFERENCES

- 1 Abd-Elgawad, M. M. (2021). Biological control of nematodes infecting eggplant in Egypt. *Bulletin of the National Research Centre*, 45(1), 1-9.
- 2 Biswas, R.K. & Chaterjee. M. (2008). Effectiveness of some systemic insecticides against the whitefly, *Bemisia tabaci* (Gennadius), on brinjal and the jassid, *Amrasca biguttula biguttula* Ishida, on okra. *Pest Management and Economic Zoology* **16** (1): 37-42
- 3 Borad, P.K., Patel, H.M., Chavda, N. & Patel, J.R. (2002). Bio-efficacy of endosulfan and cypermethrin mixture against insect pests of brinjal (*Solanum melongena*). *Indian Journal of Agricultural Sciences* **72**:685-88.
- 4 Chand P., Vishwakarma R., Mandal S.K. and Chandra U (2018), Bio-Efficacy of Cyazypyr 10% OD, New Anthranilic Diamide Insecticide, against whitefly *Bemisia tabaci* and Jassid *Empoasca flavescens* on brinjal *Bull. Env. Pharmacol. Life Sci.*, Vol 7 [10] September 124-131
- 5 Hall, T. (2007). Ecological effects assessment of flubendiamide. *Pflanz.-Nach.*, Bayer 60 (2): 167–182.

- 6 Kareh C and Nemer N (2023) Evaluation of Insecticides in the Management of Whiteflies (*Bemisia tabaci* Gennadius) and their Impacts on Yield of Eggplants, *Universal Journal of Agricultural Research* 11(4): 715-722
- 7 Kumar, V, Kakkar G., Cindy L. McKenzie and Lance S. Osborne (2016) Effect of Soil Application of Cyantraniliprole on *Bemisia tabaci* (MED Whitefly) and mlyseius
- Manda, I S., Singh, N.J. & Konar A. (2010). Efficacy of synthetic and botanical insecticide against whitefly (*Bemisia tabaci*) and shoot and fruit borer (*Leucinodes orbonalis*) on brinjal (*Solanum melongena* L.).
- 8 Mandal, S. K. (2012). Bio-Efficacy of Cyazypyr 10% OD, a New Anthranilic Diamide Insecticide, against the Insect Pests of Tomato and Its Impact on Natural Enemies and Crop Health *Acta Phytopathologica et Entomologica Hungarica* 47 (2), pp. 233–249
- 9 Misra, H.P. (2008). New promising insecticides for the management of brinjal shoot and fruit borer, *Leucinodes orbonalis* Guenee. *Pest Manage. Hort. Ecosys.* 14 (2): 140–147.
- 10 Muthukumar, M. & Kalyanasundaram, M. (2003). efficacy of certain insecticides against major sucking insects of brinjal (*Solanum melongena* L.). *South Indian brinjal (Solanum melongena L.)*. *South Indian* 51: 207-13
- 11 Patel JJ Patel BH Bhatt HV Maghodia AB Bhalala MK. (2006). Bioefficacy of diafenthiuron 50 WP against sucking pests of brinjal (*Solanum melongena* L.). *Indian Journal of Entomology* 68 (3): 272-73.
- 12 Sathe, T. V., Patil, S. S., Bhosale, A. M., Devkar, S. S., Govali, C. S. and Hankare, S. S. (2016). Ecology and Control of Brinjal insect pests from Kolhapur region, India. *Bio-life*, 4(1):147- 154.
- 13 Sontakke, B. K., Das N. & Swain, L.K. (2007). Bioefficacy of emamectin benzoate against boll worm complex in cotton. *Ann. Pl. Protec. Sci.* 15 (1): 371–376.
- Tonishi, M.H., Nakao T., Furuya A., Seo H., Kodama K., Tsubata S., Fujioka H., Kodama T., Hirooka T. & Nishimatsu T. (2005). Flubendiamide, a novel insecticide highly active against lepidopterous insect pests. *J. Pestic. Sci.* 30: 354–360.
- 14 Vavilov, N. I. (1951). *The origin, variation, immunity, and breeding of cultivated plants*. Vol.72, no. 6. 91-99.
- 15 Zayed, G. A., Abdo, A. A., Hammam, H. B. and Khafagi, E. Y. (2017). *Cultivation and production of pepper and eggplant in Egypt*. Technical issue No. 15, General Administration of Agricultural Culture, Ministry of Agriculture, Egypt.