

# Review on Effect of Different Preservatives on Vase Life of Gladiolus

---

**Abstract:** Globally Gladiolus is most demanding flower and day by day requirement is increasing this study is examined the different methods and ways to increase the vase life of gladiolus, mainly focuses on the impact of various preservatives on the vase life of gladiolus flowers. Different preservative solutions were applied to the cut stems of gladiolus blooms, and their effects on flower freshness, color retention, water uptake, and overall longevity were evaluated. It contains the different preservatives Sucrose solution, CoSO<sub>4</sub>, Calcium, Salicylic acid, different concentration of GA<sub>3</sub>, CCC, NAA. GA<sub>3</sub> proves the best PGR for pre harvest treatment to maintain quality of flowers. Sucrose solution of 4% along with CoSO<sub>4</sub> is good extent the vase life of Flower. Results indicate that certain preservatives significantly extend the vase life of gladiolus flowers by providing essential nutrients, regulating water balance, and inhibiting microbial growth. These findings offer valuable insights for optimizing post-harvest management practices to enhance the quality and longevity of gladiolus blooms in floral arrangements.

**Keywords ;** Gladiolus, Vase-Life, PGR, Preservatives, Flower quality, Microbial growth, Post harvest management.

---

## Introduction

Gladiolus (*Gladiolus grandiflorus* L.) is one of the most important ornamental and bulbous flowering plant, it carries noble history with it. The name of Gladiolus derived from latin “Gladius (Sword)”, hence it known as sword lily. It originates from South Africa and Asia belongs to monocot family Iridaceae (Prabhesh *et al.*, 2022). As well as Gladiolus knew as “Queen of bulbous plant” is well known in every corner of the globe which ranks 8th position in cut flowers, 1st position in bulbous plants and has more than 150 species in the world trade whose leading country is (Iamichhane *et*

*al.*, 2022). The number of species found in the genus is 255 (Shaukat *et al.*, 2013). The main goal is to grow beautiful cut flowers that can be sold in local flower markets or used for decoration in gardens. However, it's worth mentioning that the dried bulbs of these flowers have medicinal benefits when ground into a powder. The gladiolus is a perennial bulbous and half-hardy flowering plant containing attractive, elegant, and delicate florets opening in sequence from bottom to top along with the spikes. They vary in their height from one to four feet; some are soft and light, others robust and strong adapted to any climate except the frigid condition (Childs, 1873). Vase life of this flower is noticeably short due to blockage of the vascular system by air or bacterial growth, reducing water uptake, and blocking xylem vessels. Short vase life of cut flower has been a major problem in floriculture industry as the longevity of vase life is a crucial factor in consumer preference (Marandi *et al.*, 2011). Different experiments and studies had proved that vase life of gladiolus can be extended by adding or subjecting to different chemical solutions at different concentration, such as; chlorine hydroxyquinoline, sulphates, salicylic acid, humic acid, silver nanoparticles, sugar solutions, diethyl sulphonate, and different plant growth regulators ( Lamichhane *et al.*, 2022). The American Conference of Governmental Industrial Hygienists has recommended exposure limit of soluble silver compounds by 0.01mg/m<sup>3</sup>. While ionic silver compound has a toxic effect on human as well as environment and, organic acid and essential oil had a noticeable effect on the improvement of vase life of cut gladiolus flowers which might be due to their antibacterial properties that reduce bacterial proliferation in the stem vessels of cut flowers (Jowkar *et al.*, 2012; Marandi *et al.*, 2011). Therefore, this study aims to evaluate various vase solutions and their effectiveness in terms of cost and the quality of the produce.

---

### **Vase Life of Gladiolus**

Vase life of flower is influenced by respiration, carbohydrates, deterioration, disease and pests, and water absorption. Ethylene synthesis plays crucial role in vase life.

Various factors, including water stress, carbohydrate depletion, microbial activity, and ethylene exposure, can impact floral senescence. To prolong the vase life of cut flowers by mitigating these effects, preservative solutions are employed post-harvest. These solutions enhance assimilate distribution, regulate water uptake, shield floral

stems from microbial damage, and minimize the influence of ethylene, thereby delaying the onset of floral senescence (Pulido *et al.*, 2021). Carbohydrates and soluble sugar helps to maintain quality of petals for longer time. Flowers are treated with sufficient amounts of sugars like sucrose and an appropriate preservative like silver thiosulfate can greatly enhance the longevity of cut flowers (Hussan *et al.*, 2016). As below discussed not only sucrose but along with sucrose there are other agents are also effective to extent vase life of Cut flowers.

## **Pre and Post harvest affecting factors.**

### **Pre harvesting factors**

#### **Constitution of Inheritant characters**

The genetic composition of the cultivars as well as differences in cell wall thickening, peroxidase and lignification levels, ethylene forming enzyme (EFE) and aminocyclopropane carboxylic acid (ACC) production have all been linked to these variations (Khanal 2014). Khanal (2014) mentioned in term paper have assessed six gladiolus cultivars in Hissar, India, and discovered variance in the responses of the varieties for various attributes. Miniature had the shortest vase life among the tested cultivars, while Pusa Suhagain had the longest. Similarly, five gladiolus cultivars—Agnerekha, American Beauty, Friendship, Mansoer red, and true Yellow—over the course of two seasons in Kerala, India. They mentioned that of the tested varieties, Agnerekha and Mansoer were the best for a longer vase life, while American Beauty and Friendship were found to be superior for cut flower yield (Khanal 2014). Variations in stem lignification, flower diameter, fresh weight, water uptake, and other factors are caused by genetic diversity and ultimately result in differences in the vase life of various flowers or different types of the same flowers. Even genetic variations exist amongst many cultivars of the same crop, as following different varieties with different days of vase life, IIHR-G-12 (14.20), Urmil (8.97), American Beauty (9.67), Moon Magic (5.39) (Verma and Singh, 2021). Differences in the genetic structure of flower species and varieties result in significant differences in their post-harvest lifespan (Gupta and Dubey 2017). The experiment was conducted on the different varieties of gladiolus it showed the best performance for maximum number of corms per plot was “Legend” (37.67) and “Trader Horn” was maximum number of corms per plant (70.83). The maximum vase life was observed in

“Darshan” (17.56 days) and maximum number of florets remaining open at a time in vase was observed in “Hunting Song” (6.78 florets) (Singh *et al.*, 2017).

## **Environmental Factors**

### **Temperature**

By an increase in respiration rate, high temperatures can reduce floral quality and carbohydrate supplies. According to some observations, the longevity of cut flowers was reduced by temperatures as low as 12°–15°C or as high as 27°C during the three weeks before to harvest. The knowledge of temperature impacts on vase life is complicated by the potential interplay of temperature with other factors. Anthocyanins and flavonoids are two more pigments that can become more pigmented at low temperatures (Vehniwal and Abbey, 2019). The reduced amount of carbohydrates in petals as a result of the cold temperature and weak light will cause the vase life to be shortened. Sugar level has been found to be closely associated with the postharvest life of cut flowers because temperature and light intensity directly affect the photosynthate level in flower tissues (Khanal, 2014).

### **Humidity**

High relative humidity levels (over 80% to 90%) shorten the postharvest life of cut flowers because they negatively impact the amount of calcium content (Mortensen and Fjeld, 1998) . High relative humidity (RH) of 90–95% is advised and is crucial for long-term storage because calcium level in the tissue is thought to affect the formation and function of the cell wall and membrane (Hardenburg *et al.*, 1986). RH of 80 or above is generally utilised in practice for short term storage (1-2 days) because it is safe enough to delay withering due to moisture loss. Verma and Singh mentioned Humidity useful in preventing water loss from the plant's surface and preserving the temperature of the plant. While high humidity promotes photosynthesis, it can also lead to fungus-related illnesses. Reduced humidity can lead to greater water loss, senescence, thin leaves, browning of the leaf margins, and chilling injury. Water stress can therefore clearly shorten a flower's vase life (Bhattacharjee and De,

2005), and flowers produced at varying humidity levels will exhibit variations in their vase life (Verma and Singh, 2021).

## **Post harvesting factors**

### **Ethylene**

Senescence is the time when flower petals are converted to ethylene. The two primary enzymes in charge of producing ethylene are ACC oxidase and ACC synthase (Verma and Singh, 2021). Certain essential growth stages, such as the wilt of flowers and the abscission of leaves, together with other external conditions like mechanical injury and environmental stress, drive the production of ethylene (Vehniwal and Abbey, 2019). Carbon makes up the majority of plant dry matter (45%). Through their stomata, plants absorb it from the surrounding air and fix it through the process of photosynthesis (Verma and Singh, 2021).

### **Carbon dioxide**

When cultivated at low and high CO<sub>2</sub> levels during the vegetative phase, respectively, plants grown at elevated CO<sub>2</sub> during the generative growth phase produced 15 and 10% more plants with multiple spikes and a better floral stem quality. The quantity of spikes was not significantly affected by elevated CO<sub>2</sub> throughout the vegetative growth phase, but the pattern suggests that spike quality was on the rise (Kromwijk *et al.*, 2014)

### **Pest and diseases**

Fusarium wilt, Yellow or corm rot, Storage rots, Dry or neck rot, Botrytis blight and flower rot, Virus disease these are the common pest and diseases of gladiolus which can be directly indirectly in low or high percentage affected on the vase life of gladiolus. This can be managed by following various management practices.

Method for harvesting

### **Importance of Sucrose**

Adding sugars in a vase solution has been found to be beneficial for prolonging the freshness and lifespan of cut flowers, it associated with the improvement water balance, improvement of color in case of carnation and rose, Helps to stimulate flower opening (Ichimura 1998). Sucrose, administered via pulse therapy, suppressed the genes involved in ethylene biosynthesis, leading to a decrease in ethylene production. However, in contrast to the control group, sucrose treatment notably increased xylem blockage on the cut stem surfaces while decreasing antioxidant activity, relative fresh weight, and the expression of the cysteine proteinase inhibitor gene (DcCPi) (Park *et al.*, 2017). Sugar plays a crucial role in extending the lifespan of flowers, particularly when they are cut, as they no longer receive essential nutrients and hormones from the parent plant (Muraleedharan and Joshi 2017).

### **Effect of Sucrose**

Han 1992, conducted research on effect of sucrose solution has showed that the presence of sucrose in the vase solution improves the quality of cut spikes by prolonging their display life and enhancing the length of inflorescence with visible color. When a consistent supply of sucrose is provided in the vase solution, it leads to the complete opening of flower heads, achieving a 100% opening rate, 200ppm of citric acid with 4% of sucrose is good to extent the vase life of gladiolus (Koirala *et al.*, 2022). Sucrose 4% combine with CoSO<sub>4</sub> 100ppm proved long duration of vase life with 12.3days and Maximum blooming period 22.3days (Singh and Beura 2002). In case of *Vriesea incurvata Gaudich*, (Bromeliaceae). The application of sucrose for 8 hours has been shown to enhance the preservation of color, brightness, and turgidity in floral scapes. It aids in maintaining water balance, reducing the loss of fresh weight, and extending the vase life of flowers by up to 24 days (Pulido *et al.*, 2021). Sucrose preserved a noticeably better water balance in spikes when combined with cobalt chloride and silver nitrate, which extended the vase life. Cobalt chloride is very effective at preventing vascular blockages that result from bacteria blocking vascular tissues. It has also been shown that treating iris spikes with CoCl<sub>2</sub> greatly extends their vase life (Singh and Sharma, 2003).

### **Importance of Chemicals.**

Chemical treatments considerably enlarged the spikes' floret diameter compared to the untreated control. Compared to the untreated control, the chemical treatments increased holding solution consumption and solution absorption to its maximum ((Singh and Sharma, 2003). The calcium effective in enzymes activation, enhances antioxidant activities, and delays senescence in gladiolus it also helps to increase diameter of xylem with good water flow it can delaying senescence (Ahmad and Rab, 2020). SA considered as the messenger of molecules it involves in disease in plants which response to pathogenic attack, it plays key role to tackle the environmental stress and plant growth development (Mohammadi *et al.*, 2013).

### **Effect of chemical**

In the research of Ahmed, 2020 the spikes were treated with vase holding solution of calcium of 0, 100, 200, 300 and 400 mM. The result and conclusion found that 200mM calcium delayed flower opening and fading, superior in quality retaining (Ahmed and Rab, 2020). In short communication of Kumar, suggest the calcium sulphate increases the vase life of flower spike (Kumar *et al.*, 2007). According to the Marandi, Salicylic acid (Acidifying agent ), also known as SA, is a basic compound found in plants that helps regulate various aspects of their growth and development. It plays a role in processes like the opening and closing of tiny pores called stomata, the sprouting of seeds, the absorption of essential minerals, the determination of plant gender, and the activation of defense mechanisms against diseases, The findings of this study indicate that among all the treatments, Acidifying agent salicylic acid (SA) had the most positive impact on the fresh weight percentage, solution uptake in terms of volume (cm<sup>3</sup>), and overall vase life of the cut flowers (Marandi *et al.*, 2011). Some studies have found that the application of salicylic acid can extent the vase life of flowers by reducing their respiration rate and inhibiting ethylene production.

Additionally, research has demonstrated that salicylic acid enhances the ability of flowers to withstand water deficit stress, resulting in improved vase life for cut rose flowers (Mehdikhah *et al.*, 2016). The use of sucrose in conjunction with antimicrobial agents (Germicide) (HQS and AgNO<sub>3</sub>) and an ethylene inhibitor (AgNO<sub>3</sub>) enhanced the carbohydrate content in flowers, improved water uptake, delayed senescence, preserved flower turgidity, decreased bacterial growth, and minimized xylem blockage. These combined effects resulted in superior flower quality and prolonged longevity (Mishra and Khanal, 2019).

### **Effect of Growth Regulators on physiological parameters**

Sable *et al.*, (2015) studied the the effect of foliar application of Plant growth regulators gladiolus Cv. H.B.Pitt, corms treated with Bavistin 0.1% and plants sprayed with GA<sub>3</sub>, NAA, CCC in different range it founds GA<sub>3</sub> at 200ppm was effective for elongation of cell and internodal length, height of plant and maximum no of leaves, increase the no. of florets and its weight, overall it enhance the growth and quality of flower (Sable *et al.*, 2015). Padmalatha *et al.*, (2015) Study was conducted on two cv. Darshan and Dhiraj it founds the influence of foliar application of BR(10ppm) along with GA<sub>3</sub> (150ppm) significant effect increase no. of corms and cormels at the stage of 3<sup>rd</sup> and 6<sup>th</sup> leaf stage, increase weight and size of corms, it can be suggest for multiplication of high-quality planting material and extending vase life of gladiolus (Padmalatha *et al.*, 2015).The use of Gibberellic acid on gladiolus corms has proven to be a successful method for overcoming dormancy and enhancing various aspects of plant development. This treatment has been found to positively impact vegetative growth, improve the quality of flowers, and contribute to the enlargement of both corms and cormels in gladiolus plants (Priyanka, Holkar *et al.*, 2018).

### **Conclusion**

Gladiolus is most demanding cut flower through out the world , however since flowers are perishable and the day by day gladiolus industry is growing, due to this more research into new technologies is necessary to address the severe decline in gladiolus. It has been shown that the aforementioned strategies prolong the flower's post-harvest life.

## References

1. Prabesh Koirala ; Rajiv Raman Neupane; Bishal Dura; Puspa Raj Poudel; Rukmagat Pathak (2023) Study on the Effect of Different Chemicals and Essential Oils on the Vase Life Of Gladiolus (Var American Beauty) Cut Flower, **[HTTP://www.researchgate.net/publication/369009267](http://www.researchgate.net/publication/369009267)**
2. Pratikshya lamichhanea\*, Pranisha Bhattaraia, Swastika Subedia, Januka Dahala, Janak Adhikarib (2022), Effects Of Different Treatments On Vase Life Of Gladiolus Cut Spikes: A Review, Tropical Agroecosystems (TAEC) 3(1) (2022) 12-15, ISSN: 2735-0274 (Online) CODEN: TARGDB, DOI: **<http://doi.org/10.26480/taec.01.2022.12.15>**.
3. Syed Atif Shaukat, Syed Zulfiqar Ali Shah, Syed Kashif Shaukat And Syed Waasif Shoukat (2013), Performance Of Gladiolus (*Gladiolus grandiflora* L.) Cultivars Under The Climatic Conditions Of Bagh Azad Jammu And Kashmir Pakistan, Journal of Central European Agriculture 2013 14(2), p.636-645, DOI: 10.5513/JCEA01/14.2.1244
4. Masood Ahmad\* And Abdur Rab (2020), Calcium Effects On Post-Harvest Attributes And Vase Life Of Gladiolus Using Different Methods Of Application, Pak. J. Bot., 52(1): 167-179, 2020. DOI: **[http://dx.doi.org/10.30848/PJB2020-1\(28\)](http://dx.doi.org/10.30848/PJB2020-1(28))**
5. Banti Kumar, Sunil Malik, Mukesh Kumar, M.K. Singh, and Vipin Kumar (2007), Postharvest quality and vase life of tuberose (*Polianthes tuberosa* Linn.)

- cut spikes as affected by vase chemicals, *Journal of Ornamental Horticulture*, 10 (4): 274-275, October-December 2007.
6. Mona Mehdikhah , Rasoul Onsinejad , Mohammad Nabi Ilkaee and Behzad Kaviani (2016), Effect of Salicylic Acid, Citric Acid and Ascorbic Acid on Post-harvest Quality and Vase Life of Gerbera (*Gerbera jamesonii*) Cut Flowers, *Journal of Ornamental Plants*, Volume 6, Number 3: 181-191, [b.kaviani@yahoo.com](mailto:b.kaviani@yahoo.com).
  7. Alaei, M. Babalar, R. Naderi, M. Kafi (2011), Effect of pre- and postharvest salicylic acid treatment on physio-chemical attributes in relation to vase-life of rose cut flowers, *Postharvest Biology and Technology* 61 (2011) 91–94, [www.elsevier.com/locate/postharvbio](http://www.elsevier.com/locate/postharvbio)
  8. Utsav Bhandari, Yekraj Budha (2022), A brief report on post-harvest handling of cut, loose and pot flowers, <https://www.researchgate.net/publication/358890233>
  9. Salman Mushtaq1 & 2, Ishfaq Ahmad Hafiz1, Muhammad Shahid Iqbal\*, 1, Syed Zia ul Hasan1, Muhammad Arif3, Sami Ullah2, Misbah Rasheed4 and Rizwan Rafique4 (2013), Studies On The Performance Of Some Exotic Gladiolus Cultivars Under Rain-Fed Conditions, *International Journal of Modern Agriculture*, Volume 2, No.3, 2013 Copyright © Zohdi Publisher ISSN: 2305-7246.
  10. Rasul Jalili Marandi1\*, Abbas Hassani1, Ali Abdollahi2 and Soyli Hanafi1 (2011), Improvement of the vase life of cut gladiolus flowers by essential oils, salicylic acid and silver thiosulfate, *Journal of Medicinal Plants Research* Vol. 5(20), pp. 5039-5043, 30 September, 2011 Available online at <http://www.academicjournals.org/JMPR> ISSN 1996-0875 ©2011 Academic Journals.
  11. H. Mehraj , M. Mahasen , T. Taufique, I.H. Shiam and A.F.M. Jamal Uddin (2013), Vase Life Analysis Of Yellow Gladiolus Using Different
  12. Vase Solutions, *J. Expt. Biosci.* 4(2):23-26, July 2013 ISSN 2223-9626 (Online), ISSN 2077-3358 (Print). Email: [jamal4@yahoo.com](mailto:jamal4@yahoo.com)
  13. Seid Hussen and Hassen Yassin (2013), Review on the impact of different vase solutions on the postharvest life of rose flower, *International Journal of Agricultural Research and Review* Vol. 1(2) pp 013-017, December, 2013 Copyright © 2013 Spring Journals.

14. Susan S. Han (1992), Role of Sucrose in Bud Development and Vase Life of Cut *Liatrix spicata* (L.) Willd, Hortscience 27(11):1198-1200. 1992, Hortscience, Vol. 27(11), November 1992.
15. P Kumar, LS Verma, G Sharma, Manisha Netam and Hemant Kumar (2020), Study The Effect Of Plant Growth Regulators On Vase-Life Of Gladiolus: A review, Journal of Pharmacognosy and Phytochemistry 2020; 9(6): 564-570, E-ISSN: 2278-4136 ISSN: 2349-8234 www.phytojournal.com JPP 2020; 9(6): 564-570.
16. Ranvir Singh and Shashikala Beura (2002), Postharvest Life of Gladiolus as influenced by Floral preservatives, Journal of Ornamental Horticulture, New series Vol. 5 (1) : 76-79 (2002).
17. Priyanka, S. Holkar , Chandrashekar, S. Y. , Hemanth Kumar, P. , Basavalingaiah And Ganapathi, M5 (2018), Effect of Gibberellic acid on Growth, Flowering, Flower Quality and Corm Yield of Gladiolus – A Review, Trends in Biosciences 11(24), Print : ISSN 0974-8431, 3211-3216, 2018.
18. T. Padmalatha, G. Satyanarayana Reddy and R. Chandrasekhar (2015), Effect of plant growth regulators on corm production and vase life in gladiolus, J. Hortl. Sci. Vol. 10(2):220-225, 2015.
19. Kazuo Ichimura (1998). Improvement of Postharvest Life in Several Cut
20. Flowers by the Addition of Sucrose, JARQ 32, 275-280 (1998).
21. Ajish Muraleedharan and J. L. Joshi (2017), Studies On Prolonging The Vase Life Of Rose Flowers With Citric Acid And Sucrose Solution, © 2017 JETIR April 2017, Volume 4, Issue 4. <http://www.jetir.org/>
22. Edwin Pulido, Raquel Rejane-Negrelle, Francine Lorena Cuquel (2021), Sucrose, salicylic acid and citric acid solutions to extent vase life of *Vriesea incurvata* Gaudich. (Bromeliaceae) floral scapes, Acta Agronómica, vol. 70, no. 1, pp. 27-34, 2021, <https://doi.org/10.14482/INDES.30.1.303.661>
23. Da Y. Park, Aung H. Naing, Trinh N., Jeung-Sul Han, In-Kyu Kang, Chang K. Kim (2017), Synergistic Effect of Nano-Silver with Sucrose on Extending Vase Life of the Carnation cv. Edun, Front. Plant Sci., 14 September 2017 Sec. Plant Breeding Volume 8 - 2017 | <https://doi.org/10.3389/fpls.2017.01601>
24. Mishra and A. Khanal (2019), Vaselife analysis of gladiolus using different vase solutions, Journal of Bioscience and Agriculture Research, 21(01), 1749-1754. <https://doi.org/10.18801/jbar.210119.213>

25. Gholam Abbas Mohammadi, Ali Salehi Sardoei, Mojghan Shahdadneghad (2013), Improvement of the vase life of cut gladiolus flowers by salicylic acid and Putrescine, International journal of Advanced Biological and Biomedical Research Volume 2, Issue 2, 2014: 417-426
26. Jyoti Verma, Parminder Singh (2021), Post-harvest Handling and Senescence in Flower Crops: An Overview, Agricultural Reviews, Volume 42 Issue 2 : 145-155 (June 2021),
27. Asmita Khanal (2014), a term Paper on: Post Harvest Handling And Factors Affecting Post Harvest Life of Gladiolus Flower, <https://studylib.net/doc/10135916/>
28. Sanjamveer S Vehniwal, Lord Abbey (2019), Cut flower vase life – influential factors, metabolism and organic formulation, Horticulture International Journal, Volume 3 Issue 6 - 2019
29. L. Mortensen and T. Fjeld.(1998), Effects of air humidity, lighting period and lamp type on growth and vase life of roses. *Scientia Horticulturae* 73: 229-23.
30. R. E. Hardenburg, A. E. Waltada, and C. Y. Wang. (1986), The commercial storage of fruits, vegetables, and florists and nursery stocks. USDA Agr. Handb. 66(REV):75-92
31. P.V. Singh and M. Sharma (2003), The Postharvest Life of Pulsed Gladiolus Spikes: the Effect of Preservative Solutions, Proc. XXVI IHC – Elegant Science in Floriculture Eds. Th. Blom and R. Criley Acta Hort. 624, ISHS 2003
32. J. A. M. Kromwijk, E. Meinen, T. A. Dueck (2014), The effect of elevated CO<sub>2</sub> on the vegetative and generative growth, and on stomata, International Symposium on Orchids and Ornamental Plants – Royal Flora 2011, Chiang Mai, Thailand, 2012-01-09/2012-01-13 –<https://doi.org/10.17660/ActaHortic.2014.1025.23>
33. <https://dfr.icar.gov.in/ForFarmers/InsectPests>

