

Quantification of Microplastics in Otammiri River, Imo State, Nigeria

Abstract

This study analyzed the physicochemical and heavy metal status and also quantified microplastics from Otammiri River, Imo state, Nigeria. The physicochemical and heavy metal analysis were carried out using procedures of Federal Environmental Protection Agency. The quantification of the microplastics from Otammiri River was carried out according to the National Oceanic and Atmospheric Administration (NOAA) protocol for surface water trawling for microplastics. The microplastics quantified in this study were at the range of 0.3 mm to 5mm that were resistance to wet oxidation and exhibited flotation in a 5M NaCl. The result of the physicochemical parameters showed that the parameters were within the Federal Ministry of Environment (FMEV) permissible limits with exception to lead and mercury. The microplastics concentrations at the surface water of Otammiri River was between 0.12 g to 2.50 g and in sediments samples, between 0.01 g to 1.37 g.

1.0 Introduction

Microplastics are commonly defined as plastic particles smaller than 5 mm in at least one external dimension (Miller *et al.*, 2021; United States Environmental Protection Agency, EPA, 2021). Studies have shown that plastic is abundance in inland freshwater systems in Nigeria (Nwakanma, Blessed & Hanson, 2021; Williams & Osahon, 2021; Olarinmoye *et al.*, 2021; Enyoh, Verla, Ohiagu, & Enyoh, 2021; Babatunde & Arinze, 2018). Enyoh, Verla, Verla, & Ihenetu (2019) reported that microplastics abundance from shore of Rivers in Nwangele Local Government Area of Imo state, ranged from 440 to 1,556 particles/L, with high accumulation at downstream. Olarinmoye *et al.*, (2021) showed that 5% of the potential MP particles identified by means of FTIR and pyrolysis GC-MS were polypropylene and polyethylene. The abundance of microplastics ranged from 310–2319 particles/kg in sediment, and 139–303 particles/L in water. Cross River, Qua Iboe River and Jaja Creek in Akwa Ibom, Nigeria, suspended marine litter was reported to be of high quantity of plastic (> 5000 kg/m²). Adeogun, Ibor, Khan, Chukwuka, Omogbemil & Arukwe (2022), screened and detected MPs in the stomach of commonly consumed fish species from a municipal water supply lake (Eleyele) in Nigeria. The physical effects of MPs on marine organisms include entanglement which may lead to death, injuries and blockage of digestive or intestinal pathways, lowering of energy of productive animals, decrease in food consumption due to ingested MPs (Li, Sun, Li, Tang, Miu & Ma, 2021). Exposure routes of microplastics to humans include ingestion, inhalation, and dermal penetration (Anqi & Wen-Xiong, 2023). Among all the exposure routes, the ingestion of microplastics was regarded as the primary route. MPs/g 0.44 of

Comment [A1]: The abstract provides a concise summary of the study's objectives, methodology, and key findings. However, it could benefit from a more structured approach. Consider using subheadings like "Background," "Methods," "Results," and "Conclusion" to enhance readability. The information about physicochemical and heavy metal analysis is too brief. Specific findings, particularly regarding lead and mercury exceeding permissible limits, should be mentioned.

Comment [A2]: The introduction covers a broad range of relevant literature and provides context for the study. Ensure that references are up-to-date and correctly formatted, as some citations appear incomplete or incorrect. Clarify the rationale behind choosing Otammiri River for the study and explicitly state why this river is significant for microplastic research.

nano and microplastics were found in sugar, 0.11 MPs/g were found in salt, 0.03 MPs/g were found in alcohol, and 0.09 MPs/g were found in bottled water (Lusher, McHugh, & Thompson, 2013). Humans could also assume an estimated intake of 80 g per day of microplastics via plants (fruits and vegetable) that accumulate MPs through uptake from polluted soil (Van Franeker, J& Law, 2015). Another study in 2019, showed that microplastics in a human biological sample (feces) 50 items/g and 9 plastic types were found in the fecal sample (Anqi & Wen-Xiong, 2023). Biofilms in drinking water contain loads of microbes which are transported with the water such as tap water causing serious health menace (Nworie, 2021). Liang, de Haan, Cerdà-Domènech, M´endez, Lucena, García-Aljaro, Sanchez-Vidal & Ballesté (2023) noted that these plastics with a higher concentration of recent fecal pollution can drift to areas with lower levels of pollution, increasing the total fecal bacteria load in the water. This article focused on the findings of the study regarding the presence, distribution, and composition of microplastics in Otammiri River surface water and sediment samples.

2.0 Methodology

Study Area

The Otammiri River is situated in Imo State, Nigeria. Geographically, its watershed spans between latitudes 5°17'N and 5°30'N, and longitudes 6°58'E and 7°04'E. Originating from Egbu, Owerri, the river flows southward, passing through significant locations like Nekede, Ihiagwa, Eziobodo, Obowuumuisu, Mgbirichi, and Umuagwo, until it reaches Ozuzu in the Etche Local Government Area of Rivers State. From there, it continues its journey to the Atlantic Ocean (Fagorite, Ahirakwem, Okeke, & Onyekuru 2019). Covering an area of about 10,000 km², the Otammiri watershed experiences an annual rainfall ranging between 2250 and 2500 mm. Predominantly, the watershed is characterized by depleted rainforest vegetation, with an average temperature of 27°C maintained throughout the year (Fagorite *et al.*, 2019).

Comment [A3]: The methodology section is detailed, covering various aspects of sample collection and analysis. However, include more information on the statistical methods used for analyzing the data. Ensure the descriptions of the protocols, such as the NOAA protocol for microplastic quantification, are clear and complete, including any modifications made to standard procedures.




Fig 1: Location and Accessibility Map of the Study Area

Collection Points of Samples

The location and accessibility points for samples collection from Otammiri River, Imo State Nigeria are listed below.

- a) Ekwuato 464115, Lat 5.334717⁰ and Long 6.965682⁰ Imo, Nigeria (downstream)
- b) Ihiagwa Otammiri Road, Ihiagwa 460113, Lat 5.388439⁰ and Long 6.987726⁰ (middle stream 1)
- c) Nekede 460106, Lat 5.442928⁰ and Long 7.022863⁰ (middle stream 2)
- d) Wetheral, Owerri, 460281, Lat 5.471891⁰ and Long 7.0416⁰ (upstream).

Collection of physicochemical and heavy metals samples from surface water of Otammiri

This study was carried out during the dry season (January, 2024). Samples were collected using procedures of Federal Environmental Protection Agency (FEPA 2003).

Collection of microplastics from surface water of Otammiri

The collection of surface water microplastic samples was conducted using a surface manta trawl, adhering to the protocol outlined by the National Oceanic and Atmospheric Administration (NOAA) for surface water trawling for microplastics (Bikker, Lawson, Wilson & Rochman, 2020; Kerubo, Muthumbi, Onyari, Kimani & Robertson-Andersson, 2020). Samples were gathered against the water current, with the manta trawl equipped with a mesh net size of 0.335 mm and an opening measuring 25 × 60 cm.

Collection of microplastics from sediment samples of Otammiri

The sediment samples were obtained utilizing a grab sampler, as described by Miller *et al.* (2021) and deposited into clean glass jars. At each sampling point, five sediment samples were collected, ensuring representation across the area.

Quantification of Microplastics from Otammiri River

The quantification of microplastics from the Otammiri River followed the protocol established by the National Oceanic and Atmospheric Administration (NOAA) for surface water trawling (Bikker *et al.*, 2020; Kerubo *et al.*, 2020).




Fig 2: Flow diagram for the analysis of microplastics in water samples (Miller *et al.*, 2021).




Fig 3: Flow diagram for the analysis of microplastics in sediment samples (Miller *et al.*, 2021).

3.0 **Result**

Physicochemical and Heavy Metal Parameters of Otammiri River

From table 1, it was observed that the surface water at Ekwuato location (464115) was clear; Ihiagwa (460113), sparsely loaded with debris; Nekede (460106), heavily loaded with debris and Wetheral (460281) (the upstream), clear. Their sediment samples at Ekwuato Location (464115) were sparsely loaded with debris, Ihiagwa (460113), loaded with debris, Nekede (460106), and heavily loaded with debris while Wetheral location (460281) was clear. Odour was unobjectionable across the sampling points.

Table 2, showed that pH for the water sample ranged from 6.57 ± 0.04 (Wetheral 460281) to 7.42 ± 0.05 (Nekede). Temperature of the water ranged from 20.50 ± 0.14 (Wetheral 460281) to 22.70 ± 0.14 (Nekede 460106) while temperature for the sediment samples ranged from 19.70 ± 0.14 (Wetheral) to 20.60 ± 0.28 (Ekwuato). The surface water conductivity had least and highest values at 16.10 ± 0.28 (Ekwuato), 25.10 ± 0.42 (Wetheral) and Nekede (28.50 ± 0.42). There is no significant difference ($P \leq 0.05$) in the pH, temperature and conductivity values of the water and sediment samples across the sampling locations.

The turbidity levels of Otammiri River at various sampling points were assessed, revealing notable variations. Ihiagwa exhibited the highest turbidity levels (17.6 ± 1.41 NTU), followed by Nekede (6.17 ± 0.49 NTU), Ekwuato (2.93 ± 0.32 NTU), Wetheral (1.7 ± 0.17 NTU). Statistical analysis indicated no significant difference ($P \leq 0.05$) in turbidity levels among the sampling locations.

Comment [A4]: The results section presents comprehensive data on physicochemical parameters, heavy metals, and microplastic concentrations. Use more visual aids like graphs and tables to make the data more accessible and easier to interpret. Some figures and tables need better formatting and labeling for clarity. Provide a more detailed statistical analysis of the data, including significance testing where applicable.

Table 1: Appearance and odour of Otammiri Surface water and Sediment at their sampling locations


	FMEnv Standard	Location	Ekwuato (464115)	Ihiagwa (460113)	Nekede	Wetheral
Appearance	Clear	Surface	Clear	sparse loaded with debris	heavily loaded with debris	Clear
		Sediment	sparse loaded with debris	loaded with debris	heavily loaded with debris	heavily loaded with debris
Odour	Unobjectionable	Surface	Unobjectionable	Objectionable	Objectionable	objectionable
		Sediment	Unobjectionable	Unobjectionable	Unobjectionable	Unobjectionable

Table 2: pH and Temperature Conductivity and Turbidity of Otammiri River

		FMEnv Standard	Ekwuato (464115)	Ihiagwa (460113)	Nekede	Wetheral
Water	Ph	6.5 – 8.50	7.15±0.12	7.21±0.13	7.42±0.05	6.57±0.04
	Temperature (°C)	-	22.30±0.14	22.35±0.07	22.70±0.14	20.50±0.14
	Conductivity (µS/cm)	1000	16.10±0.28	23.75±5.02	28.50±0.42	25.10±0.42
	Turbidity (NTU)	10.00	2.93±0.32	17.6±1.42	6.17±0.49	1.70±0.17
Sediment	pH	6.5 – 8.50	7.03±0.20	7.18±0.10	7.81±0.45	6.81±0.18
	Temperature (°C)	-	20.60±0.28	20.4±0.14	20.35±0.21	19.70±0.14
	Conductivity (µS/cm)	1000	10.80±0.14	19.55±1.77	50.10±3.25	24.50±0.57

± Standard deviation; µS/cm- microSiemens per centimeter; NTU- Nephelometric Turbidity Units

UNDER PEER REVIEW



Hardness, Total Solids, Total Dissolved Solids, Total Suspended Solids, Chemical Oxygen Demand, Biochemical Oxygen Demand

The result from table 3 showed that Otammiri River had the highest values for “Hardness” (mg/l) at Nekede (60.00 ± 7.07), and least at Ekwuato (40.00 ± 5.66). The sediment samples recorded highest values for “Hardness” at Nekede (80.00 ± 5.66), and least at Wetheral (35.00 ± 4.24) locations. Total Solid, (mg/l) in the water samples revealed to be higher in Nekede (6.10 ± 0.65) and least at Ekwuato (4.56 ± 0.32). The sediment samples showed to have higher total solids at Ihiagwa (12.47 ± 0.92), and least at Wetheral (9.39 ± 0.32) locations. Total Suspended Solid (mg/l) in the water samples revealed to be higher at Ihiagwa, (5.66 ± 0.29) and least at Ekwuato (1.35 ± 0.07) Locations. The sediment samples showed to have higher Total Suspended Solid at Nekede (4.60 ± 0.40), and least at Wetheral (3.19 ± 0.96).

The table 3 also revealed Chemical Oxygen Demand, COD (mg/l) of the water samples to be higher in Nekede location (178.67 ± 33.94) and least at Ekwuato (165.34 ± 60.34) locations. COD (mg/l) of the sediment samples was higher in Wetheral location (124.00 ± 5.66) and least at Ekwuato (76.00 ± 1.88) locations. The Biochemical Oxygen Demand, BOD (mg/l) result of the water samples showed that Nekede location (165.45 ± 5.44) had higher BOD and least at Wetheral (128.70 ± 18.24).

There is no significant difference ($P \leq 0.05$) across the sampling locations for both water and sediment samples.

Table 3: Hardness, Total Solids, Total Dissolved Solids, Total Suspended Solids, Chemical Oxygen Demand, Biochemical Oxygen Demand

	Water					Sediment				
	FMEnv Standard	Ekwuato	Ihiagwa	Nekede	Wetheral	FMEnv Standard	Ekwuato	Ihiagwa	Nekede	Wetheral
Hardness (mg/l)	200.00	40.00±5.6	54.00±2.83	60.00±7.07	41.00±1.41	-	62.00±4.24	64.50±3.53	80.00±5.66	35.00±4.24
Total Solid, (mg/l)	500.00-1000.00	4.56±0.32	5.62±0.45	6.10±0.65	4.70±0.08	-	9.86±0.42	12.47±0.92	9.58±0.18	9.39±0.32
Total Dissolved Solid (mg/l)	500.00	2.53±0.01	3.99±0.50	4.72±0.09	2.67±0.19	-	5.15±0.81	6.61±0.70	6.67±0.55	7.87±0.52
Total Suspended Solid (mg/l)	<10.00	1.35±0.07	5.66±0.29	4.47±0.60	2.13±0.21	-	3.36±0.35	4.24±1.13	4.60±0.40	3.19±0.96
COD (mg/l)	NS	165.34±60.34	173.14±37.88	178.67±33.94	172.00±35.82	NS	76.00±1.88	91.03±1.94	94.67±16.97	124.00±5.66
BOD (mg/l)	NS	130.15±0.78	155.50±9.69	165.45±5.44	128.70±18.24	NS	126.80±3.96	158.96±40.54	176.00±40.73	127.40±25.74

± Standard deviation; mg/l: milligram /liter; NS- Not Stated

Chlorine, Nitrate, Nitrate, Phosphorous, Sulphate and Fluoride

The result from table 4 showed that nitrate (mg/l) was found to be higher at Ihiagwa, (4.11 ± 0.14) and least at Ekwuato location (1.49 ± 0.70). The sediment samples recorded highest values for at Nekede (6.45 ± 0.09), and least for Wetheral (4.77 ± 0.35). Nitrite (mg/l) was found in the water samples to be higher at Ihiagwa, (0.67 ± 0.01), and least at Wetheral (0.24 ± 0.04). The sediment samples recorded highest values at Ihiagwa, (3.75 ± 0.31) and least values at Wetheral (2.64 ± 0.33). Phosphorous (mg/l) was found in the water samples to be higher at Nekede (2.00 ± 0.07) and least at Wetheral (1.54 ± 0.07). The sediment samples recorded highest values at Ekwuato (2.61 ± 0.07) and least at Nekede (2.37 ± 0.14). Sulphate (mg/l) was found in the water samples to be higher at Nekede (13.37 ± 0.87), and least at Ekwuato (11.31 ± 2.04). The sediment samples recorded highest values at Nekede (86.62 ± 1.45) and least at Wetheral (45.68 ± 1.75). Fluoride (mg/l) was found in the water samples to be higher at Nekede (1.06 ± 1.00) and least at Ekwuato (0.23 ± 0.07). The sediment samples recorded highest values at Nekede (2.23 ± 0.24).

The heavy metals analysis shown in table 5 revealed cadmium (mg/l) to be 0.01 for both water and sediment samples across the sampling locations. Lead (mg/l) in the water samples ranged from Ekwuato (0.01 ± 0.00) to Wetheral location (7.00 ± 9.90). The sediment lead concentration was between 0.01 ± 0.01 (Wetheral and Ihiagwa locations) and 0.02 ± 0.00 (Ekwuato location). Mercury (mg/l) in the water samples was between 0.01 ± 0.00 (Wetheral location) and 0.03 ± 0.00 (Nekede and Ihiagwa locations). Arsenic (mg/l) in the water samples had higher values in Ekwuato location (0.04 ± 0.05) while the sediment samples had the highest arsenic values in Wetheral location (0.04 ± 0.01). There is no significant difference ($P \leq 0.05$) across the sampling locations for both water and sediment sample.

Table 4: Chlorine, Nitrate, Nitrite, Phosphorous, Sulphate and Fluoride

	Water					Sediment				
	FMEnv Standard	Ekwuato	Nekede	Ihiagwa	Wetheral	FMEnv Standard	Ekwuato	Ihiagwa	Nekede	Wetheral
Chlorine (mg/l)	250.00	95.00±2.8	112.00±2.23	104.00±2.83	94.5±4.95	NS	155.00±2.83	142.50±17.67	160.00±1.41	79.00±4.24
Nitrate (mg/l)	50.00	1.49±0.70	3.76±0.28	4.11±0.14	2.24±0.18	20.00	5.96±0.48	6.21±1.06	6.45±0.09	4.78±0.35
Nitrite (mg/l)	0.30	0.33±0.02	0.58±0.01	0.67±0.01	0.24±0.04	NS	3.13±0.59	3.75±0.31	3.53±0.64	2.64±0.33
Phosphorous (mg/l)	5.00	1.78±0.02	2.00±0.07	1.92±0.07	1.54±0.07	>100.00	2.61±0.07	2.29±0.22	2.37±0.14	2.39±0.13
Sulphate (mg/l)	200-400.00	11.32±2.04	13.37±0.87	13.16±0.00	11.92±0.00	100.00	51.44±1.75	71.88±2.38	86.62±1.45	45.68±1.75
Fluoride (mg/l)	NS	0.23±0.07	1.06±1.00	0.60±0.03	0.30±0.08	NS	0.47±0.15	2.23±0.24	2.18±0.06	0.47±0.15

± Standard deviation; mg/l: milligram /liter; NS- Not Stated

Table 5: Heavy Metal Analysis of Otammiri River

	Water					Sediment				
	FME _{env} Standard	Ekwuato	Ihiagwa	Nekede	Wetheral	FME _{env} Standard	Ekwuato	Ihiagwa	Nekede	Wetheral
Cadmium (mg/l)	0.01	0.01±0.00	0.01±0.00	0.01±0.01	0.01±0.01	0.01	0.01±0.00	0.01±0.00	0.01±0.01	0.01±0.00
Lead (mg/l)	0.05	0.01±0.00	0.01±0.01	0.03±0.01	7.00±9.90	0.05	0.02±0.00	0.01±0.01	0.01±0.01	0.01±0.01
Mercury (mg/l)	0.01	0.02±0.00	0.03±0.00	0.03±0.00	0.01±0.00	0.001	0.04±0.00	0.03±0.00	0.04±0.01	0.04±0.00
Arsenic (mg/l)	0.2	0.04±0.05	0.03±0.00	0.02±0.00	0.02±0.00	0.05	0.03±0.00	0.03±0.00	0.02±0.00	0.04±0.01

± Standard deviation; mg/l: milligram /liter

Total Solutes of Samples trawled from surface water and sediment of Otammiri River for Microplastics studies

As shown in fig 1, the total solids for the surface water ranged from Ekwuato, 2.24 ± 0.28 g to Ihiagwa, 49.27 ± 1.41 g.

As seen in fig 2, the sediments total solids ranged from Wetheral, 61.6 ± 18.67 g to Ekwuato, 185.17 ± 15.34 g.

Concentration of Microplastics in the Surface Water

Examining the concentration of microplastics in surface water across the Otammiri sampling points in fig 3, followed the order: Nekede > Ihiagwa > Wetheral > Ekwuato. There is no significant difference between the microplastic concentrations found in Otammiri surface water at the different sampling points.

Correlation between Turbidity of Otammiri River and the microplastics concentration

Fig 4, "Correlation between Turbidity of Otammiri River and the microplastics concentration", showed that the microplastics at the surface of the river followed the trend of the turbidity levels. The downstream had the least turbidity and microplastics values (2.93 ± 0.32 NTU and 0.12 ± 0.01 g) while the middle stream had the highest turbidity and microplastics values (17.60 ± 1.42 NTU and 2.50 ± 1.56 g) respectively. At $P \geq 0.05$ statistical level, correlation statistical result showed that increment in turbidity levels is directly proportional to the concentrations of the microplastics at the different sampling points as the middle had the highest concentration of the microplastics.

Concentration of Microplastics in the Sediment Samples

Similarly, the concentration of microplastics in sediment samples across the Otammiri sampling points as shown in fig 5 followed the order: Ihiagwa > Nekede > Ekwuato > Wetheral. There is a significant difference ($P \geq 0.05$) in the microplastics concentration at the different sampling points.




Fig 4: Total Solids of samples trawled from Otammiri River surface water from different sampling points

g: grams.




Fig 5: Total Solids of sediment samples from Otammiri River from different sampling points

g: gram




Fig 6: Concentration of Microplastics in Surface Water across Otammiri River Sampling Points

± Standard error of mean g: grams Lat: Latitude Long: Longitude




Fig 7: Correlation between Turbidity of Otammiri River and the microplastics concentration

NTU: Nephelometric Turbidity Units

g: gram




Fig 8: Concentration of Microplastics in Sediment Samples across Otammiri River Sampling Points

± Standard error of mean g: grams Lat: Latitude Long: Longitude g: gram

Discussion

Physicochemical parameters including pH, temperature, dissolved oxygen levels, turbidity, conductivity and even nutrient concentration which are influenced by various natural and anthropogenic activities shapes the ecological characteristics of river systems. These factors also influence microplastics abundance in river ecosystem. The appearance of the surface water was clear at the downstream and upstream but was heavily loaded with debris at the middle stream (Ihiagwa and Nekede) with objectable odour as well. This indicates the presence of pollution in this water body at these points which do not confirm to safe drinking water standards (WHO, 2011).The pH of the Otammiri river and sediments were at acceptable limits of the Federal Ministry of Environment (FMEV) (6.5 – 8.50). The upstream (Wetheral location) surface water and sediment were observed to be slightly acidic while the middle and downstream were observed to be slightly alkaline. In acidic conditions, certain plastics may degrade more rapidly due to hydrolysis or other chemical reactions. Conversely, in alkaline conditions, degradation might be slower. This can influence the rate at which macroplastics break down into microplastics (Andrady, 2011).

The temperature values of the surface water and sediments were within the ambient temperature range stipulated by FMEV which is in agreement with the findings of Cosmas,

Comment [A5]: The discussion effectively links the findings to broader environmental and health implications. However, discuss the potential sources of microplastics and heavy metals in more detail, such as industrial activities, urban runoff, and other anthropogenic factors. Compare the findings with other similar studies globally and regionally to highlight the significance of the results. Address any limitations of the study and suggest areas for future research.

Ahamefula & Ahiarakwem (2015) and Okeke & Adinna (2013) in Otammiri River. The temperature values reported were due to the climatic condition of the study area (Eze, Onwukeme, Ogbuagu, Aralu, Aduaka & Okoro, 2024; Ubaka, Ekwonu & Ogah 2019; Okechi & Chukwura, 2020). Warmer temperatures can accelerate the degradation of plastic materials, leading to increased fragmentation and subsequent release of microplastics into the environment (Moore, 2008). Temperature influences the metabolic rates and activities of microbial communities responsible for the degradation and decomposition of organic matter, including plastics (Zettler, Mincer & Amaral-Zettler, 2013). Higher temperatures can enhance microbial colonization and enzymatic breakdown of microplastics, potentially affecting their abundance and persistence in aquatic environments (Wright, Erni-Cassola, Zadjelovic & Gibson, 2020).

The mean electrical conductivity values of the surface water and sediments were within the acceptable FMEV limit ($1000 \mu\text{scm}^{-1}$). This infers that there is less impact of dissolved ions in both the surface water and sediment samples. Electrical conductivity can influence the transport and distribution of microplastics. For example, higher conductivity levels may indicate higher concentrations of dissolved salts, which could affect the buoyancy and settling rates of microplastic particles (Enders, Lenz, Stedmon & Nielsen, 2015). This can indirectly influence the abundance and distribution of microplastics in aquatic environments by altering their movement and deposition patterns.

The mean total hardness (mg/l) values of the surface water and sediment were below the recommended limit stipulated by FMEV. The middle streams (Nekede and Ihiagwa) were observed to have higher values for total hardness than the upstream and downstream. The hardness in water is due to the presence of cations such as Ca^{2+} and Mg^{2+} , anions such as HCO_3^- , Cl^- and SO_4^{2-} in surface water. Studies conducted by Akubugwo, Nwachukwu & Odika (2013) and Eze *et al.* (2021) in Njaba River, Imo State, Nigeria, reported a similar finding. Hard water contains higher concentrations of these ions, which can interact with microplastic surfaces through electrostatic attractions and facilitate the aggregation of microplastic particles with suspended solids and organic matter (Hüffer, Hofmann & Griebler, 2017). This aggregation process may affect the transport and settling of microplastics in aquatic environments, potentially influencing their overall abundance.

The mean suspended solids values in both the surface water and sediment were lower than the permissible limit in the upstream and downstream locations but marginally increased in

the middle stream. This informs the high microplastics concentration observed in middle stream. The effect of consuming water with high solids has been reported by Akubugwo, Nwachukwu & Odika (2013). Total suspended solids in water can serve as carriers for microplastic particles, facilitating their transport and aggregation. Microplastics can adsorb onto suspended solids through physical interactions and surface properties, leading to the formation of larger aggregates that settle out of the water column (Nizzetto, Futter & Langaas, 2016).

The level of chemical oxygen demand (COD) in the sampling points exceeds the 10 mg/l COD of the maximum permissible limit for drinking water and aquatic life. COD determines the amount of oxygen required for chemical oxidation of organic compounds. High levels of COD can lead to death of fishes and can cause dysentery in humans who use the polluted water. The findings are in tandem with the work of Amadi, Olasehinde, Okosun, & Yisa (2010) in his quality assessment of Otammiri River. High COD levels can impact the composition and activity of microbial communities in water bodies. Variations in COD levels may alter microbial processes involved in the breakdown of organic matter and plastics, indirectly influencing the persistence and fate of microplastics in aquatic ecosystems (Pittura, Avio, Giuliani, d'Errico, Keiter, Cormier... Regoli, 2020).

The result obtained from this study showed biochemical oxygen demand (BOD) values to exceed the recommended maximum allowable concentration of the European Union for good quality water for aquaculture (3.0 – 6.0 / 3-5 (pmm). Unpolluted water usually has values 2 or less whereas those receiving waste water/discharge may have values up to 10 mg/l or more (Amadi, Olasehinde, Okosun, & Yisa 2010). The BOD values are high probably due to the discharge of municipal wastes and poorly executed agricultural activities near the river banks which was observed during the survey of the study site. Okeke & Adinna (2013) observed similar values in their study. Elevated BOD levels may enhance microbial activity and organic matter availability, potentially affecting the sorption and degradation rates of microplastics in aquatic ecosystems (Su, Xue, Li, Yang, Kolandhasamy, Li & Shi, 2019). Microorganisms play a crucial role in the degradation and biotransformation of organic pollutants and microplastics in water bodies (Zhang, Zhang, Qian & Jiang, 2019). Changes in BOD levels can alter the composition and activity of microbial communities, indirectly influencing the degradation and fate of microplastics in aquatic environments (Tian, Jiang, Liu, Chen, Shen, & Zhu, 2021).

The nitrate mean values were also observed to be below permissible levels of FMEV. This shows that there is no much runoffs from the uplands into the river due to the dry season. This study is at variance with the study conducted in wet season by Joseph *et al.* (2019) where high concentration of nitrate was observed. Nitrite pollution can also affect the composition and activity of microbial communities in aquatic environments (Wang, Xie, Zhao & Liu, 2018). Changes in nitrite levels may therefore influence microbial processes involved in the breakdown and fate of microplastics, potentially affecting their abundance and persistence in water bodies. High concentrate of nitrate in water can cause the development of methaemoglobinaemia (blue baby syndrome) in infants (Joseph, Majesty, Friday, 2019). Increased nutrient availability due to nitrate pollution may stimulate algal growth and primary productivity, leading to the formation of biofilms and organic aggregates in which microplastics can become entrapped or sorbed (Lares, Ncibi, Sillanpää & Sillanpää, 2018).

Phosphate mean values were below the permissible standards. Changes in phosphate levels may influence microbial processes involved in the breakdown and fate of microplastics, potentially affecting their abundance and persistence in water bodies (Zhang *et al.*, 2019).

Sulphate mean values are lower than the permissible limits by FMEV Standard, although the river sediment had higher values for sulphate than the surface water. Drinking water containing higher amounts of magnesium sulphate or sodium sulphate may lead to intestinal discomfort, diarrhea and consequent dehydration especially in drinking water containing >500 mg/l of sulphate (Al-Wabel, Usman, El-Saeid, 2016). Sulfate ions can participate in chemical reactions and interactions with microplastic surfaces. Changes in sulfate concentrations may affect the availability and speciation of metal ions and other contaminants that can adsorb onto microplastic particles (Sun, Wang, Han, Jin, Yu, Zhao & Shi, 2021). These chemical interactions may influence the fate and bioavailability of microplastics in aquatic ecosystems, potentially impacting their abundance and ecological effects.

The mean values of the studied heavy metals were within the permissible limits in the upstream (Wetheral) that recorded 7.00 ± 9.90 mg/l and mercury that had values that exceeded the permissible limits in all sampled locations. This could result from weathering of parent rocks or anthropogenic activities (Okechi & Chukwura, 2020). It is important to note that these heavy metals may be part of the leachate from waste dumps and landfills around the sampling points which were later washed down the Otammiri River. The heavy metals also

owe their sources mainly from discharged electronic wastes (e-wastes)(Okechi & Chukwura, 2020). Heavy metals can adsorb onto the surface of microplastic particles in aquatic environments due to electrostatic attractions and surface properties (Xue *et al.*, 2020). This sorption process can lead to the accumulation of heavy metals on microplastics, potentially altering their distribution and behavior in water bodies. Conversely, microplastics can also serve as carriers for heavy metals, facilitating their transport and bioaccumulation in aquatic organisms (Song *et al.*, 2021). Heavy metals sorbed onto microplastics may undergo remobilization under changing environmental conditions, leading to the release of heavy metals back into the water column (Hüffer, Hofmann & von der Kammer, 2017). This remobilization process can contribute to the redistribution of heavy metals and microplastics in aquatic ecosystems, potentially increasing their exposure and ecological impacts on aquatic organisms (Sun *et al.*, 2021). Heavy metals adsorbed onto microplastic surfaces may become more bioavailable to aquatic organisms, leading to potential toxicity and ecological risks (Li *et al.*, 2019). The presence of heavy metals on microplastics can affect their interactions with biota and influence physiological responses in exposed organisms (Luís, Ferreira, Fonte, Torres & Soares, 2020).

The turbidity of the Otammiri River was between 1.7 ± 0.17 NTU and 17.6 ± 1.41 NTU. Ihiagwa and Nekede locations (middle streams) had higher turbidity levels. This could be attributed to the heavy anthropogenic activities (sanding mining and waste dumping) in the Ihiagwa and Nekede axis of Otammiri River. This is in consonance with the report of Bashir, Kimiko, Mak, Fag & Goncalves (2021) who stated that the turbidity of freshwaters around busy metropolis tends to increase due to high anthropogenic activities. The increment in turbidity levels directly proportional to the concentrations of the microplastics the different sampling points as the middle had the highest concentration of the microplastics. Research has shown that higher turbidity levels are associated with increased microplastics abundance (He, Han, Zheng, Yue, Li, Wu & Li, 2021; Ziajahromi, Kumar & Neale, 2020). Microplastics concentrations of Otammiri River surface water was between 0.12 g to 2.50 g. Nekede and Ihiagwa locations (middle streams) had the highest concentration of microplastic sampled (2.50 g and 0.59 g respectively). This may be attributed to the industrial and domestic wastewater discharged into the Otammiri River at that point. The upstream sampling point at which less microplastic concentration was seen, was located at Wetheral station. This is to show that anthropogenic activities are far below the levels obtainable in the middle streams. The upstream had higher microplastics concentrations than the downstream (upstream, 0.39

g; downstream, 0.39 g), denoting that the microplastics did not travel long distances as the heavy concentration seen in the middle was far less in the downstream. This can be attributed to the self-filtration system of water bodies as noted by Toumi, Abidli & Bejaoni, (2019). The lack of long-distance carriage of the microplastics may be due to the low flow velocity of the river. He, Smith, Egodawalta, Ayoko, Rimtoul & Goonetilleke (2021) had stated that high water velocity was required to transport more microplastics along a river course. The lack of long-distance carriage observed in the current research may also be due to the general pollution of the river with many large-sized items, which may be absorbing microplastics onto their surfaces, thereby limiting their transport along the river. The spatial variations of the microplastics abundance in the water samples vividly reveals the influence of anthropogenic activities around the sampling areas. This finding is in line with the study of Olarinmoye, Stock, Scherf, Whenu, Asenime & Ganzallo (2021) who noted that the microplastics load in rivers are governed by their proximity to urban areas and human activities. In terms of flowing systems in Nigeria, studies (Gideon, Adewumi, Kehinde, Michael, Temitop, Oluwatoyin, Oluwatosin, & Ademola, 2024 – Osun River; Ogbomida, Obazele, Aganmwonyi, Chukwuka, Emeribe & Omoigberale, 2023 - Ikpoba Rivers, Edo State; Aliyu, Okunola, Awe & Musa., 2023 - Kaduna; Nwonumara, Okoro, & Okogwu, 2021 – Ndibe, Cross River; Ebere Wirnkor, Verla & Ihenetu, 2019) have reported microplastics abundances in river water samples.

In particular, the levels of microplastics found in Otammiri is cause to worry. Not only does this water serve for domestic and agricultural purposes, most community dwellers depend on the fish for its protein needs. It may therefore be inferred that local people in the area are directly and in advertently exposed to microplastics through domestic use and consumption of the aquatic animals. The high-level microplastic pollution of Otammiri River may not be unconnected with the wide use and indiscriminate disposal of plastic materials, particularly the single-use ones (Yalwaji, John-Nwagwu & Sogbanmu, 2022). This situation is exacerbated by the lack of specific laws and policies prohibiting single-use plastics in Nigeria and non-enforcement of already established environmental protection laws.

The sampled sediments of Otammiri had microplastics concentration at the range of (0.01 g to 1.37 g). Microplastics concentrations in the Otammiri sediments were generally lower than the water samples. However, the spatial distribution pattern is similar to that observed for the water samples. Concentration of the microplastics increased from the upstream to the middle stream, and thereafter decreased at the downstream. This finding supports the fact that

sediments are a sink for MPs (Olarinmoye, Stock, Scherf, Whenu, Asenime & Ganzallo, 2021). Florian, Joris, Ian & Michael (2020) noted that turbidity currents potentially distribute and bury large quantities of microplastics in seafloor sediments. Overall, the spatial variation of the microplastics in the sediments reflects the impact of Owerri city's poor plastic waste management on the sediments microplastics contents. Eze, Anaebonam, Nweze, Onyemeka, Frank-Ogu, Justice-Alucho & Chiroma (2021) reported that due to available space along Otammiri Riverbank, the Owerri Municipal Council in 2000 approved the site as the official dumpsite for most of the solid waste collected from the municipality. Consequently, solid waste has been dumped at this site for more than ten years, on a surface area approximately 11 hectares in size, 6 meters high and not compacted and capped. Nearly 30 tons of commingled wastes are dumped here daily. Vast majority of Nigeria people also dispose plastics indiscriminately in the environment, where they degrade gradually over time, to release plastic fragments of various sizes. Such wastes are carried by run-off into streams and rivers, where they also breakdown to release small plastic particles. Ting, Haiqing, Fang, Fuhong, Youjun, Hao, Dongxia, Xingxuan, & Qianqian, (2023) noted that microplastics abundance in waterbodies is closely related to urbanization, human industrial activities, and wastewater impacts.

4.0 Conclusion

The physicochemical status of the Otammiri River is within the acceptable limits but high presence of lead and mercury is a cause for public health concern. The microplastics results revealed varying quantities of microplastics in Otammiri River surface water (ranging from 0.12 g to 2.50 g) and sediment (ranging from 0.01 g to 1.37 g). This information underscores the significant issue of plastic pollution, particularly the presence of microplastics, which pose substantial threats to aquatic ecosystems and human health due to their persistent, bioaccumulative nature and ability to adsorb harmful pollutants.

References

- Adeogun, A., Ibor, O., Khan, A., Chukwuka, A., Omogbemi, E. & Arukwe, A. (2022). Detection and occurrence of microplastics in the stomach of commercial fish species from a municipal water supply lake in southwestern Nigeria. *Environmental Science and Pollution Research*, 27:31035–31045.
- Aliyu, A. O., Okunola, O. J., Awe, F. E. & Musa, A. A. (2023). Assessment of microplastics contamination in River water, bottled water, sachet water and branded table salt

Comment [A6]: The conclusion summarizes the main findings and their implications succinctly. Reinforce the need for regulatory measures and public awareness campaigns to address microplastic pollution. Highlight any immediate actions that can be taken based on the study's findings.

Comment [A7]: Ensure the manuscript is free from grammatical errors and awkward phrasing. Consider professional editing to enhance readability. Maintain consistency in terminology and units of measurement throughout the manuscript. Ensure all figures and tables are of high quality, properly labeled, and referenced in the text. Consider adding captions that fully explain the content without needing to refer back to the text. Overall, the manuscript presents significant findings on microplastic pollution in the Otammiri River. By addressing the suggestions above, the manuscript can be improved in clarity, readability, and scientific rigor.

- samples in Kaduna Metropolis, Nigeria. *J. Appl. Sci. Environ. Manag.* 27 (6):1105–1118.
- Al-Wabel, M. I., Usman, A. R. A. & El-Saeid, M. H. (2016). Levels, Sources, and Risk Assessment of Polychlorinated Biphenyls (PCBs) in Soils from Industrial Areas: A Case Study from Saudi Arabia. *Polycyclic Aromatic Compounds*, 38(5): 420-433.
- Akubugwo, E. I., Nwachukwu, M. I. & Odika, P. C. (2013). Water quality assessment of Njaba River, Nigeria. *Journal of Environmental Science, Toxicology and Food Technology*, 4(6): 33-37.
- Amadi, A. N., Olasehinde, P. I., Okosun, E. A. & Yisa, J. (2010). Assessment of the water quality index of Otamiri and Oramiriukwa Rivers. *Physics International Journal*, 1(2):116 – 123.
- Andrady, A. L. (2011). Microplastics in the marine environment. *Marine Pollution Bulletin*, 62(8):1596-1605.
- Anqi, S. & Wen-Xiong, W. (2023). Human Exposure to Microplastics and Its Associated Health Risks. *Environment and Health*, 1:139–149
- APHA, (2005). Standard Methods for the Examination of Water and Wastewater 19 (American Public Health Association, APHA, AWWA Washington. DC.
- Babatunde, B. B. & Arinze, O. U. (2018). Suspended marine litter in Akwa Ibom state, Nigeria: A case study of cross river, QUA Iboe river and Jaja Creek. *Journal of Geography, Environment and Earth Science International*, 18: 1-13.
- Bashir, S. M., Kimiko, S., Mak, C. W., Fang, J. & Gonçalves, D. (2021). Personal care and cosmetic products as a potential source of environmental contamination by microplastics in a densely populated Asian City. *Front. Mar. Sci.* 8:683482
- Cosmas, A., Ahamefula, S. & Ahirakwem, C. (2015). Seasonal Variations in physiochemical and Bacteriological Parameters of Rivers. *Journal of Environmental Protection*, 5: 1094-1110.
- Ebere, C. E., Wirnkor, V. A., Verla, E. V. & Ihenetu, S. C., (2019). Macrodebris and microplastics pollution in Nigeria: first report on abundance, distribution and composition. *Environ. Anal. Health Toxicol.* 34 (4), e2019012

- Enders, K., Lenz, R., Stedmon, C. A., & Nielsen, T. G. (2015). Abundance, size and polymer composition of marine microplastics $\geq 10\mu\text{m}$ in the Atlantic Ocean and their modelled vertical distribution. *Marine Pollution Bulletin*, 100(1):70–81.
- Enyoh, C. E., Verla, A. W., Verla, E. N. & Ihenetu, S. C. (2019). Macrodebris and microplastics pollution in Nigeria: first report on abundance, distribution and composition. *Environmental Analysis Health and Toxicology*, 34(4): 1-15.
- Enyoh, C. E., Verla W. A., Ohiagu, F. O. & Enyoh, E.C. (2021): Progress and future perspectives of microplastic research in Nigeria. *International Journal of Environmental Analytical Chemistry*, 3:4-10.
- Eze C. C., Anaebonam, E, Nweze, K., Onyemeka, R, Frank-Ogu, N, Justice-Alucho, C. & Chiroma, I. (2021). Impact of municipal solid waste on the water quality of Otamiri River in Owerri, South-Eastern Nigeria. *World Journal of Biology Pharmacy and Health Sciences*, 07 (03), 065–072
- Eze, V. C., Onwukeme, V. I., Ogbuagu, J. O., Aralu, C. C., Aduaka, C. N. & Okoro, N. J. (2024). Evaluation of selected physicochemical properties of River Otamiri, Imo State. *Health Environ*, 4(1):209-216.
- Fagorite, V. I., Ahirakwem, C. A., Okeke, O. C., & Onyekuru, S. O. (2019). Physico-Chemical Characteristics of Otamiri River and Its Sediments in Parts of Owerri. *Elixir Geology. Elixir International Journal*, 131: 53223-53229.
- Federal Environmental Protection Agency (FEPA) 2003. Guidelines and Standards for Environmental Pollution Control in Nigeria. 238 pp.
- Florian, P., Joris, T., Ian, A. & Michael, A. (2020). Transport and Burial of Microplastics in Deep-Marine Sediments by Turbidity Currents. *Environment Science Technologies*, 54, 4180–4189
- Gideon, A., Adewumi, Y., Kehinde, O., Michael, O., Temitope O., Oluwatoyin, K., Oluwatosin, S. & Ademola F (2024). Why Nigeria should ban single-use plastics: Excessive microplastic pollution of the water, sediments and fish species in Osun River, Nigeria. *Journal of Hazardous Materials Advances*, 13, 100409

- He, B., Smith, M., Egodawatta, P., Ayoko, G., Rintoul, L. & Goonetilleke, A. (2021). Dispersal and transport of microplastics in river sediments. *Environ. Pollut.* 279:116884.
- He, C., Han, Y., Zheng, Y., Yue, W., Li, L., Wu, M. & Li, Z. (2021). Influence of physicochemical parameters on microplastics distribution in urban rivers. *Science of the total environment*, 799: 149380.
- Hüffer, T., Hofmann, T., & Griebler, C. (2017). Sorption of organic compounds by microplastics in the aquatic environment: A review. *Water Research*, 125: 305–317.
- Hüffer, T., Hofmann, T., & von der Kammer, F. (2017). Sorption of Metals onto Microplastics in Freshwater. *Environmental Science & Technology*, 51(24): 13472–13480.
- Kerubo, O. J., Muthumbi, A. W., Onyari, M. J., Kimani, E. N. & Robertson-Andersson, D. (2020). Microplastic pollution in the surface waters of creeks along the Kenyan coast, Western Indian Ocean (WIO). *Western Indian Ocean Journal of Marine Science*, 19 (2) 2020 75-88.
- Lares, M., Ncibi, M. C., Sillanpää, M. & Sillanpää, M. (2018). Occurrence, identification and removal of microplastic particles and fibers in conventional activated sludge process and advanced MBR technology. *Water Resources*, 133:236–246
- Li, J., Liu, H., Paul Chen, J., Xie, Z., & Li, Y. (2019). Sources and fate of microplastics in marine and freshwater environments: A review. *Water Research*, 162:191–206.
- Li, Y., Sun, N., Li, J., Tang, R., Miu, Y. & Ma, X. (2021). Research on the Influence of Microplastics on Marine Life. IOP Conference Series: *Earth and Environmental Science*, 6 (31):012006
- Liang, H., de Haan, P., Cerdà-Domènech, M., M'endez, J., Lucena, F., García-Aljaro, C., Sanchez-Vidal, A. & Ballester, E. (2023). Detection of faecal bacteria and antibiotic resistance genes in biofilms attached to plastics from human-impacted coastal areas. *Environmental Pollution* 319:120983
- Luís, L. G., Ferreira, P., Fonte, E., Torres, T., & Soares, A. M. V. M. (2020). Microplastics increase mercury bioconcentration in gills and bioaccumulation in the liver, and cause oxidative stress and damage in *Dicentrarchus labrax* juveniles. *Environmental Pollution*, 266(3): 115286

- Lusher, A. L., McHugh, M. & Thompson, R. C. (2013) Occurrence of microplastics in the gastrointestinal tract of pelagic and demersal fish from the English Channel. *Marine Pollution Bulletin*, 67, 94–99.
- Miller, E., Sedlak, M., Lin, D., Box, C., Holleman, C., Rochman, C., & Sutton, R. (2021). Recommended best practices for collecting, analyzing, and reporting microplastics in environmental media: Lessons learned from comprehensive monitoring of San Francisco Bay. *Journal of Hazardous Materials*, 409 (2021) 124770
- Moore, C. J. (2008). Synthetic polymers in the marine environment: A rapidly increasing, long-term threat. *Environmental Research*, 108(2): 131–139.
- Nizzetto, L., Futter, M. N., & Langaas, S. (2016). Are agricultural soils dumps for microplastics of urban origin? *Environmental Science & Technology*, 50(20): 10777–10779.
- Nwakanma, C., Blessed, T. & Hanson, H. (2021). Assessment of Microplastics Physical and Chemical Property Concentration in Surface Stream Water of Uyo Metropolis. *African Journal of Environment and Natural Science Research*, 4(1):62-83. ISSN: 2689-9434
- Nwonumara, G. N., Okoro, P. O. & Okogwu, O. I. (2021). Assessment of the incidence of microplastics at Ndibe, Cross River, Nigeria. *The Zoologist*, 19:46-51.
- Nworie, F. S. (2021). Microplastics: Holistic overview of source, identification, interaction, health and environmental implications and strategies of abatement. *Acta Chemica Malaysia (ACMY)*, 5(1): 18-23
- Ogbomida T., Obazele O., Aganmwonyi, I., Chukwuka, O., Emeribe, C. & Omoigberale, M. (2023). Assessment of Microplastics in Water, Sediment, and Fish of Ikpoba Rivers of Edo State, Nigeria. *African Scientist*, 24, 3:366-376
- Okechi, R. N., & Chukwura, E. I. (2020). Physicochemical and Bacteriological Qualities of Otamiri River Water and Sediment in South Eastern Nigeria. *Frontiers in Environmental Microbiology*, 6 (2): 18-26.
- Okeke, P. N. & Adinna, E. N. (2013). Water Quality Study of Otamiri Otamiri-oché River in Owerri, Nigeria. *Universal Journal of Environmental Research and Technology*, 3(6): 641-649.

- Olarinmoye, M., Stock, C., Scherf, N., Whenu, O., Asenime, C. & Ganzallo, S. (2021). Microplastic Presence in Sediment and Water of a Lagoon Bordering the Urban Agglomeration of Lagos, Southwest Nigeria. *Geosciences*, 10, 494;
- Pittura, L., Avio, C. G., Giuliani, M. E., d'Errico, G., Keiter, S. H., Cormier, B., ... Regoli, F. (2020). Microplastics as Vehicles of Environmental PAHs to Marine Organisms: Combined Chemical and Physical Hazards to the Mediterranean Mussels, *Mytilus galloprovincialis*. *Frontiers in Marine Science*, 7:1–13.
- Shen, Z., Jia, L., & Wen, Q. (2020). Nitrate Pollution in Groundwater and Its Health Effects on the Residents in Coastal Areas of Northern Jiangsu Province, China. *Archives of Environmental & Occupational Health*, 75(5):311–319.
- Song, Y., Li, J., Hu, J., Hu, T., Wu, Z., & Cheng, Y. (2021). Heavy metal(loid)s in microplastics from the coastal environments in China. *Environmental Pollution*, 277: 116764.
- Su, L., Xue, Y., Li, L., Yang, D., Kolandhasamy, P., Li, D., & Shi, H. (2019). Microplastics in Taihu Lake, China. *Environmental Pollution*, 252(2): 1882–1891.
- Sun, Q., Wang, X., Han, L., Jin, X., Yu, Z., Zhao, J., ... Shi, D. (2021). Dynamics of sulfur transformation and its driving factors in the sediment of eutrophic and hypertrophic lakes. *Chemosphere*, 263: 128136
- Tian, H., Jiang, L., Liu, L., Chen, M., Shen, X., & Zhu, Y. (2021). Enhanced bioaccumulation of microplastics by bacterial diversity changes under high biochemical oxygen demand condition. *Science of the Total Environment*, 773: 144970.
- Ting, P., Haiqing, L., Fang, Y., Fuhong, S., Youjun, G., Hao, Y., Dongxia, F., Xingxuan, Z. & Qianqian, W. (2023). Review of microplastics in lakes: sources, distribution characteristics, and environmental effects. *Carbon Research*, 2:25
- Toumi, H., Abidli, S. & Bejaoui, M. (2019). Microplastics in freshwater environment: the first evaluation in sediments from seven water streams surrounding the lagoon of Bizerte (Northern Tunisia). *Environ. Sci. Pollut. Res.* 26 (14), 14673–14682.
- Ubaka, K. G., Ekwonu, A. M. & Ogah, J. O. (2019). Effect of pollution on the physicochemical characteristics of Otammiri River, Imo State. *American Academic & Scholarly Research Journal*, 11(4)

- United States Environmental Protection Agency, US EPA (2021). A community science for sampling microplastic pollution. *EPA's Microplastic Beach Protocol*, 1-22
- Van Franeker, J. A. & Law, K. L. (2015). Seabirds, gyres and global trends in plastic pollution. *Environmental Pollution*, 203, 89–96.
- Wang, Z., Xie, X., Zhao, X., & Liu, Y. (2018). Seasonal variations in denitrification rates and potential controlling factors in a subtropical estuary (Min River Estuary, Southeast China). *Environmental Science and Pollution Research International*, 25(4): 3734–3743.
- Williams, J. N. & Osahon, N. T. (2021). Assessment of Microplastic Degrading Potential of Fungal Isolates from an Estuary in Rivers State, Nigeria. *South Asian Journal of Research in Microbiology*, 9(2): 11-19.
- Wright, R. J., Erni-Cassola, G., Zadjelovic, V., & Gibson, M. I. (2020). Christie-Oleza, J. A. Accelerating rates of plastic pollution in terrestrial ecosystems. *Nature Communications*, 11(1): 3487.
- Xue, X., Li, J., Wu, C., Liu, F., & Jia, F. (2020). Interactions of Pb(II) and Cd(II) with microplastics: Effect of pH, concentration and adsorption time. *Journal of Hazardous Materials*, 384: 121324.
- Yalwaji, B., John-Nwagwu, H. O. & Sogbanmu, T. O. (2022). Plastic pollution in the environment in Nigeria: a rapid systematic review of the sources, distribution, research gaps and policy needs. *Sci. Afr.* 16, e01220.
- Zettler, E. R., Mincer, T. J., & Amaral-Zettler, L. A. (2013). Life in the "Plastisphere": Microbial Communities on Plastic Marine Debris. *Environmental Science & Technology*, 47(13): 7137–7146.
- Zhang, D., Zhang, Z., Qian, Y., & Jiang, S. (2019). Contribution of bacteria to the bioaccumulation of microplastics in aquatic organisms. *Science of the Total Environment*, 693: 133499.
- Ziajahromi, S., Kumar, A. & Neale, P. (2020). Correlation of microplastics abundance and physicochemical properties of urban water bodies and its implications for stormwater management. *Journal of Hazardous Materials*, 385:121582