

1  
2 **EXPLORING IMPACT OF TELEREHABILITATION IN**  
3 **POST VIRAL FATIGUE SYNDROME FOLLOWING**  
4 **COVID-19: A NARRATIVE REVIEW OF LITERATURE**

5  
6  
7  
8  
9 **ABSTRACT**

10  
11  
**Aims:** The purpose of this study was to see how telerehabilitation affected post-viral tiredness following covid19.

**Study design:** Narrative Review.

**Materials and methods:**

Various computerised data bases were used to conduct a structured Literature search. PubMed, Google Scholar, Pedro, American Physical Therapy Association (APTA), World Health Organization (WHO), World Confederation Physical Therapy (WCPT), Medscape, Research-gate Embase, American Physical Therapy Association (APTA), World Health Organization (WHO), World Confederation Physical Therapy (WCPT), Medscape, Research-gate Embase Clinical trials, randomised controlled trials, cross-sectional studies, cohort studies, and literature reviews were all part of the search strategy. The search was restricted to English literature alone.

**Result:**

A total of 75 items were found in various electronic databases. Based on inclusion and exclusion criteria, 57 articles containing actual substance were shortlisted. Telerehabilitation in post-viral fatigue syndrome following covid 19 had a good effect on the patients, according to the study. It helps individuals with Post-Viral Fatigue Syndrome increase their breathing rate, physical strength, and overall function.

**Conclusion:**

This narrative review of the literature discovered that physiotherapy therapies, particularly Telerehabilitation, have a good effect in post-viral fatigue following covid19. Telerehabilitation has a greater impact on reducing the symptoms of post viral fatigue syndrome patients and will aid in their recovery. The things taught in Telerehabilitation also assist a lot, and the exercise improves the cardiovascular system by increasing HR (heart rate), breathing rate, body temperature, and blood pressure (Blood pressure).

12 *Keywords:* Telerehabilitation , covid19 , respiratory diseases, Post viral fatigue syndrome

## 14 **Introduction**

15 Telerehabilitation is defined as the use of technology to assist people with disabilities "Data and communication  
16 technologies are used to deliver recovery and rehabilitation services. Assessment, evaluation, checking,  
17 anticipating, intervening, oversight, instruction, meetings, and coaching are all examples of telerehabilitation  
18 administrations" <sup>[1]</sup> Telerehabilitation is a new and innovative methodology that can provide real assistance  
19 during the home restoration procedure for the improvement of engine, cognitive, or mental disorders. It can  
20 provide services such as physiotherapy, language training, word-related treatment, patient telemonitoring, and  
21 teleconsultation to help patients who are confined to their homes and do not have access to specialists or other  
22 medical professionals <sup>[2]</sup>. There are a couple of studies that have involved telerehabilitation in patients with  
23 COVID-19. This mediation has been demonstrated to be viable in respiratory, cardiovascular, metabolic, and  
24 neurological pathologies <sup>[3]</sup>post-viral condition, otherwise called post-viral weariness, is a sensation of fatigue  
25 and shortcoming that continues after an individual has recuperated from a viral sickness. It can grow even after  
26 a typical disease like the flu <sup>[4]</sup>.

27 For a long time, post-viral fatigue has been recognised and investigated. Previous research on post-viral  
28 tiredness is reviewed in this article <sup>[6]</sup>. "Upper respiratory tract infections frequently result in short-term fatigue,  
29 emphasising the importance of considering time periods after the person has recovered from the acute  
30 symptoms" <sup>[6,8]</sup>. "More serious infections can cause long-term exhaustion, and human herpesviruses and  
31 enteroviruses have been linked to chronic fatigue syndrome. It's not surprising, then, that a severe disease like  
32 COVID-19 can result in long-term symptoms. Long Covid is a condition that has only recently been identified,  
33 and the symptoms are discussed here. Researchers are currently in the process of identifying the condition" <sup>[6,7]</sup>.

34 "Post-viral weariness condition (PVFS) is an ailment that is sorted among different problems of the sensory  
35 system in the 11th update of the International Classification of Diseases. As per the current codification  
36 framework distributed by the World Health Organization in 2019, PVFS covers constant exhaustion disorder  
37 (CFS) and harmless myalgic encephalomyelitis (ME), baffling conditions recently assigned as individual  
38 substances, and presently arranged under the PVFS umbrella. Albeit a few CFS/ME cases are not gone before  
39 by a viral contamination. <sup>[5]</sup>. Post-viral exhaustion is incessant and normal, and can be noticed after  
40 contamination with various infections. By and large, the greatness of exhaustion is connected with the  
41 seriousness of the disease, and may to a limited extent mirror the recuperation saw in any extreme ailment and it  
42 is to depict the relentless impacts seen after COVID-19 contamination, and the condition currently alluded to as  
43 Long Covid" <sup>[6]</sup>. "The COVID-19 pandemic, which was caused by the Severe Acute Respiratory Syndrome  
44 Coronavirus 2 (SARS-CoV-2), has had a devastating effect on the global population, with a high fatality rate.  
45 The current study examines the literature on the short- and long-term health effects of previous epidemics and  
46 illnesses in order to determine whether there are any health risks related with COVID-19 recovery. Such

47 problems, according to previous research on post-epidemic and post-infection recovery, include the  
48 development of significant weariness. Certain factors, such as the severity of infection, in addition to the  
49 ‘cytokine storm’ experienced by many COVID-19 patients, may contribute to the development of later health  
50 problems”<sup>[8]</sup>.

51 Rehabilitation professionals plays an important role in speeding up the recovery of post viral fatigue syndrome  
52 patients<sup>7</sup>.

53  
54  
55  
56  
57  
58

**Table 1. Review of literature**

AUTHOR & YEAR OF PUBLICATION	TOPIC	METHOD	RESULT
<p>Marcelo Dalbosco-Salas et.al.,Rodrigo Torres-Castro et.al.,Andres Rojas Leyton et.al.,2020</p>	<p>Effectiveness of a Primary Care Telerehabilitation Program for Post-COVID-19 Patients: A Feasibility Study</p>	<p>In Chile, an observational, prospective investigation was carried out in seven primary care centres. Adult patients (&gt;18 years) who had previously been infected with SARSCoV-2 were included in the study. The telerehabilitation program included 24 supervised home-based exercise training sessions. Before and after the intervention, the efficacy was assessed using the 1-minute sit-to-stand test (1-min STST), the 36-Item Short Form Health Survey (SF-36), weariness, and dyspnea symptoms. A total of 115 patients (55.4 percent female) with an average age of 55.6 12.7 years were included in the study. Fifty-seven patients (50%) had previous hospitalizations, with 35 (30.4%) being admitted to the ICU [11].</p>	<p>After the intervention, the 1-min STST improved from 20.5 10.2 (53.1 25.0 percent expected) to 29.4 11.9 (78.2 28.0 percent projected) repeats (p 0.001). From 39.6 17.6 to 58.9 20.5, the SF-36 global score improved dramatically. After the intervention, fatigue and dyspnea were greatly reduced. Despite the lack of a control group, this study found that a telerehabilitation programme used in primary health care is practical and helpful in increasing physical capacity, quality of life, and symptoms in adult COVID-19 survivors [9].</p>
<p>Jianan Li et.al., Wenguang Xia et.al., Chao Zhan et.al., 2022</p>	<p>A telerehabilitation programme in post-discharge COVID-19 patients (TERECO): a randomised controlled trial</p>	<p>Randomized controlled trial with 1:1 block randomization in parallel groups.</p> <p>Setting China's Jiangsu and Hubei provinces have three large hospitals.</p> <p>Participants were 120 COVID-19 survivors who had previously been hospitalised and still had dyspnoea problems, with 61 being assigned to the control group and 59 to the TERECO group.</p> <p>Intervention Unsupervised 6-week home-based workout program that includes breathing control and</p>	<p>At post-treatment, the adjusted between-group difference in 6MWD was 65.45 m (95 percent CI 43.8 to 87.1; p0.001), and at follow-up, it was 68.62 m (95 percent CI 46.39 to 90.85; p0.001). The post-treatment effect for LMS was 20.12 s (95 percent CI 12.34 to 27.9; p0.001), and the follow-up effect was 22.23 s (95 percent CI 14.24 to 30.21; p0.001). Except for post-treatment maximum voluntary breathing, there were no variations in lung function between the groups. The TERECO group had a higher increase in the SF-12 physical component, with treatment effects of 3.79 (95 percent CI 1.24 to 6.35; p=0.004) at post-</p>

		<p>thoracic expansion, aerobic exercise, and LMS exercise given through smartphone and remotely monitored with heart rate monitoring.</p> <p>Outcomes The 6 minute walking distance (6MWD) in meters was the primary outcome. Squat time in seconds, pulmonary function as determined by spirometry, HRQOL as defined by the Short Form Health Survey-12 (SF-12), and mMRC-dyspnoea were all secondary outcomes. At 6 weeks (post-treatment) and 28 weeks, the outcomes were evaluated (follow-up) <sup>[12]</sup>.</p>	<p>treatment and 2.69 (95 percent CI 0.06 to 5.32; p=0.045) at follow-up<sup>[10]</sup>.</p>
<p>Abubeker Alebachew et.al., Ahmed Adem Mohammed et.al., 2020</p>	<p>Effectiveness and feasibility of telerehabilitation in patients with COVID-19: a protocol for a systematic review and meta-analysis</p>	<p>From the beginning through the end of November 2021, the databases PubMed, Web of Science, Science Direct, Physiotherapy Evidence Database, Google Scholar, and the Cochrane Library will be searched. There will be randomised controlled trials looking into the effectiveness of telerehabilitation in the treatment of COVID-19. Functional ability, cardiopulmonary exercise testing, and quality of life will be the major outcomes. Anxiety/depression levels, sleep quality, mortality rate, completion rate, cause for withdrawal, adverse events, service satisfaction, cost-effectiveness, and other potential factors will all be secondary outcomes. Two reviewers will independently screen and collect data from included research, as well as assess their quality. To assess the risk of bias, the Cochrane risk of bias instrument will be utilised. For</p>	<p>The I test will be used to analyse the heterogeneity of the included studies. The data will be analysed using a fixed-effects model first. It is assumed that there is significant heterogeneity among the included trials if <math>I &gt; 0.5</math> or <math>p &lt; 0.1</math>, and a random-effects model will be utilised in this situation. Sensitivity analysis will be used to establish the source of heterogeneity by removing trials one by one <sup>[11]</sup>.</p>

		statistical analysis, the Review Manager V.5.4 (Cochrane Collaboration) programme will be used <sup>[13]</sup> .	
Shrushti Prashant Jachak et.al., Pratik Arun et.al., 2021	Great Awakening – Telerehabilitation in Physiotherapy during Pandemic and Impact of COVID-19	<p>The number of diseases and deaths continues to climb, and the impact of the crisis is expanding across emerging countries. Reactions, attitudes, and outcomes differed all throughout the world. The outbreak has had negative mental health consequences and symptoms. This pandemic has had an impact on the healthcare system, which treats patients with diseases other than corona. Despite the fact that the power and severity of the slowdown varies from transitory to long-term recession, they all agree that the slowdown will have a significant influence on various sectors of the economy. Most importantly, panic among consumers and businesses has distorted regular purchase patterns and resulted in market inconsistency.</p> <p>In this difficult period, digitalization and automation have shown to be the answer intense impact amongst various sectors of the economy. Most importantly, some panic among consumers and firms has disfigured normal patterns of consumption and caused market inconsistency. Digitalization and automation have proved to be the solution in this challenging phase<sup>[14]</sup>.</p>	<p>As the world scrambles to alleviate the virus's massive healthcare burden, most economies are bracing for the damage the infection will wreak. Many governments have already proposed a series of 'economic packages' to aid struggling businesses, workers, and healthcare systems. The Indian government and the Reserve Bank of India have also taken a number of steps to combat the corona virus sickness and improve the country's economic situation. As a result of the social distancing used to break the chain of infection, healthcare is also impacted, with patients unable to visit clinics and hospitals. In this difficult period, digitalization and automation have shown to be the answer. With the help of numerous smartphone applications and websites for online consultation, telemedicine and telephysiotherapy have revolutionised the way people spend their time in a good way. Whether it's a live session or a recorded one, Psychological hazards are reduced in individual or group sessions. It is beneficial to the patients as well as the motivation of the staff. As a result, telemedicine and tele physiotherapy are becoming more popular the amount of care provided to those suffering from both mental and physical illnesses health and mental well-being, reduced hospital costs bolster traditional face-to-face practise and increase adherence to therapy and happiness with it<sup>[12]</sup></p>

<p>Cleofas Rodriguez- Blanco et.al.,</p> <p>Juan Jose Gonzalez et.al.,</p> <p>2021</p>	<p>Short-Term Effects of a Conditioning Telerehabilitation Program in Confined Patients Affected by COVID-19 in the Acute Phase. A Pilot Randomized Controlled Trial</p>	<p>A total of 40 volunteers were randomly assigned to one of two groups: an experimental group that underwent muscle training and a control group that did not engage in any physical activity. The one-week intervention was completed by 36 individuals, 18 in each group. The six-minute walking test, multidimensional dyspnoea-12, thirty-second sit-to-stand test, and Borg Scale were all used to collect data<sup>[15]</sup>.</p>	<p>At the start, both groups were similar. The experimental group showed statistically significant improvement (<math>p &lt; 0.05</math>) over the control group. There were no gender differences found (<math>p &gt; 0.05</math>).<sup>[13]</sup>.</p>
<p>Juan Jose Gonzalez et.al.,</p> <p>Martin Botella- Rico et.al.,</p> <p>2020</p>	<p>Therapeutic pulmonary telerehabilitation protocol for patients affected by COVID-19, confined to their homes: study protocol for a randomized controlled trial</p>	<p>In the regions of Andalusia, Murcia, and Valencia, patients will be recruited (Spain). Patients will be confined to their homes, where they will follow their prescribed exercise programme, which will be monitored remotely. Evaluators will be on hand to conduct telematically controlled measurements at the start, middle, and end of the study. The patients will be randomly assigned to one of three groups: two will participate in a home exercise programme (breathing exercises or non-specific muscle toning exercises), and the third will engage in sedentary activities while using mental activation techniques as a sham group. Physical tests, effort, and perceived weariness will be used to assess respiratory variables and other physical state factors<sup>[16]</sup>.</p>	<p>Researchers look at how two types of exercise regimens, toning and respiratory, influenced patients with COVID-19 during their home confinement phase in terms of fatigue and perceived exertion, physical health, and respiratory activity maintenance. Researchers want to look into a field that has never been investigated before, such as the effects of a toning and respiratory exercise programme in these patients, in historical circumstances that no one has ever seen before in Spain, because the general population has never been forced to remain confined in their homes due to a coronavirus pandemic (COVID-19)<sup>[14]</sup>.</p>
<p>Kelsey Rosen et.al.,</p> <p>Monika Patel et.al.,</p> <p>2020</p>	<p>Delivering Telerehabilitation to COVID-19 Inpatients: A Retrospective Chart Review Suggests it is a Viable Option</p>	<p>A study was performed a retrospective chart assessment of COVID-19 patients who underwent either telerehabilitation alone or a mix of telerehabilitation and in-person rehabilitation. COVID-19</p>	<p>Twelve of the 33 COVID-19 inpatients who underwent telerehabilitation, in-person rehabilitation, or both received solely telerehabilitation (age range, 33 to 65 years; all but one male). They were discharged home after</p>

		<p>inpatients were chosen for telerehabilitation based on the algorithm if they were able to ambulate freely, use technology, had stable vital signs, require minimum supplementary oxygen, and are cognitively intact. Data from inpatients who solely received telerehabilitation, which comprised patient education, therapeutic activities, and breathing techniques, was evaluated<sup>[20]</sup>.</p>	<p>demonstrating independence with their tailored home exercise programmes in one to two sessions, without the need for an in-person rehabilitation consultation, without the need for extra oxygen, and with no exacerbation of symptoms<sup>[15]</sup>.</p>
<p>Jose- Manuel Pastora- Bernol et.al., Marin Jose et.al., 2021</p>	<p>Telerehabilitation Intervention in Patients with COVID-19 after Hospital Discharge to Improve Functional Capacity and Quality of Life. Study Protocol for a Multicenter Randomized Clinical Trial</p>	<p>This study uses a single blind multicenter randomised clinical trial in patients with a COVID-19 diagnosis who have been released from the hospital in Andalusia (south of Spain). This study will follow the SQUIRE (Standards for Quality Improvement and Excellence in Reporting) recommendations and will follow the CONSORT (Consolidated Standards of Reporting Trials) criteria<sup>[21]</sup>.</p>	<p>This study should give information on the feasibility of implementing TR programs in COVID-19 patients after release from the hospital, as well as health resources and costs, which will allow the development of new intervention policies for this group of patients<sup>[16]</sup>.</p>
<p>Snehil Dixit et.al., Audrey Borghi-Silva2 et.al., Kalyana Chakravarthy Bairapareddy et.al., 2021</p>	<p>Revisiting pulmonary rehabilitation during COVID-19 pandemic: a narrative review</p>	<p>During the coronavirus pandemic of 2019, there was an apparent link between the chances of problems associated with SARSCoV-2 infection and people with a history of chronic respiratory disease (COVID-19). In cardiopulmonary management, SARS-CoV-2 poses a serious risk. Furthermore, in the event of a pandemic, chronic respiratory disorders may increase the risk of morbidity and mortality among the affected population<sup>[48]</sup>.</p>	<p>The purpose of this review is to highlight the importance of pulmonary rehabilitation (PR) in people with chronic respiratory disorders (COPD and Asthma) in the COVID-19 era. Following initial medical therapy and acute recovery, exercise-based pulmonary rehabilitation (PR) may play a critical role in long-term management and recovery in the population with a pre-existing pulmonary diagnosis who have contracted SARS-CoV-2. To combat post-COVID-19 tiredness, energy saving measures will be useful in PR of low to moderate intensity instances. Furthermore, there is an immediate need to manage post-</p>

			COVID-19 anxiety and sadness, which is impacting the PR delivery system <sup>[49]</sup> .
Abayomi Salawu et.al., Angela Green et.al., Nina brixey et.al., 2020	A Proposal for Multidisciplinary Tele-Rehabilitation in the Assessment and Rehabilitation of COVID-19 Survivors	In February 2020, a global pandemic of COVID-19, a new extremely contagious disease caused by coronavirus (severe acute respiratory syndrome (SARS)-Cov-2) infection, was declared. Although it is usually transmitted through the respiratory system, it can damage other organ systems in the body. In severe circumstances, 20% of individuals affected require hospitalisation and mechanical ventilation. About half of those who survive the disease have lasting functional abnormalities that require multimodal therapy. Existing community services are unable to provide the manpower needed to give the essential rehabilitation input. To monitor these patients, strict medical follow-up rules need scheduled assessments within 12 weeks of release. Existing treatment pathways are unlikely to be able to meet demand because to the short timeframe in which these occurrences occur The following is a proposal for an innovative integrated post-discharge care pathway to ease follow-up by acute medical teams (respiratory and intensive care) as well as a specialty multidisciplinary rehabilitation team <sup>[49]</sup> .	The monitoring and provision of complete medical assessments and multidisciplinary rehabilitation will be possible with such a pathway. This research recommends that a tele-rehabilitation model be added into the pathway, utilising digital communication technologies to provide these patients with speedy remote assessment and efficient therapy delivery. Following hospital discharge, tele-rehabilitation provides a speedy and effective way to respond to the specialised rehabilitation needs of COVID-19 survivors <sup>[17]</sup> .
Pamela Tanguay et.al., Nicole Marquis et.al., Isabelle Gaboury et.al.,	Telerehabilitation for Post-Hospitalized COVID-19 Patients: A Proof-of-Concept Study During a Pandemic	A pre-experimental, pre-post pilot research was done. COVID-19 patients who had been discharged from the hospital were recruited. Eight weeks of supervised	We booked 64 supervised sessions with seven individuals who had only minor technical difficulties. Initial results revealed that pulmonary symptoms had a moderate to

2021		physiotherapy sessions were part of the intervention. We kept track of technical concerns, recruitment strategy success, and participant participation during supervised sessions. The effect of pulmonary symptoms on quality of life and functional health was studied <sup>[50]</sup> .	significant influence on quality of life. On the EuroQol-Visual Analog Scale (EQ-VAS) instrument, all patients had improved from 10 to 45 points after eight weeks, suggesting clinical significance <sup>[18]</sup> .
Siddhi Ghodge et.al., Prajakta Tilaye et.al., Swati Nerkar et.al., 2020	Effect of Pulmonary Telerehabilitation on Functional Capacity in COVID Survivors; An Initial Evidence	This study included 10 participants (8 males, 2 females) who met the inclusion criteria of post-covid-19 survivors with respiratory system involvement who underwent at least 6 weeks of pulmonary telerehabilitation at least 3 days per week. Breathing exercises, an incentive Spirometer, a variety of aerobic and strengthening exercises, and patient education were used in pulmonary telerehabilitation. The visual analogue scale (VAS) for tiredness, SPO2 using a pulse oximeter, single breath count, one minute sit to stand test, and one minute squat test were measured before and after 6 weeks of rehabilitation <sup>[51]</sup> .	The descriptive statistics had been completed. According to the normality analysis, non parametric data was subjected to the Wilcoxon sign rank test, while parametric data was subjected to the Paired t test. A p value of less than 0.05 was deemed significant.  The one minute sit to stand test (p=0.005), one minute squat test (p=0.007), resting SPO2 (p=0.025), post exercise SPO2 (p=0.003), and post exercise VAS for tiredness (p=0.017) all indicated statistically significant differences between pre and post rehabilitation. In the single breath count test (p=0.415), there was no statistically significant difference. Resting, post-exercise SPO2, and the VAS for exhaustion all demonstrated clinically meaningful differences with substantial effect sizes <sup>[19]</sup> .

59  
60

61 **Methodology**

62 **3.1 PURPOSE**

63 The purpose of this study was to determine if telerehabilitation has an effect on post-viral fatigue following  
64 COVID-19.

65 **3.2 OBJECTIVE**

66 The goal of this study is to evaluate the impact of telerehabilitation on post-viral fatigue following COVID-  
67 19.

68 **3.3 STUDY DESIGN:** Narrative review of literature.

70 **3.4 MATERIALS AND METHODS:**

71 Various computerised data bases were used to conduct a structured Literature search.

73 **3.5 SOURCES OF DATA:** PubMed, Google Scholar, Pedro, American Physical Therapy Association (APTA),

74 World Health Organization (WHO), World Confederation Physical Therapy (WCPT), Medscape, Research-gate


75 Embase, American Physical Therapy Association (APTA), World Health Organization (WHO), World

76 Confederation Physical Therapy (WCPT), Medscape, Research-gate Embase Clinical trials, randomised

77 controlled trials, cross-sectional studies, cohort studies, and literature reviews were all part of the search

78 method. Only English literature was included in the search.

81 **FIG 1. PRISMA CHART**



109

110

111

112

### 3. RESULTS AND DISCUSSION

According to the findings of this study, telerehabilitation provides various benefits in postviral fatigue syndrome patients, including decreasing symptoms and assisting recovery. Exercise improves the cardiovascular system by increasing HR (heart rate), breathing rate, body temperature, and blood pressure, hence telerehabilitation lessons are also beneficial (Blood pressure).

Shrushti Prashant Jachak et.al., Pratik Arun et.al., in 2020 stated that Telemedicine and Telephysiotherapy for online consultation have positively impacted time. "Patients' psychological risks are reduced when live sessions are conducted, and it is also advantageous to the therapy's motivation. Tele-rehabilitation improves the quality of treatment for persons with physical and mental health problems, lowers hospital costs, and strengthens traditional face-to-face therapy"<sup>[12]</sup>.

In 2021 Cleofas Rodriguez- Blanco et.al., Juan Jose Gonzalez et.al., did a study and stated that In COVID-19 individuals with mild to moderate symptomatology in the acute stage, a one-week telerehabilitation programme based on muscle toning exercise is beneficial, safe, and feasible<sup>[13]</sup>.

Juan Jose Gonzalez et.al., Martin Botella- Rico et.al., in 2020 did a study and stated that this article provides a detailed description of a randomised controlled trial that looked at the effects of two types of exercise programmes, toning and respiratory, on fatigue and perceived exertion, physical health, and respiratory activity maintenance in patients with COVID-19 who were confined to their homes<sup>[16]</sup>.

In 20 Kelsey Rosen et.al., Monika Patel et.al., did a study and stated that Inpatient telerehabilitation appears to be a viable alternative for COVID-19-positive hospitalised patients, and it could be a safe way to give inpatient rehabilitation to isolated or at-risk groups. Inpatient telerehabilitation minimised staff exposure while offering critical teaching and services to patients at our hospital. "No research has looked into the use of telerehabilitation for hospitalised patients, including those with COVID-19, to our knowledge. Our findings indicate that this novel strategy merits further investigation"<sup>[15]</sup>.

In 2021 Jose- Manuel Pastora- Bernol et.al., Marin Jose et.al., did a study and concluded that the viability of executing TR programmes in COVID-19 patients after discharge from the hospital, as well as health resources and costs, in order to design new intervention policies for this group of patients<sup>[16]</sup>

"The current concerns about COVID-19's consequences should not deter us from taking action; rather, they should encourage us to look for alternatives that will bring pulmonary rehabilitation closer to patients who require it after being discharged from the hospital"<sup>[20]</sup>. Telerehabilitation is viewed in

143 this light as a method that could give an efficient and cost-effective treatment to the health issues  
144 highlighted. “Physiotherapy will be a cornerstone in the interdisciplinary health team for the recovery  
145 of the sequelae, which should lead to an improvement in the quality of life of COVID-19 patients, not  
146 only in intensive care units and in hospitalised patients, but also in the interdisciplinary health team for  
147 the recovery of the sequelae, which should lead to an improvement in the quality of life of patients  
148 with COVID-19” [20,21].

149 “Unlike other studies that require software to be installed on specific devices, our intervention has  
150 minimal drawbacks because it can be accessed from any location and on any device that has an  
151 internet connection, which most patients have (personal computer, laptop, tablet, smartphone)” [33,34].  
152 “Other investigations, by contrast, necessitate a highly complex technological platform, software  
153 installation, and multidirectional cameras for controlled clinical control between the healthcare  
154 professional and the patient” [22]. The speedy installation process is ensured because the patient  
155 employs his own technical devices.

156 In 2020 Siddhi Ghodge et.al., Prajakta Tilaye et.al., Swati Nerkar et.al., did a study and stated that  
157 Pulmonary telerehabilitation shows statistically and clinically significant improvement in functional  
158 capacity in COVID survivors [16].

159 However, this may result in a selection bias for patients who do not have access to technology. With  
160 the use of technology, technical issues (disconnection, device failures) and technological difficulties  
161 may arise [23].

#### 162 163 164 165 166 **4. CONCLUSION**

167 This narrative review of the literature discovered that physiotherapy interventions, notably  
168 Telerehabilitation, are effective in reducing post-viral tiredness after covid19 infection.  
169 Telerehabilitation has a higher influence on lowering symptoms and aiding recovery in individuals with  
170 postviral fatigue syndrome. According to the preliminary findings, pulmonary symptoms had a  
171 moderate to major impact on quality of life. After eight weeks, all patients had improved from 10 to 45  
172 points on the EuroQol-Visual Analog Scale (EQ-VAS), indicating clinical relevance.

173 Telerehabilitation lessons are also very beneficial, because exercise improves the cardiovascular  
174 system by boosting HR (heart rate), breathing rate, body temperature, and blood pressure (Blood  
175 pressure). As a result, telemedicine and telephysiotherapy are becoming more popular, increasing

176 the amount of care provided to those suffering from both mental and physical illnesses, lowering  
177 hospital costs, bolstering traditional face-to-face practise, and increasing adherence to therapy and  
178 satisfaction with it.

## 179 180 **ACKNOWLEDGEMENTS**

181 Acknowledged with LLMs  
182  
183  
184  
185  
186

## 187 188 **AUTHORS' CONTRIBUTIONS**

189 Authors may use the following wordings for this section: "NR and JY' designed the study, performed the statistical  
190 analysis, wrote the protocol, andwrote the first draft of the manuscript. 'Author BS' and 'Author FP' managed the analyses  
191 ofthe study. 'Author NR and BS' managed the literature searches..... All authors read and approved the final manuscript."  
192  
193  
194  
195  
196  
197  
198  
199  
200

## 201 202 **REFERENCES**

- 203  
204  
205 1. Rosen K, Patel M, Lawrence C, Mooney B. Delivering telerehabilitation to COVID-19 inpatients: a  
206 retrospective chart review suggests it is a viable option. HSS Journal®. 2020 Nov;16(1\_suppl):64-70.  
207
- 208 2. Maggio MG, De Luca R, Manuli A, Calabrò RS. The five 'W'of cognitive telerehabilitation in the Covid-  
209 19 era. Expert Review of Medical Devices. 2020 Jun 2;17(6):473-5.  
210
- 211 3. Dalbosco-Salas M, Torres-Castro R, Rojas Leyton A, Morales Zapata F, Henríquez Salazar E, Espinoza  
212 Bastías G, Beltrán Díaz ME, Tapia Allers K, Mornhinweg Fonseca D, Vilaró J. Effectiveness of a primary  
213 care telerehabilitation program for post-COVID-19 patients: a feasibility study. Journal of clinical  
214 medicine. 2021 Jan;10(19):4428.  
215
- 216 4. Brancatella A, Ricci D, Cappellani D, Viola N, Sgrò D, Santini F, Latrofa F. Is subacute thyroiditis an  
217 underestimated manifestation of SARS-CoV-2 infection? Insights from a case series. The Journal of  
218 Clinical Endocrinology & Metabolism. 2020 Oct;105(10):e3742-6.  
219

- 220 5. Ostojic SM. Diagnostic and pharmacological potency of creatine in post-viral fatigue syndrome. *Nutrients*.  
221 2021 Feb;13(2):503. Islam MF, Cotler J, Jason LA. Post-viral fatigue and COVID-19: lessons from past  
222 epidemics. *Fatigue: Biomedicine, Health & Behavior*. 2020 Apr 2;8(2):61-9.  
223
- 224 6. Simani L, Ramezani M, Darazam IA, Sagharichi M, Aalipour MA, Ghorbani F, Pakdaman H. Prevalence  
225 and correlates of chronic fatigue syndrome and post-traumatic stress disorder after the outbreak of the  
226 COVID-19. *Journal of Neurovirology*. 2021 Feb;27(1):154-9. Islam MF, Cotler J, Jason LA. Post-viral  
227 fatigue and COVID-19: lessons from past epidemics. *Fatigue: Biomedicine, Health & Behavior*. 2020 Apr  
228 2;8(2):61-9.  
229
- 230 7. Dalbosco-Salas M, Torres-Castro R, Rojas Leyton A, Morales Zapata F, Henríquez Salazar E, Espinoza  
231 Bastías G, Beltrán Díaz ME, Tapia Allers K, Mornhinweg Fonseca D, Vilaró J. Effectiveness of a primary  
232 care telerehabilitation program for post-COVID-19 patients: a feasibility study. *Journal of clinical  
233 medicine*. 2021 Jan;10(19):4428.  
234
- 235 8. Xia W, Zhan C, Liu S, Yin Z, Wang J, Chong Y, Zheng C, Fang X, Cheng W, Reinhardt JD. A  
236 telerehabilitation programme in post-discharge COVID-19 patients (TERECO): a randomised controlled  
237 trial. *Thorax*. 2021 Jul 26.  
238
- 239 9. Behan PO, Behan WM, Bell EJ. The postviral fatigue syndrome—an analysis of the findings in 50 cases.  
240 *Journal of Infection*. 1985 May 1;10(3):211-22.  
241
- 242 10. Seid AA, Aychiluhm SB, Mohammed AA. Effectiveness and feasibility of telerehabilitation in patients  
243 with COVID-19: a protocol for a systematic review and meta-analysis. *BMJ open*. 2022 Jan  
244 1;12(1):e058932.  
245
- 246 11. Jachak SP, Phansopkar PA, Naqvi WM, Kumar K. Great Awakening--Telerehabilitation in Physiotherapy  
247 during Pandemic and Impact of COVID-19. *Journal of Evolution of Medical and Dental Sciences*. 2020  
248 Nov 9;9(45):3387-94.  
249
- 250 12. Rodriguez-Blanco C, Gonzalez-Gerez JJ, Bernal-Utrera C, Anarte-Lazo E, Perez-Ale M, Saavedra-  
251 Hernandez M. Short-term effects of a conditioning telerehabilitation program in confined patients affected  
252 by COVID-19 in the acute phase. a pilot randomized controlled trial. *Medicina*. 2021 Jul;57(7):684.  
253

- 254 13. SPERANZA E, BOLZAN V, Giulio S, VITALI F. Impact of telerehabilitation after sports injuries on  
255 psychological outcomes during the COVID-19 pandemic. *Medicina dello Sport*. 2021 Dec;74(4):657-71.
- 256
- 257 14. Gonzalez-Gerez JJ, Bernal-Utrera C, Anarte-Lazo E, Garcia-Vidal JA, Botella-Rico JM, Rodriguez-  
258 Blanco C. Therapeutic pulmonary telerehabilitation protocol for patients affected by COVID-19, confined  
259 to their homes: study protocol for a randomized controlled trial. *Trials*. 2020 Dec;21(1):1-9.
- 260
- 261 15. Rosen K, Patel M, Lawrence C, Mooney B. Delivering telerehabilitation to COVID-19 inpatients: a  
262 retrospective chart review suggests it is a viable option. *HSS Journal®*. 2020 Nov;16(1\_suppl):64-70.
- 263
- 264 16. Pastora-Bernal JM, Estebanez-Pérez MJ, Molina-Torres G, García-López FJ, Sobrino-Sánchez R, Martín-  
265 Valero R. Telerehabilitation Intervention in Patients with COVID-19 after Hospital Discharge to Improve  
266 Functional Capacity and Quality of Life. Study Protocol for a Multicenter Randomized Clinical Trial.  
267 *International Journal of Environmental Research and Public Health*. 2021 Jan;18(6):2924.
- 268
- 269 17. Salawu A, Green A, Crooks MG, Brixey N, Ross DH, Sivan M. A proposal for multidisciplinary tele-  
270 rehabilitation in the assessment and rehabilitation of COVID-19 survivors. *International journal of*  
271 *environmental research and public health*. 2020 Jan;17(13):4890.
- 272 18. Tanguay P, Marquis N, Gaboury I, Kairy D, Touchette M, Tousignant M, Décary S. Telerehabilitation for  
273 post-hospitalized COVID-19 patients: a proof-of-concept study during a pandemic. *International Journal*  
274 *of Telerehabilitation*. 2021;13(1).
- 275
- 276 19. Ghodge S, Tilaye P, Deshpande S, Nerkar S, Kothary K, Manwadkar S. Effect of pulmonary  
277 telerehabilitation on functional capacity in COVID survivors; initial evidence. *Int J Health Sci Res*.  
278 2020;10(10):123-9.
- 279 20. Lista-Paz, A.; González-Doniz, L.; Souto-Camba, S. Qué papel desempeña la Fisioterapia en la pandemia  
280 mundial por COVID-19. *Fisioterapia* 2020, 42, 167–169.
- 281 21. Concepción Martín Cortijo, Francisco Javier Zarza Bejarano EGD, Marta Sánchez Cortés VTRFHU 12 de,  
282 (Madrid) O. Guía. De Fisioterapia en el Paciente con Sospecha De COVID-19 O COVID-19 Confirmado.
- 283 22. Guzik TJ, Mohiddin SA, Dimarco A, Patel V, Savvatis K, Marelli-Berg FM, Madhur MS, Tomaszewski  
284 M, Maffia P, D'acquisto F, Nicklin SA. COVID-19 and the cardiovascular system: implications for risk  
285 assessment, diagnosis, and treatment options. *Cardiovascular research*. 2020 Aug 1;116(10):1666-87.
- 286 23. Hwang R, Bruning J, Morris N, Mandrusiak A, Russell T. A systematic review of the effects of  
287 telerehabilitation in patients with cardiopulmonary diseases. *Journal of cardiopulmonary rehabilitation and*  
288 *prevention*. 2015 Nov 1;35(6):380-9.

