

Correlation and Path Coefficient analysis for selecting morphological and biochemical traits in Potato (*Solanum tuberosum* L.) in the 'Terai' region of Uttarakhand

Abstract

Knowledge regarding the direct and indirect effects of traits and understanding of traits' relationship with important morphological and biochemical attributes on yield amplification of breeding materials are vital prerequisites for crop improvement. The main focus of the research was to assess the correlation and path analysis in 32 CPRI advanced potato breeding lines using a randomized block design with three replications to direct proper selection criteria for tuber yield in the Terai region of Uttarakhand. The Central Potato Research Station has a collection of advanced potato breeding lines that have been selected and maintained for a long time. It has characterized its genetic makeup and usefulness for the breeding program. The results revealed that the average weight of tuber per plant, number of tubers per plant, tuber length, tuber girth, number of branches per plant and plant height contributed the highest positive correlation with tuber yield. The path coefficient result ~~showed~~revealed that the average weight of tuber per plant, number of tubers per plant and number of branches per plant had the highest positive direct effect on yield, whereas the effect of ~~others~~other traits ~~were~~was low. Therefore, these specific characters should be given more importance when selecting for yield performance.

Comment [MI1]: Can be removed

Keywords: Potato, Correlation analysis, Path coefficient analysis, Yield

Comment [MI2]: Consider different key words

INTRODUCTION

The potato (*Solanum tuberosum* L.) is a widely grown food crop, ranked third in human consumption after rice and wheat (FAOSTAT, 2021), and is grown worldwide as a primary crop,

a secondary crop, or an intercrop. The potato is an Andean native, and with a vegetative propagation strategy, it developed a short-day dependence on tuber development. Potato is a member of the Solanaceae family and is specifically a member of the *Petota* section of the *Solanum* genus (Spooner et al. 2014). The potato has a vast eco-geographical range and among all other major food crops in the world, it shows a unique nature in that it produces stolons or underground stems which in the presence of congenial environmental factors swell to produce tubers (Hijmans, 2001). The majority of potato cultivars are autotetraploid ($2n = 4x = 48$), highly heterozygous, suffer from acute inbreeding depression, and are vulnerable to a wide range of harmful diseases and pests. Autotetraploid cultivars that are propagated from seed tubers constitute the majority ~~in-of~~ commercial potato production. Potato is a wholesome food that provides proteins, carbohydrates, vitamin B complex, minerals, vitamin C, and high value dietary fibres and phenolic compounds (Woolfe, 1987). In 2019, China, India, and Russia were the three leading countries in potato production. China produced 91,818,950 tons of potatoes, accounting for 24.78% of the world's total potato production. Similarly, India produced 50,190,000 tons, which is 13.54% of the world's potato production, while Russia produced 22,436,581 tons, which is 5.95% of the world's potato production. Together, these three countries accounted for over 44% of the world's potato production. Potato production has seen a significant rise in the past 17 years, according to FAOSTAT (2019). The production has increased by approximately 96.54%, from 349,000 tons in 1993 to 863,348 tons in 2010. India has become the second-largest potato producer globally, with production of 51.3 million tons from a cultivated area of 2.16 million hectares with 23.77 tons per hectare productivity (FAOSTAT,2022). In India, potatoes are mainly used for table consumption, accounting for approximately 68% of the total usage. About 7.5% of potatoes are used for processing, 8.5% for

seed, and the remaining 16% of produce is wasted due to poor handling during pre- and post-harvest stages. Since the establishment of the Central Potato Research Institute in Shimla in 1949 has developed and released a total of 56 potato varieties that are suitable for various potato-growing regions in the country (Luthra *et al.* 2018). Tuber yield in potatoes is a complex polygenic characteristic (Killick, 1977) that is the result of interaction of diverse factors and has a low heritability. Consequently, more understanding of genetic variation among different potato attributes and how it affects yield should be useful to breeders for further crop improvement (Bhatt, 1973).

The correlation coefficient studies between various quantitative traits provide insight into the relationship between yield and its attributing characteristics. This information is utilized to select a superior plant type in the potato breeding program. Several researchers have reported correlations between morphological and biochemical features with potato yield (Mahgoub *et al.* 2015; Rajini, 2015). The estimate of the correlation coefficient only reveals the degree and nature of the association between yield and its constituents. However, it does not provide information about the direct and indirect effects of various yield variables on yield. The most important objective in potato breeding is to achieve high tuber yield while maintaining good quality. The yield of tubers is a complex trait that involves many interrelated components. Therefore, the path coefficient analysis is useful in showing the direct and indirect effects of casual variables on the response variable (Shubha and Singh, 2018). Researchers have widely utilized this method to evaluate the significance of yield components (Sidhu, 1992). The current analysis aims to examine the interrelationship between 32 different agro-morphological and biochemical variables and their effects on crop yield.

Material and methods

The present investigation ~~was~~ conducted ~~on~~ in the experimental field of the Vegetable Research Centre of the GBPUAT Pantnagar, District Udham Singh Nagar (Uttarakhand), during the rabi season of 2017-2018. Geographically, Pantnagar is situated in the Shivalik hills, at 29.5° N and 79.3° E, with an elevation of 244 m above mean sea level, which ~~are~~ is part of the 'Terai' mountain range of the outer Himalayas. The region experiences a humid and subtropical climate, with frost expected from the ~~final~~ last week of December to the end of January.

Experimental planting material and field trail:

The 32 diverse potato genotypes were used in the study along with three checks (Table 1), maintained in the Vegetable Research Centre of the University. The experiment was carried out in a randomized block design with three replications. In a plot 20 tubers with comparable size per genotype were planted with two rows spaced 60 cm between rows and 20 cm within rows. The fertilizer dose of (NPK ~~Kg per ha~~) was applied in a ratio of 160:100:120 Kg per ha in the form of Urea, Single super phosphate, and muriate of potash, respectively. Standard cultural practices were carried out in the experimental plot during the crop season.

Various growth parameters and tuber yield analysis :

The observations related to different growth parameters and yield-related traits were collected from five randomly ~~selected~~ chosen ~~competitive~~ plants in each plot from each replication. The mean ~~values of five plants was~~ were used for statistical analysis. Plant emergence ~~per cent~~ percent at 30 days after planting, plant height at 60 days after planting (cm), tuber girth (cm), tuber length (cm), number of tubers per plant, average weight of tuber yield per plant and tuber yield per plot were measured.

Biochemical analysis:

For tuber dry matter, total soluble solids (TSS), specific gravity, ascorbic acid, and protein, biochemical analysis was done in ~~the lab of~~ the Department of Vegetable Science at the G. B. Pant University of Agriculture and Technology.

Statistical analysis

The direct and indirect effects and correlation were estimated through path coefficient analysis as described by Dewey and Lu (1959) and Snedecor and Cochran (1987), respectively, using SAS 9.2 statistical package.

Results and discussion

Character associations

The phenotypic and genotypic correlations among various traits are presented in **Table 2**. All thirteen characters under study showed higher genotypic correlation coefficients than phenotypic correlation coefficients. Tuber yield per plot showed significant and positive correlation with average weight of tuber per plant (0.961 and 0.932), number of tuber per plant (0.517 and 0.510), tuber length (0.352 and 0.343), tuber girth (0.324 and 0.314), plant height (0.318 and 0.308), and specific gravity of tuber (0.2056 and 0.194) at genotypic and phenotypic level respectively. ~~Number~~ The number of branches per plant was found to be significant and positive correlated with plant height (0.493 and 0.467), tuber length (0.351 and 0.333) and tuber girth (0.349 and 0.332) both at genotypic and phenotypic ~~level~~ levels respectively. Tuber girth had a significantly high and positive correlation for tuber length (0.641 and 0.626) and tuber yield per plot (0.324 and 0.314) both at genotypic and phenotypic ~~level~~ levels respectively. ~~Dry~~

Comment [M13]: Please mention the unit of measurement in the results section as well.

matter content of tubers showing high significant and positive correlation for the specific gravity of the tubers (0.335 and 0.326) both at genotypic and phenotypic levels respectively. Ascorbic acid content of tuber exhibited significantly high and positive correlation with protein content of tuber (0.506 and 0.487) both at genotypic and phenotypic level respectively. Panigrahi *et al.* (2017) reported total tuber yield at early harvest showed significant positive correlation with germination percent, marketable tuber yield. Patel *et al.* (2018) they reported that a significantly significant positive correlation were-was observed between total tuber yield with plant height, number of tubers per plant, average weight of tubers per plant and marketable tuber yield. A previous worker also found similar results, positive correlation of tuber yield with number of tubers per plant also reported by Many other researchers have reported similar results for positive correlation of tuber yield with number of tubers per plant (Shubha, and Singh, (2018); Lavanya *et al.* (2020), Gebreselassie *et al.* (2022); Nigussie *et al.* (2023) and Sandilya *et al.* (2023). Similar results on exhibiting a correlation between tuber yield and weight of tubers was were reported by Shubha, and Singh (2018), Singh *et al.* (2021), Gebreselassie *et al.* (2022) and Tessema *et al.* (2023). Highly significant positive correlation of tuber yield with plant height was also reported by Tessema *et al.* (2023) and Tsagaye *et al.* (2023); Gusain (2010) and Hunde *et al.* (2022) reported that tuber size and specific gravity of tuber has a positive correlation with tuber yield.

Comment [M14]: Incomplete sentence, please rephrase.

Path coefficient analysis

~~Genotypic~~ The genotypic correlation coefficient of various yield attributing characters of tuber yield was further partitioned-categorized into their direct and indirect effects.

Direct effect on tuber yield

Comment [M15]: Reads more like statements rather than explaining the results with the help of available related studies

The values of the path coefficient for tuber yield per plot and its components are shown in **Table 3**. The highest positive direct effect which contributed towards tuber yield per plot was observed via average weight of tuber yield per plant (1.769), followed by number of branches per plant (0.251), number of tubers per plant (0.120), protein content of tuber (0.018) and specific gravity of tuber (0.017), whereas the effect of other traits were low (≤ 0.013). Negative direct effects on tuber yield per plant was also exhibited by plant emergence percent 30 days after planting (-0.033), dry matter content of tuber (-0.025), tuber length (-0.017) and ascorbic acid content of tuber (-0.003). In several studies, positive direct effect of tuber weight per plant on tuber yield has been reported (Tripura *et al.* 2016 and Lavanya *et al.* 2020 and Tripura *et al.* 2016). Similarly, Lavanya *et al.* 2020 and Sahu *et al.* 2023 observed observed positive direct effect on tuber yield per plot. Shubha and Singh (2018); Lavanya *et al.* 2020; Kumar *et al.* 2020; Kumar *et al.* 2022 and Sandilya *et al.* 2023 reported the highest direct effect of number of tubers per plant on tuber yield. Sahu *et al.* 2023 and Sandilya *et al.* 2023 reported highest direct effect of number of branches per plant on tuber yield. Shubha and Singh (2018) reported tuber size, specific gravity of tuber, total soluble solid content of tuber, protein content of tuber have high positive direct effect on tuber yield. Gusain (2010) reported positive direct effect on tuber yield by protein content of tuber, specific gravity of tuber and total soluble solid content of tuber. Patel *et al.* (2018) reported positive direct effect of total soluble solid content of tuber on total tuber yield. Sandilya *et al.* 2023 and Panigrahi *et al.* 2017 reported plant emergence have direct negative effect on tuber yield. Sattar *et al.* 2007 and Panigrahi *et al.* 2017 observed dry matter content have direct negative effect on tuber yield.

Indirect effect of yield components on tuber yield

Comment [M16]: Please re-write

Comment [M17]: Confusing statement

Comment [M18]: Please follow a certain style and remain consistent throughout the whole draft

The indirect effect of various traits on tuber yield per plot ~~are is~~ presented in **Table 3**. The plant emergence ~~per cent~~ exhibited a positive indirect effect on tuber yield per plot through dry matter content of tuber (0.011). ~~Number~~ ~~The number~~ of branches per plant had a positive indirect effect on tuber yield per plot via height of the plant at 60 days after planting (0.003). The positive indirect effect of number of ~~tuber~~ tubers per plant on tuber yield per plot was exhibited through ~~the~~ total soluble solid content of ~~the~~ tuber (0.012) and ~~the~~ number of branches per plant (0.005). ~~Average~~ ~~The average~~ weight of tuber per plant had a positive indirect effect on tuber yield per plot was exhibited through number of tubers per plant (0.914), tuber length (0.623), tuber girth (0.574), plant height at 60 days after transplanting (0.563), specific gravity of tuber (0.365), weight loss of tuber at 20 days after harvesting (0.301) and plant emergence ~~per cent~~ at 30 days after planting (0.132). The positive indirect effect of plant height at 60 days after planting on tuber yield per plot was exhibited through ~~the~~ number of branches per plant (0.006) and it also showed negative indirect effects with ~~the~~ number of tubers per plant (-0.001). The positive indirect effect of tuber girth on tuber yield per plot was ~~exhibited~~ ~~estimated~~ through length of ~~the~~ tuber (0.004) and it also showed negative indirect effects with ~~the~~ number of tubers per plant (-0.001). The positive indirect effect of tuber length on tuber yield per plot was exhibited through ~~the~~ protein content of tuber (0.006) and it also showed negative indirect effects with ~~the~~ number of branches per plant (-0.006). The dry matter content of ~~the~~ tubers have negative direct effect on tuber yield per plot exhibited through ~~the~~ specific gravity of ~~the~~ tuber (-0.008) and it also showed positive indirect effects with ~~the~~ number of tubers per plant (0.006). The positive indirect effect of ~~the~~ specific gravity of ~~the~~ tuber on tuber yield per plot was exhibited through plant emergence ~~per cent~~ at 30 days after planting (0.003), dry matter content of tuber (0.006) and total soluble solid

content of tuber (0.001). Weight loss of tuber at 20 days after harvesting had a positive indirect effect on tuber yield per plot ~~was exhibited~~ through tuber length (0.004). Total soluble solid content of tuber had ~~positive indirect effect on tuber yield per plot was exhibited~~ through plant emergence percent at 30 days after planting (0.004) and specific gravity of tuber (0.001).

Comment [M19]: What does it? Author(s) have used this term very frequently. Author(s) can consider several other different words

Comment [M10]: Getting tired of the excessive use of these words

~~Ascorbic-The ascorbic~~ acid content of tuber had a positive indirect effect on tuber yield per plot was exhibited through plant emergence ~~per centpercent~~ at 30 days after planting (0.001). The positive indirect effect of ~~the~~ protein content of tuber on tuber yield per plot was exhibited through number of tubers per plant (0.005) and ascorbic content of tuber (0.009). Path coefficient analysis revealed the average weight of tuber per plant, number of tubers per plant, height of plant at 60 days after planting, girth of tuber and number of branches per plant being the most important variables to increase tuber yield.

The above findings of path studies are ~~in accordance with~~ under the findings of Lavanya *et al.* 2020 for ~~the~~ number of branches per plant, plant height, number of tubers per plant, and tuber yield per plant. Shubha and Singh (2018) reported tuber size ~~have-has~~ ~~negative~~ ~~negative~~ indirect effects ~~with-on~~ the number of tubers per plant and ~~the~~ similar results were found for ~~the~~ specific gravity of ~~the~~ tuber, total soluble solid content of ~~the~~ tuber, ~~the~~ ascorbic acid content of tuber and protein content of tuber. Tripura *et al.* 2016 also found tuber length ~~have-has~~ indirect negative effect ~~with-on~~ number of branches per plant. Gebreselassie *et al.* 2022 found that ~~the~~ dry matter content of tuber have a positive indirect effect ~~with-on~~ the number of tubers per plant.

Conclusion

In the present investigation, tubers yield per plot showed a highly significant and positive correlation with the average weight of tubers per plant, number of tubers per plant, tuber length, tuber girth, number of branches per plant, and plant height. These research findings, which may allow for the direct selection of genotypes for the future breeding programme. Direct The direct effect of the average weight of tubers per plant, number of tubers per plant, number of branches per plant, plant height, tuber girth, specific gravity of tuber, total soluble solid content and protein content of tuber shows these are the most important direct influencing yield contributing characters. For improving potato tuber yield, the traits with the highest positive effects may be used in genotype selection. Based on character association and path analysis, it has been determined that simultaneous selection of the above mentioned traits average weight of tuber per plant, number of tubers per plant, number of branches per plant, plant height, tuber length, tuber girth and specific gravity of tuber will be more fruitful when choosing desirable genotypes because these characteristics exhibit significant and positive correlations as well as a strong direct influence on tuber yield.

REFERENCE

Asghari M, Kalantar M, Hassanpanah D, Dehghani Zahedani M 2022 Genetic varieties of the obtained hybrids from the Potato (*Solanum tuberosum* L.) tuber intersections in the spring using the factor and cluster analysis. *Journal of Vegetables Sciences* 5(2): 163-79.

Bhatt GM (1973) Significance of path coefficient analysis in determining the nature of character association. *Euphytica* 22(2):338-43.

Comment [M11]: Please double-check and follow the journal's guidelines for references. Also, remain consistent throughout this section. In some places, author(s) italicized journal names and did not italicize in many other references.

Dewey DR, Lu K (1959) A correlation and path coefficient analysis of components of crested wheatgrass seed production 1. *Agronomy journal* 51(9):515-8.

FAOSTAT, F (2019) Food and Agriculture Organization of the United Nations-Statistic Division <https://www.fao.org/faostat/en/#data>.

FAOSTAT. Available online: <https://www.fao.org/faostat/en/#data/QCL> (accessed on 23 November 2021).

Food and Agriculture Organization (FAO) (2022) FAOSTATdatabase. Available at: <http://www.fao.org/faostat/en/#data/QC> (Accessed 19 February 2022).

Gebreselassie H, Ajema L (2022) Correlation, Path Coefficient and Multivariate Analysis for Yield and Yield associated Traits among Potato (*Solanum tuberosum* L.) Genotypes Grown in Eastern Ethiopia. *World J. Agric. Soil Sci.* 8(2):1-7.

Gusain, P (2010) Evaluation of potato (*Solanum tuberosum* L.) germplasm for growth and yield characters under *TERAI* condition in Uttarakhand (Thesis M.Sc. GB Pant University of Agriculture and Technology, Pantnagar-263145, Uttarakhand, India).

Hijmans RJ (2001) Global distribution of the potato crop. *American journal of potato research* 78:403-12.

Hunde NF, Galalcha DT, Limeneh DF (2022) Correlation and path coefficient analyses of tuber yield and yield components among potato (*Solanum tuberosum* L.) genotypes at Bekoji, Southeastern Ethiopia. *International Journal of Agricultural Research, Innovation and Technology* 12(2):144-54.

Killick RJ (1977) Genetic analysis of several traits in potatoes by means of a diallel cross. *Annals of applied biology* 86(2):279-89.

Kumar R, Prakash S, Luthra SK, Singh B, Chand P, Kumar V, Singh R, Alam K (2022) Analysis of correlation and path coefficient among the yield and yield attributes characters in potato (*Solanum tuberosum* L.). *InBiol.* 916-922.

Kumar V, Mishra DP, Kumar V, Tiwari AK (2020) Correlation and path coefficient analysis for yield and components traits in different genotypes of potato (*Solanum tuberosum* L.) under Eastern Uttar Pradesh condition. *IJCS.*8:2866-70.

Lavanya KS, Srinivasa V, Ali S, Devaraju, Lakshmana D, Kadian MS (2020) Correlation and path analysis for yield and yield-related traits of potato (*Solanum tuberosum* L.) in Karnataka. *National Academy Science Letters* 43(2):137-40.

Luthra SK, Gupta VK, Kaundal B, Tiwari JK (2018) Genetic analysis of tuber yield, processing and nutritional traits in potato (*Solanum tuberosum*). *Indian Journal of Agricultural Sciences* 88(8):1214-21.

Mahgoub HA, Eisa GS, Youssef MA (2015) Molecular, biochemical and anatomical analysis of some potato (*Solanum tuberosum* L.) cultivars growing in Egypt. *Journal of Genetic Engineering and Biotechnology* 13(1):39-49.

Nigussie HL, Anilay AM, Getie MA, Kasssa MW (2023) Variability and association among tuber yield and yield related traits of potato (*Solanum tuberosum* L.) genotypes in Debre markos district, north west Ethiopia. *Ecological Genetics and Genomics* 10:100180.

Panigrahi KK, Pradhan J, Panigrahi P, Sarkar KK (2017) Genetic variability, character association and path coefficient analysis of yield attributes for medium and late maturing potato cultivars. *International Journal of Current Microbiology and Applied Sciences* 6(7):2558-66.

Patel RA, Bhavani RT, Patel DK, Patel JR (2018) Character association and path coefficient studies for yield and its components in Potato (*Solanum tuberosum* L.). *Int. J. Curr. Microbiol. App. Sci.* 7(8):1942-50.

Rajani. Morphological, biochemical and molecular characterization in potato (*Solanum tuberosum* L.) germplasm (Doctoral dissertation, GB Pant University of Agriculture and Technology, Pantnagar-263145, Uttarakhand, India). 2015

Sahu P, Chaurasiya PC, Rangare NR, Kumar V, Shandilya VK, Singh T, Dehari B (2023) Study on genetic variability, correlation and path analysis in F1 potato genotypes (*Solanum tuberosum* L.).

Sandilya VK, Rangare NR, Saran D, Kumar R, Kumar V, Ameen G, Dhiwar K, Dehari B, Sahu P, Khokhar D (2023) Character association and path coefficient analysis between tuber yield and yield components in potato (*Solanum tuberosum* L.) genotypes.

Sattar MA, Sultana N, Hossain MM, Rashid MH, Islam AA (2007) Genetic variability, correlation and path analysis in potato (*Solanum tuberosum* L.). *Bangladesh Journal of Plant Breeding and Genetics* 20(1):33-8.

Shubha K, Singh D (2018) Selection of yield-associated morphological and biochemical traits using correlation and path coefficient analysis in potato (*Solanum tuberosum* L.) in the foothills of North-Western Himalayas. *Potato Research* 61:273-81.

Singh S, Mishra DP, Singh N, Singh S, Singh SP (2021) Genetic variability and associations studies for yield and its component traits in potato (*Solanum tuberosum* L.).

Snedecor W, Cochran WG (1987) Statistical methods. Oxford and IBM Calcutta p 593.

Spooner DM, Ghislain M, Simon R, Jansky SH, Gavrilenko T (2014) Systematics, diversity, genetics, and evolution of wild and cultivated potatoes. *The botanical review* 80:283-383.

Sukhchain, Sidhu BS (1992) Correlation and path coefficients analysis for reproductive traits in Guinea grass. *Euphytica* 60:57-60.

Tessema L, Mohammed W, Abebe T (2023) Association of traits in potato (*Solanum tuberosum* L.) cultivars evaluated in Central Ethiopia Ethiopian Journal of Science and Technology 16(3):225-36.

Tripura A, Das A, Das B, Priya B, Sarkar KK (2016) Genetic studies of variability, character association and path analysis of yield and its component traits in potato (*Solanum tuberosum* L.). *Journal of Crop and Wee* 12:56-63.

Tsagaye D, Fikre D, Fufa N, Gebretensay F, Ali A, Wegayehu G (2023) Agronomic Evaluation of Potato (*Solanum tuberosum* L.) Genotypes for Tuber Yield and Yield Attributing Traits. *International Journal of Bio-resource and Stress Management* 14:719-25.

Woolfe JA, Poats SV (1987) The potato in the human diet. Cambridge University Press.

Table 1 List of Potato (*Solanum tuberosum* L.) genotypes under study and source of seed

| Sl. No. | Genotype | Source | Sl. No. | Genotype | Source |
|---------|----------|--------------|---------|----------|--------------|
| 1 | K. Surya | CPRI, Shimla | 17 | K.Khyati | CPRI, Shimla |
| 2 | P-29 | CPRI, Shimla | 18 | P-25 | CPRI, Shimla |
| 3 | PH-3 | CPRI, Shimla | 19 | P-23 | CPRI, Shimla |
| 4 | C-8 | CPRI, Shimla | 20 | P-30 | CPRI, Shimla |
| 5 | P-33 | CPRI, Shimla | 21 | P-31 | CPRI, Shimla |

| | | | | | |
|----|------------|--------------|----|-----------|--------------|
| 6 | P-11 | CPRI, Shimla | 22 | P-40 | CPRI, Shimla |
| 7 | P-7 | CPRI, Shimla | 23 | P-27 | CPRI, Shimla |
| 8 | P-12 | CPRI, Shimla | 24 | C-14 | CPRI, Shimla |
| 9 | P-14 | CPRI, Shimla | 25 | MM-11 | CPRI, Shimla |
| 10 | K.Sindhuri | CPRI, Shimla | 26 | P-34 | CPRI, Shimla |
| 11 | C-17 | CPRI, Shimla | 27 | C-6 | CPRI, Shimla |
| 12 | C-20 | CPRI, Shimla | 28 | K.Lalit | CPRI, Shimla |
| 13 | P-9 | CPRI, Shimla | 29 | P-4 | CPRI, Shimla |
| 14 | P-22 | CPRI, Shimla | 30 | K.Puskar | CPRI, Shimla |
| 15 | C-15 | CPRI, Shimla | 31 | K.Frysona | CPRI, Shimla |
| 16 | C-28 | CPRI, Shimla | 32 | P-15 | CPRI, Shimla |

Table 2 Phenotypic and genotypic coefficient of correlation for tuber yield and its attributing traits in potato

| Characters | | No. of branches per plant | Plant height at 60DAP (cm) | Tuber girth (cm) | Tuber length (cm) | No. of tubers per plant | Average weight of tuber per plant(g) | Tuber dry matter (%) | Specific gravity of tuber (g/cm ³) | Total soluble solid content of tuber (%) | Ascorbic acid content of tuber (mg/100g) | Protein per cent | Weight loss of tuber 20 DAH (%) | Tuber yield per plot (Kg) |
|--|----|---------------------------|----------------------------|------------------|-------------------|-------------------------|--------------------------------------|----------------------|--|--|--|------------------|---------------------------------|---------------------------|
| Plant emergence per cent at 30 DAP | rp | 0.140 | 0.218* | 0.198 | 0.030 | -0.154 | 0.049 | -0.236* | 0.111 | 0.275** | 0.271** | 0.158 | 0.173 | 0.049 |
| | rg | 0.241* | 0.320** | 0.338** | -0.003 | -0.234* | 0.075 | -0.308** | 0.156 | 0.366** | 0.373** | 0.231* | 0.075 | 0.075 |
| No. of branches per plant | rp | | 0.467** | 0.332** | 0.333** | -0.249* | 0.342 | 0.086** | -0.065 | 0.054 | 0.010 | -0.016 | 0.249* | 0.342** |
| | rg | | 0.493** | 0.349** | 0.351** | -0.263** | 0.336 | 0.091** | -0.084 | 0.063 | -0.021 | -0.016 | 0.272** | 0.336** |
| Plant height at 60 DAP (cm) | rp | | | 0.178 | 0.353** | -0.086 | 0.308** | 0.212* | 0.111 | -0.107 | 0.095 | -0.010 | 0.265** | 0.308** |
| | rg | | | 0.182 | 0.385** | -0.084 | 0.318** | 0.214* | 0.108 | -0.118 | 0.092 | -0.004 | 0.270** | 0.318** |
| Tuber girth (cm) | rp | | | | 0.626** | -0.162 | 0.314** | -0.078 | 0.088 | 0.164 | 0.051 | -0.049 | 0.149 | 0.314** |
| | rg | | | | 0.641** | -0.164 | 0.324** | -0.088 | 0.095 | 0.169 | 0.048 | -0.052 | 0.147 | 0.324** |
| Tuber length (cm) | rp | | | | | -0.140 | 0.343** | 0.072 | 0.267** | -0.002 | -0.311** | -0.333** | 0.309** | 0.343** |
| | rg | | | | | -0.151 | 0.352** | 0.077 | 0.278** | 0.001 | -0.319** | -0.352** | 0.320** | 0.352** |
| Number of tubers per plant | rp | | | | | 0.510** | -0.231* | 0.183 | -0.524** | -0.066 | 0.247* | | -0.161 | 0.511** |
| | rg | | | | | | 0.517** | -0.233* | 0.195 | -0.535** | -0.060 | 0.258* | -0.163 | 0.517** |
| Average weight of tuber per plant(g) | rp | | | | | | | -0.012 | 0.194 | -0.315** | -0.170 | -0.049 | 0.167 | 0.932** |
| | rg | | | | | | | -0.011 | 0.206* | -0.317** | -0.165 | -0.050 | 0.170 | 0.961** |
| Tuber dry matter (%) | rp | | | | | | | | 0.326** | 0.145 | -0.237* | 0.090 | -0.039 | -0.012 |
| | rg | | | | | | | | 0.335** | 0.147 | -0.246* | 0.092 | -0.044 | -0.011 |
| Specific gravity of tuber (g/cm ³) | rp | | | | | | | | | 0.073 | -0.159 | 0.015 | -0.069 | 0.194 |
| | rg | | | | | | | | | 0.069 | -0.171 | 0.014 | -0.074 | 0.206* |
| Total soluble solid content of tuber (%) | rp | | | | | | | | | | -0.028 | -0.121 | -0.067 | -0.315** |
| | rg | | | | | | | | | | -0.031 | -0.123 | -0.077 | -0.317** |
| Ascorbic acid content of tuber (mg/100g) | rp | | | | | | | | | | | 0.487** | -0.223* | -0.171 |
| | rg | | | | | | | | | | | 0.506** | -0.235* | -0.165 |
| Protein per cent of tuber | rp | | | | | | | | | | | | -0.440** | -0.049 |
| | rg | | | | | | | | | | | | -0.459** | -0.050 |
| Tuber yield per plot (Kg) | rp | | | | | | | | | | | | | 1.000 |
| | rg | | | | | | | | | | | | | 1.000 |

*,** Significant at P = 5% and 1% levels, respective

Table 3 Genotypic path coefficient showing direct and indirect effects of different characters on tuber yield in potato

| Characters | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 | |
|------------|-------|--------|--------|--------|--------|-------|--------|--------|-------|--------|--------|--------|--------|--------|
| 1 | | -0.033 | -0.008 | -0.011 | -0.011 | 0.001 | 0.008 | -0.003 | 0.011 | -0.005 | -0.012 | -0.012 | -0.008 | -0.008 |
| 2 | 0.001 | | 0.251 | 0.003 | 0.001 | 0.001 | -0.002 | 0.000 | 0.001 | 0.000 | 0.000 | 0.000 | 0.001 | 0.001 |
| 3 | 0.004 | 0.006 | | 0.013 | 0.002 | 0.005 | -0.001 | 0.004 | 0.003 | 0.001 | -0.002 | 0.001 | -0.001 | 0.003 |

| | | | | | | | | | | | | | |
|----|-------|--------|--------|--------------|---------------|--------------|--------------|---------------|--------------|--------------|---------------|--------------|--------------|
| 4 | 0.003 | 0.003 | 0.001 | 0.007 | 0.004 | -0.001 | 0.002 | -0.001 | 0.001 | 0.001 | 0.001 | -0.004 | 0.001 |
| 5 | 0.000 | -0.006 | -0.007 | -0.011 | -0.017 | 0.003 | -0.006 | -0.003 | -0.005 | 0.000 | 0.006 | 0.006 | -0.006 |
| 6 | 0.005 | 0.005 | 0.002 | 0.003 | 0.003 | 0.120 | -0.011 | 0.005 | -0.004 | 0.012 | 0.001 | -0.005 | 0.003 |
| 7 | 0.132 | -0.063 | 0.563 | 0.574 | 0.623 | 0.914 | 1.769 | -0.019 | 0.365 | -0.561 | -0.293 | -0.088 | 0.301 |
| 8 | 0.008 | -0.002 | -0.005 | 0.002 | -0.002 | 0.006 | 0.001 | -0.025 | -0.008 | -0.004 | 0.006 | -0.003 | 0.001 |
| 9 | 0.003 | -0.002 | 0.002 | 0.002 | 0.005 | 0.003 | 0.004 | 0.006 | 0.017 | 0.001 | -0.003 | 0.002 | -0.001 |
| 10 | 0.004 | 0.001 | -0.001 | 0.002 | 0.000 | -0.005 | -0.003 | 0.002 | 0.001 | 0.011 | -0.001 | -0.002 | -0.001 |
| 11 | 0.001 | 0.001 | -0.001 | -0.001 | 0.001 | 0.001 | 0.001 | 0.001 | 0.001 | 0.001 | -0.003 | -0.001 | 0.001 |
| 12 | 0.004 | -0.003 | -0.001 | -0.009 | -0.006 | 0.005 | -0.001 | 0.007 | 0.002 | -0.002 | 0.009 | 0.018 | -0.008 |
| 13 | 0.003 | 0.003 | 0.003 | 0.002 | 0.004 | -0.002 | 0.002 | -0.001 | -0.001 | -0.001 | -0.003 | -0.006 | 0.012 |

Note: Bold diagonal value indicate direct effect

Residual effect = 0.0028

* and ** Significant at P = 5% and 1% levels respectively.

1: Plant emergence per cent at 30 days after planting, 2: Number of branches per plant, 3: Plant height at 60 days after planting (cm), 4 : Tuber girth (cm), 5: Tuber length (cm), 6: Number of tubers per plant, 7: Average weight of tuber per plant(g), 8: Dry matter content of tuber (per cent), 9: Specific gravity of tuber (g per cm³), 10: Total soluble solid content of tuber (per cent), 11 : Ascorbic acid content of tuber (mg per 100g) 12: Protein content of tuber (per cent) and 13: Weight loss of tuber at 20 days after harvesting (per cent).