

## Quality control of the biological properties of hydrazide-hydrazonederivatives from strains responsible for food poisoning isolated from foods sold on the corridors of the town of Daloa (Côte d'Ivoire)

### Abstract :

Bacteria are a real public health problem because of their involvement in many diseases. Their resistance to antibiotics has become one of the most important problems in the fight against infectious diseases worldwide. The aim of this work is to evaluate the antibacterial activity of hydrazide-hydrazonederivatives against environmental strains using the agar diffusion method in cups and the macro-dilution method in liquid medium.

In this work, a survey was conducted among food vendors on the four security corridors of the city of Daloa. A structured questionnaire was administered to 127 vendors and a total of 60 food samples were collected from the four corridors and analysed.

The results indicate that the majority of food sales at these security corridors are carried out by women (88.12%). The vendors were mostly adults (59.84%) aged between 15 and 35, with no formal education (50.39%). Microbiological analysis of the 2 types of food showed the presence of enterobacteria above the recommended thresholds. With regard to the biological properties explored, four substances (AEV5, AB2, AB3 and AB5) revealed antibacterial activity. The MBC/MIC ratio of all these substances being less than 4 indicates that these synthesised substances all have bactericidal activity. As a result, substances derived from S-alkylation and Mannich bases can be used to combat bacterial infections.

**Keywords:** biological properties, hydrazide-hydrazonederivatives, food poisoning, bacterial strains, enterobacteria, food, Daloa

### Introduction

Food is the first consumer good used to satisfy the primary needs of every human being. The quality of this food is vital to the health of consumers (Fung *et al.*, 2018). Food is the main source of micro-organisms when it is not eaten in a healthy way. Every year, several million people suffer from food-borne illnesses (Kirk *et al.*, 2015 ; Fung *et al.*, 2018). These food-borne illnesses are generally known as food poisoning. They are diseases that are often infectious and accidental, contracted as a result of ingesting food contaminated with pathogenic bacteria. Such contamination generally results from inadequate food handling, preservation or cooking methods (Ferreira *et*

*al.*, 2013). These infectious diseases are a global concern. The number of cases of food poisoning is constantly rising around the world. Every year, there are more than 600 million cases of food poisoning in the world (OMS, 1996; Fleckenstein *et al.*, 2021). There have been numerous cases of food poisoning in Côte d'Ivoire. In May 2022, in the town of Bondoukou, in the east of the country, around a hundred people aged between 2 and 32 suffered from food poisoning after eating a dairy product sold in an ice cream parlour (Ministère de la santé, 2022). Food-borne diseases are therefore a major concern for governments (Bernard, 2014). Most are caused by bacteria such as *Listeria monocytogenes*, *Escherichia coli*, *Staphylococcus aureus*, *Salmonella spp*, *Clostridium spp* and *Campylobacter spp* (Antunes *et al.*, 2020; Vieira *et al.*, 2021). Antibiotics are most commonly used to treat this condition. Unfortunately, the misuse of antibiotics and the horizontal transfer of resistance genes have led to the emergence of bacteria resistant to certain antibiotics. Among the concerns raised by this issue, bacterial resistance to antibiotics has been recognised by the World Health Organisation as one of the most serious threats to global health, food security and development. Its spread could change medicine in the coming years. Antibiotic resistance is all the more worrying given that the use of antibiotics is often uncontrolled (Pons *et al.*, 2014). Faced with the constant threat of infectious agents, it is essential to find new antimicrobial molecules. Hydrazide-hydrazon derivatives could offer an alternative for better management of cases of food poisoning (Popiołek, 2017). Indeed, these substances are a potential source for strengthening the fight against bacteria that are increasingly resistant to antibiotics. The main objective of this work is to evaluate the antibacterial activity of hydrazide-hydrazon derivatives synthesised on strains isolated from street food sold in the commune of Daloa.

The specific objectives were to

- To search for the presence of germs responsible for food poisoning in food stuffs sold on the four security corridors in the town of Daloa;
- To demonstrate the antibacterial effect of hydrazide-hydrazon derivatives on strains from food stuffs;
- determine the antibacterial parameters (Minimum Inhibitory Concentration, Minimum Bactericidal Concentration) of these substances.

## **Materials and methods**

### **Materials**

## Presentation of the study area

Located in the centre-west of Côte d'Ivoire, with a population of 705,378 (INS, 2021), the town of Daloa is 141 km from the political capital Yamoussoukro and 383 km from Abidjan, the economic capital. Geographically, Daloa is located at latitude 6°53 north and longitude 6°27 west. It is the capital of the Haut Sassandra region. It is the largest town in the west of the country. It is a major crossroad linking the West with the North and the West with the South of the country via national roads. From north to south, it links the towns of Issia and Vavoua, and from east to west, it connects Duékoué to Yamoussoukro.




Figure 1: Map of the town of Daloa (Bolou, 2021)

## Biological material

### Bacterial strains

The microbial support consists of :

- Bacterial strains isolated and identified from foodstuffs sold on the various corridors. A total of six (6) strains were selected. These were four (4) strains of *E. coli* coded *E. coli* A1, *E. coli* A2, *E. coli* A3, *E. coli* A4 and two (2) strains of Enterobacteriaceae coded *Ent* P1, *Ent* P2.

- A strain of *E. coli* resistant to third generation cephalosporins (C3G ®) from the bio-collection of the Centre Nationale de Référence des Antibiotiques of the Institut Pasteur (Abidjan, Côte d'Ivoire) used for quality control of the tests.

### Other materials

- **Hydrazide-hydrazones derivatives and mannich bases** : 4 codified substances (AEV3, AEV5, AEV6, AEV8) derived from benzimidazole S-alkylation and 5 codified mannich base substances (AB2, AB3, AB4, AB5 and AB7) synthesised and described as new bioactive

molecules were provided by the Laboratory of Constitution and Reaction of Matter, Department of Sciences of Structures of Matter and Technology, Félix Houphouët-Boigny University, Abidjan, Côte d'Ivoire for this study.

## **Methods**

### **Socio-demographics survey**

A socio-demographic survey of the various security corridors (Man, Issia, Bouaflé and Vavoua) was carried out over a period of one month (from 31 March to 30 April 2022). The survey sought information on the age, gender and level of education of the vendors. It also looked at the hygienic state of food preparation, conditions of sale, conditions of food preparation and storage, and the level of knowledge about hygiene practices among male and female vendors.

### **Sample collection and transport**

Food samples were collected on four security corridors in the city of Daloa, namely the Issia corridor (South), the Vavoua corridor (North-West), the Bouaflé corridor (East) and the Man corridor (West). A total of 30 solid food samples and 30 liquid food samples were collected from vendors. Samples were taken aseptically. The samples were packaged in labelled sterile jars, then placed in a cooler containing an ice accumulator and transported to the laboratory.

### **Research and isolation of strains**

The preparation of the mother suspension of samples and decimal dilutions was carried out in accordance with standard NF EN ISO 6887-1, 2017, which defines the general rules for the preparation of the mother suspension and dilutions. Culture media were prepared and used in accordance with the manufacturers' instructions. Two types of inoculation were carried out. Surface inoculation for the detection of *E. coli* on Rapid'*E. coli* medium (Bio-Rad, France) and inoculation in the mass with VRBL (Biokar, France) and VRBG (Biokar, France) media for the detection of Enterobacteriaceae. Cultures were incubated at 37°C for 24 hours. Colonies were read and counted, identified and preserved in 10% glycerol water.

### **Preparation of solutions of hydrazide-hydrazonederivatives and mannich bases**

Stock solutions of the chemical substances were prepared at a concentration of 16 µg/mL. The substances in powder form were first diluted in ethanol at 96°C.

### **Sensitivity testing and determination of inhibition diameters**

The Mueller-Hinton agar diffusion technique used by Bssaibi *et al.* in 2009 was used in this study. This technique involved obtaining young colonies transplanted onto nutrient agar

using the streak method. A bacterial inoculum estimated at  $10^6$  bacteria/mL adjusted by opacity to 0.5 McFarland was prepared in physiological water (NaCl 0.9%). 1 mL of this inoculum was then swabbed onto Muller-Hinton agar (GMH, Bio-Rad, France). Cups were made by sinking the large end of a sterile Pasteur pipette into the inoculated Muller-Hinton agar and left to dry. These plates were then filled with 50  $\mu$ L of the chemical preparation and incubated at 37°C for 24 hours. The inhibition diameters were then determined by caliper measurement. The effectiveness of the substances was assessed according to the criteria of Ponce *et al* (2003). Thus, a substance is said to be ineffective if the inhibition diameter is less than 8 mm, effective if the diameter is between 9 and 14 mm, very effective for a diameter between 15 and 19 mm and extremely effective if the diameter is greater than 20 mm.

#### **Determination of antibacterial parameters Minimum Inhibitory Concentration (MIC) and Minimum Bactericidal Concentration (MBC)**

Antibacterial parameters were determined for the so-called effective substances using the liquid macro-dilution method described by (Koné *et al.*, 2004).

#### **Preparation of the bacterial inoculum**

A bacterial inoculum was prepared from a young colony in 10 mL of sterile MH broth and incubated at 37°C for 3 hours. Then 0.3 mL of this pre-culture was diluted in 10 mL of sterile MH broth and homogenised.

#### **Preparation of test concentration ranges**

A concentration range for the test substances was obtained using the double dilution method. A stock solution with a concentration of 1000  $\mu$ g/mL (2 mg in 2 mL of 70° ethanol) was prepared. A series of 2-fold dilutions were made from this solution to obtain concentration ranges from 500  $\mu$ g/mL to 7.81  $\mu$ g/mL.) Then, in sterile haemolysis tubes, 1 mL of each concentration range of the substance was brought into contact with 1 mL of bacterial inoculum. In a growth control tube, 1 mL of sterile distilled water was added to the inoculum, while 2 mL of sterile Muller-Hinton broth (MHB) was added to the sterility control. The preparations were incubated for 24 hours at 37°C. Afterwards, observation with the naked eye was carried out and the lowest concentration for which no bacterial growth was observed with the naked eye corresponded to the MIC.

The MBC is determined by transferring a sample of the tubes showing no growth to a fresh medium devoid of the molecules to be tested. This involved diluting the starting inoculum from  $10^{-1}$  to  $10^{-4}$  and inoculating these different dilutions using a calibrated 2  $\mu$ L loop in 5 cm long streaks on Muller-Hinton agar, then incubating at 37°C for 24 hours (preparation A).

After reading the MICs, the contents of tubes in which there was no visible growth were used to inoculate the GMH on 5 cm long streaks (preparation B). The MIC was determined by comparing the bacterial growth of preparations A and B. The smallest concentration in the tube with less than 0.01% viable bacteria compared to the initial inoculum is the MBC.

The MBC/MIC ratio was also measured to show the bacteriostatic and bactericidal effects of betelleaves. If the ratio is  $\leq 2$ , the samples are considered to be bactericidal agents. The bacteriostatic mode of action is reflected when the ratio is  $\geq 4$  (Konaté *et al.*, 2012 ; Abdi et Dego, 2019 ; Nayaka *et al.*, 2021).

### **Statistical analysis**

All tests were carried out in triplicate and the numerical values obtained are expressed as the arithmetic mean plus the corresponding statistical standard deviation. STATISTICA 7.1 software was used for these analyses. Analyses of variance (ANOVA) based on multiple DUNCAN tests with a significance level of 5% were performed in order to compare the variables measured on different strains with the substances tested. Means, standard deviations and graphs were produced using EXCEL 2016.

## **Results and discussion**

### **Results**

#### **Socio-demographic profile of the people surveyed and their level of knowledge about hygiene**

The results of the survey of food vendors in the four corridors are shown in table I below. In terms of gender, the results show that 11.82% of vendors are men and 88.12% are women. In terms of level of education, vendors are poorly educated. Only 7.08% had completed higher education, 27.55% had attended secondary school, 14.96% had completed primary school, and 50.39% had no education at all. In terms of age, 15.74% of sellers are under 14, 59.84% are between 15 and 35, and 24.42% are over 36. In terms of knowledge of hygiene, the results show that 54.5% of respondents have some knowledge of hygiene and 45.5% have no knowledge of hygiene.

**Table I: Socio-demographic profiles of vendors on the various corridors**

<b>Socio-demographic profiles Staff Fréquency (%)</b>	
<b>Sex Male</b>	11,82
<b>Female</b>	88,12

<b>Age</b>	0-14 years	20	15,74
	15-35 years	76	59,84
	35 years and over	31	24,42
<b>Level of study</b>	No level of education	64	50,39
Primary	19	14,96	
Secondary	5	27,55	
	Supérieur	9	7,08
<b>Level of hygiene knowledge</b>	Oui	69	54,5
	Non	58	45,5

UNDER PEER REVIEW

### Antibacterial effect of chemicals derived from benzimidazole S-alkylation on bacterial strains

Table II below shows the inhibition diameters of the four benzimidazole S-alkylation substances (AEV3, AEV5, AEV6 and AEV8) on strains isolated from food. It should be noted that the four substances were ineffective on the 4 *E. coli* strains isolated because they had an inhibition diameter of less than 9 mm. The substances AEV5 and AEV8 were highly effective and effective respectively only on the Ent P2 strain with diameters of 17.3 mm and 10 mm. On the Ent P2 strain, all the substances in this chemical group were found to have no antibacterial effect.

**Table II: Diameters (mean  $\pm$  SD) of the zones of inhibition of chemical substances derived from benzimidazole S-alkylation on bacterial strains**

Bacterial strains	Inhibition diameters (mm)			
	S-alkylation benzimidazole			
	AEV3	AEV5	AEV6	AEV8
<i>E. coli</i> A1	0,8 $\pm$ 0,1 <sup>b</sup>	5,3 $\pm$ 0,6 <sup>a</sup>	0,6 $\pm$ 0,2 <sup>b</sup>	0,5 $\pm$ 0,0 <sup>b</sup>
<i>E. coli</i> A2	0,7 $\pm$ 0,2 <sup>a</sup>	0,8 $\pm$ 0,1 <sup>a</sup>	0,7 $\pm$ 0,2 <sup>a</sup>	0,7 $\pm$ 0,0 <sup>a</sup>
<i>E. coli</i> A3	0,7 $\pm$ 0,1 <sup>b</sup>	7,2 $\pm$ 0,6 <sup>a</sup>	0,7 $\pm$ 0,2 <sup>b9</sup>	0,5 $\pm$ 0,0 <sup>b</sup>
<i>E. coli</i> A4	0,7 $\pm$ 0,2 <sup>a</sup>	0,7 $\pm$ 0,1 <sup>a</sup>	0,6 $\pm$ 0,2 <sup>a</sup>	0,9 $\pm$ 0,0 <sup>a</sup>
Ent P1	0,6 $\pm$ 0,2 <sup>c</sup>	17,3 $\pm$ 0,6 <sup>a</sup>	0,8 $\pm$ 0,1 <sup>c</sup>	10 $\pm$ 0,0 <sup>b</sup>
Ent P2	0,8 $\pm$ 0,1 <sup>a</sup>	0,6 $\pm$ 0,2 <sup>a</sup>	0,7 $\pm$ 0,2 <sup>a</sup>	0,5 $\pm$ 0,1 <sup>a</sup>

NB: Values with the same letter are statistically identical, at the 5% threshold.

### Effect of various Mannich-based chemicals on the *in vitro* growth of bacterial strains

The results of the sensitivity of bacterial strains to Mannich-based substances are presented in Table III. All the substances were ineffective on the Ent P2 strain. No zone of inhibition was observed. The results also show that the substances AB2, AB3, AB4 and AB7 are extremely effective on the *E. coli* A2 strain with diameters between 21 and 27 mm and similarly for AB2 and AB3 on the *E. coli* A1 strain with diameters of 21.0  $\pm$  0.0 and 23.0  $\pm$  0.0 respectively. The largest inhibition zone diameters were observed on *E. coli* A2 (27.0  $\pm$  0 mm) for AB2 (Table III).

**Table III: Diameters (mean ± SD) of the zones of inhibition of Mannich-based substances on bacterial strains**

Bacterial strains	Inhibition diameters (mm)				
	Mannich-based				
	AB <sub>2</sub>	AB <sub>3</sub>	AB <sub>4</sub>	AB <sub>5</sub>	AB <sub>7</sub>
<i>E. coli</i> A1	21,2±0,8 <sup>b</sup>	23,0±0,0 <sup>a</sup>	—	18,3±0,5 <sup>c</sup>	13,0±1,6 <sup>d</sup>
<i>E. coli</i> A2	27,9±1,4 <sup>a</sup>	24,5±2,5 <sup>b</sup>	21,2±0,8 <sup>c</sup>	16,3±0,2 <sup>d</sup>	22,6±1,4 <sup>bc</sup>
<i>E. coli</i> A3	17,2±0,7 <sup>b</sup>	17,0±0,2 <sup>b</sup>	12,3±0,7 <sup>c</sup>	19,0±0,0 <sup>a</sup>	—
Ent P1	17,0±1,3 <sup>a</sup>	15,3±0,7 <sup>a</sup>	14,0±0,8 <sup>a</sup>	19,0±1,2 <sup>a</sup>	10,0±0,8 <sup>a</sup>
Ent P2	—	—	—	—	—

NB: Values with the same letter are statistically identical, at the 5% threshold.

#### Antibacterial parameters of synthesised chemicals

Table IV presents the results of the determination of antibacterial parameters (MIC, MBC and MBC/MIC) of the substances that showed an inhibitory action on the bacterial strains by their inhibition diameter. These were AEV5 and AEV8 on the *Ent P1* strain, AB2 on *E. coli* A2, AB3 on *E. coli* A1 and AB5 on *Ent P1*. The MBC/MIC ratio for all the substances was less than 4. The substances all have bactericidal power on the bacterial strains used.

**Table IV: Antibacterial parameters of benzimidazole S-alkylation derivatives and Mannich bases**

	S-alkylation benzimidazole		Mannich-based		
	AEV5	AEV8	AB2	AB3	AB5
	<b>Bacterial strains</b>				
	<i>Ent P1</i>	<i>Ent P1</i>	<i>E. coli</i> A2	<i>E. coli</i> A1	Ent P1
MIC (µg/ mL)	31,25	31,25	125	62,5	62,5
MBC (µg/ mL)	62,5	62,5	250	62,5	125
MBC/MIC	2	2	2	1	2
Effect	Bactericidal	Bactericidal	Bactericidal	Bactericidal	Bactericidal

## Discussion

The aim of this work was to evaluate the antibacterial activity of the chemical substances synthesised on bacterial strains responsible for food poisoning. The strains were isolated from foods sold on the various security corridors in the town of Daloa. The socio-demographic survey on the sale of food at these various corridors revealed that this activity is mainly carried out by women (88.12%), compared with 11.82% for men. Vendors are generally very young, with 59% aged between 15 and 35, and the vast majority have no formal education. These same observations have been made by Khomotso and Chelule (2020) and also by Attienet *et al* (2021), in fact the commercial activity of street food in Africa is mostly carried out by women and especially that linked to catering. In fact, according to Rahman *et al* (2016), the predominance of women in the street food sector could be due to the fact that women are the main breadwinners in the majority of households. The results of the microbiological analysis of the various foodstuffs revealed the presence of microorganisms such as: enterobacteria, faecal coliforms and *E. coli*. These germs have also been found in street foods in the work of certain authors such as Mosupye and Von Holy (2000), Attienet *et al*, (2022) and Zébré *et al*, (2022). The presence of these germs in these foods is thought to be due to handling by the vendors. These germs are indicators of poor hygiene. They are evidence of faecal contamination (Koffi-Nevryet *al.*, 2012). Hygiene conditions and the practices of food handlers in sales outlets are generally inadequate (Attienet *et al.*, 2022; Zébré *et al.*, 2022). A lack of understanding of the concept and importance of hygiene is thought to be behind the high levels of contamination in the various samples analysed (Cardinale *et al.*, 2005; Attienet *et al.*, 2021). Consumer health is therefore becoming a global concern. In order to combat food-borne illnesses effectively in the face of the growing threat of bacteria resistant to conventional antibiotics, new chemical molecules derived from benzimidazole S-alkylation and Mannich bases have been synthesised and tested on bacteria isolated from food. The results obtained with the benzimidazole S-alkylation substances show that only the Ent P1 strain is sensitive to the AEV5 and AEV8 substances. This observation is less surprising given that enterobacteria are highly resistant to antimicrobials (Iredellet *et al.*, 2016; Li *et al.*, 2021). According to Zniberet *et al*, (2009), the lack of effect could be explained by taking into account the structure of the molecule tested (the size of the molecules) as well as the physiology and genetic potential of the target cell. But according to Iredellet *et al*, (2016), Enterobacteriaceae resistance to antibiotics, particularly  $\beta$ -lactam antibiotics, is increasingly dominated by the mobilisation of single continuously expressed genes that code for effective drug-modifying enzymes. As for

Mannich-based substances, they have almost all shown antibacterial activity on all strains with the exception of the *Ent P2* strain. The antibacterial activity of these substances has been demonstrated in several previous studies. Indeed, this family of substances is effective on both Gram+ and Gram- bacteria (Gheorghe, 2015). Determination of the antibacterial parameters shows that the substances have an antibacterial effect. The MBC/MIC ratio is less than 4. These substances could therefore be used as antibacterial agents.

### **Conclusion**

This work made it possible to demonstrate the effectiveness and in vitro antibacterial properties of derivatives resulting from S-alkylation and Mannich-based substances on six bacterial strains (four (4) strains of *E. coli* and two (2) strains of enterobacteria) isolated from foods. The MICs obtained ranged between 31.25 and 125 µg/mL. As for the CMBs, they were between 31.25 and 250 µg/ML. CMB/MIC ratios indicated chemicals with bactericidal activity. These results made it possible to conclude that some of these synthesized chemical compounds have antibacterial properties. Thus, these molecules could contribute to the fight against bacterial infections.

### **References**

- Abdi R. D., Dego K. O. (2019). Antimicrobial activity of *Persicaria pensylvanica* extract against *Staphylococcus aureus*. *European Journal of Integrative Medicine*, 29 :1-7
- Antunes P., Novais C., Peixe L. (2020). Food-to-human bacterial transmission. *Microbiol Spectrum* 8 (1) : 1-26
- Attien P., Toe E., Kouassi K. A., Zébré A. C., Gomé M. N., Sina H., Dadie A. (2022). Evaluation of health risks related to the consumption of fish from the guéssabo river. *Food and Nutrition Sciences*, 13(1) : 55-64.
- Attien Y. P., Zebre A. C., Sina H., Angaman, D. M., Baba-Moussa L., Adjehi T. D. (2021). Assessment of the Sanitary Quality of the Dishes Sold in Street's Restaurants with High Frequentation in Daloa (Ivory Coast). *Int. J. Curr. Microbiol. App. Sci*, 10(12), 206-215.
- Bernard H., Faber M., Wilking H., Haller S., Höhle M., Schielke A., Ducomble T, Siffczyk C., Merbecks S. S., Fricke G., Hamouda O., Stark K., Werber D. (2014). Large multistate outbreak of norovirus gastroenteritis associated with frozen strawberries, Germany, 2012. *Eurosurveillance*, 19 (8)

- Bolou G.A. (2021). Etalement urbain et accès aux services socio-collectifs dans la ville de Daloa (Centre- Ouest de la Cote d'Ivoire). *Revue scientifique spécialisée en Géographie*, 5 : 26-46.
- Bssaibis, F., Gmira N., Meziane M. (2009). Activitéantibactérienne de *Dittrichiaviscosa* (L.) W. Greuter. *Revue de Microbiologie Industrielle, Santé et environnement*, 3: 44-55
- Cardinale E., Perrier Gros-Claude J.D., Tall F., Gueye E.F., Salvat G. (2005). Risk factorsforcontamination of ready-to-eatstreet-vendedpoultrydishes in Dakar, Senegal. *It. J. Food Microbiol*, 25 : 157-165
- Ferreira A., PetrettiC.,Vanina B. (2013). Les toxi-infections alimentaires collectives, *In* : Biologie de l'alimentation Humaine, France 477-493
- Fleckenstein J. F., Kuhlmann M., Sheikh A. P. (2021). Acute bacterialgastroenteritis. *Gastroenterol Clin North Am*, 50(2): 283-304
- Fung F., Huei-Shyong W., Suresh M. (2018). Food safety in the 21st century. *Biomedical Journal*, 41 (2) :88-95
- Gheorghe R. (2015). Mannich bases in medicinalchemistry and drugdesign.*European Journal of medicinalchemistry*, 89 : 743-816.
- INS.(2021). Recensement générale de la population et de l'habitat : Résultat globaux. *Ministère du plan et du Développement de Côte d'Ivoire*, Abidjan (Côte d'Ivoire) 37p
- Iredell J., Brown J., TaggK. (2016). Antibioticresistance in *Enterobacteriaceae*:mechanisms and clinical implications. *TheBmj*, 352 :1-19
- Kirk D. M., Pires M. Sara., Robert E. Black., Caipo M., Crump A. J., Devleesschauwer B., Döpfer D., Fazil A., Fischer-Walker C.L., Hald T., Hall J. A., Keddy H. K., Lake J. R., Lanata C. F., Torgerson P. R., Havelaar A. H., Angulo F. J. (2015). World HealthOrganizationEstimates of the Global and RegionalDiseaseBurden of 22 FoodborneBacterial, Protozoal, and Viral Diseases, 2010: A Data Synthesis. *PLoS Med*,12(12).
- Khomotso J. M., Chelule P. K. (2020). Safe Food Handling Knowledge and Practices of Street Food Vendors in Polokwane Central Business District. *Foods*. 9 (11):1-10
- Koffi-Nevry R., Assi C. B. J., Assemmand E. F., Wognin A. S., Koussémon M. (2012). Origine des témoins de contamination fécale de l'eau d'arrosage de la laitue (*lactuca sativa*) cultivée dans la zone péri urbaine d'Abidjan. *Journal of Applied Biosciences*, 52: 3669-3675
- Konate K.,Mavoungou J. F.,Lepengue A. N., Aworet-Samseny R. R., Hilou A., Souza A., Dicko M. H., M'Batchi B. (2012).Antibacterialactivityagainstbeta- lactamase

producing Methicillin and Ampicillin-resistant *Staphylococcus aureus*: fractional Inhibitory Concentration Index (FICI) determination, *Ann. Clin. Microbiol. Antimicrob.* 11(18) :1-12

Koné W. M., Atindehou K. K., Terreaux C., Hostettmann K. U. R. T., Traore D., Dosso M. (2004). Traditional medicine in North Côte-d'Ivoire: screening of 50 medicinal plants for antibacterial activity. *Journal of ethnopharmacology*, 93(1) : 43-49

Li Y., Wang Z., Ajani J. A., Song S. (2021). Drug resistance and Cancer stem cells. *Cell Communication and Signaling*, 19(1) :1-11

Ministère de la santé et de l'hygiène publique, Côte d'Ivoire. (2022). Intoxication alimentaire à Bondoukou. 4 Mai 2022. In: bondoukou, dépêches, santé/AIP

Mosupye M., Von Holy A. (2000). Microbiological hazard identification and exposure assessment of street food vending in Johannesburg, South Africa. *International Journal of Food Microbiology*, 61 (2000) 137-145

Nayaka, N.M.D.M.W., Sasadara, M.M.V., Sanjaya, D.A., Yuda, P.E.S.K., Dewi, N.L.K.A.A., Cahyaningsih, E., Hartati, R. (2021). Piper betle (L): Recent Review of Antibacterial and Antifungal Properties, Safety Profiles, and Commercial Applications. *Molécules*, 26 (2321) :1-21

Organisation Mondiale de la Santé (OMS). (1996). La restauration collective publication régionale, série européenne Genève, 71p

Ponce A. G., Fritz R., Del Vallec., Rouras I. (2003). Antimicrobial activity of essential oils on the native microflora of organic Swiss chard. *Society of Food Science and Technology*, 36(7): 679-684

Pons M.J., Mosquitto S., Gomes C., Del V. L.J., Ochoa T.J., Ruiz J. (2014). Analysis of quinolone-resistance in commensal and diarrheagenic *Escherichia coli* isolates from infants in Lima, Peru. *Trans R Soc Trop Med Hyg*, 108 :22-8

Popiołek L. (2017). Hydrazide-hydrazones as potential antimicrobial agents: overview of the literature since 2010. *Med Chem Res*, 26:287-301

Rahman M.M., Arif M.T., Bakar K., Talib Z. (2016). Food safety knowledge, attitude and hygiene practices among the street food vendors in Northern Kuching City, Sarawak. *Borneo Sci*, 31:95-103

Vieira K. C. de O., Silva, H. R. A. da, Rocha, I. P. M., Barboza, E., Eller, L. K. W. (2021). Foodborne pathogens in the omic era. *Critical Reviews in Food Science and Nutrition*, 62(24) :1-16

Zébré A. C., Yao P. A., Sina H, Gome M., Konate I., Baba-Moussa L. (2022). Sanitary Investigation on Microbial and Heavy Metals of Grilled Fish Sold in the Night Street Market of Daloa (Ivory Coast). *Journal of Advances in Microbiology*, 22 (12): 1-9

Zniber R., Ahmed M., Redouane A., Mostafa E. G., Abdelkrim F. M., Abdel I. M. Abdelkamel H. (2009). Activités biologiques de dérivés du benzimidazole. *Revue Roumaine de Chimie*, 54(8) : 643-650

UNDER PEER REVIEW