

FACTORS CONTRIBUTING TO THE PREVALENCE OF PRENATAL DEPRESSION AMONG ADOLESCENT MOTHERS SEEKING ANTE-NATAL CARE AT WAJIR COUNTY REFERRAL HOSPITAL

Abstract

This thesis investigates prenatal depression among adolescent mothers receiving antenatal care at Wajir County Referral Hospital. The study, utilizing questionnaires, reveals a significant prevalence of prenatal depression in this population. Factors such as age, education, and socioeconomic status contribute to the vulnerability of adolescent mothers. Psychosocial stressors like social stigma and lack of support networks also compound the risk. The research explores the experiences of adolescent mothers, emphasizing the need for tailored interventions and integrating mental health support into antenatal care services. The findings suggest implications for healthcare policies and community-based initiatives to address prenatal depression in resource-constrained settings like Wajir County, promoting better maternal and child outcomes.

Keywords: Prenatal depression, Adolescent mothers, Prevalence, Psychosocial stressors, antenatal care

1. Introduction

This text discusses the significance of prenatal depression among adolescent mothers, focusing on its prevalence and associated factors. The document highlights that depression during pregnancy can occur in various forms, affecting both the mother and child. It emphasizes the negative outcomes linked to depressive symptoms in adolescent mothers, such as decreased maternal confidence and increased likelihood of subsequent pregnancies. The global prevalence of prenatal depression is mentioned, with higher rates in Low- and Middle-Income Countries (LMICs). In the African context, cultural and social factors contribute to the mental health challenges faced by pregnant and postpartum adolescents.

The study's statement of the problem underscores the need to address prenatal depression among adolescents, considering factors like dysfunctional family structures, low socioeconomic status, and lack of support. The text justifies the research, citing the lack of studies on prenatal depression in Africa and the need for specific interventions tailored to the context of Wajir

County in Kenya. The objectives of the study include determining the prevalence of prenatal depression and identifying associated socio-cultural, maternal, and healthcare factors among adolescent mothers in Wajir County Referral Hospital.

The research questions outline specific inquiries, such as the prevalence of prenatal depression and the factors associated with it. The significance of the study is highlighted in addressing the increasing cases of prenatal depression among adolescent mothers in Wajir County Referral Hospital. The study aims to contribute valuable insights for healthcare professionals and policymakers to develop targeted interventions and policies.

Limitations and delimitations of the study are acknowledged, including the confined scope to Wajir County Referral Hospital, limiting generalizability, and the potential for reverse causality in establishing causation.

2. Literature Review

Depression is a prevalent mental disorder characterized by persistent sadness, lack of interest in enjoyable activities, and various physical symptoms. It has significant global implications, being a leading cause of disability. Antenatal depression in pregnant women can lead to adverse birth outcomes and postnatal depression, making routine screening crucial.

Various screening instruments, including the Edinburgh Postnatal Depression Scale (EPDS), Beck Depression Inventory (BDI), Mini-international Neuropsychiatric Interview (MINI-Plus), and Hamilton Depression Rating Scale (HAM-D), are used for assessing prenatal depression. The EPDS scale is widely employed and has demonstrated reliability, sensitivity, and specificity, particularly in resource-constrained settings.

The prevalence of prenatal depression among adolescent mothers tends to be higher than in other groups. Rates of depressive symptoms in HICs are around 10-15%, with variations across trimesters. In LMICs, a meta-analysis reported a prevalence of 15.6%, while studies in Africa found rates of 11.3% and 18.3% for prenatal and postnatal depression, respectively.

Several factors influence prenatal depression, including economic status, socio-demographic factors, cultural aspects, adverse life events, and healthcare-related factors. Low economic status, low education levels, and unemployment are associated with higher risks. Social support, marital satisfaction, and perceived support act as protective factors, while adverse life events and lack of healthcare access contribute to increased risks.

Commented [A1]: Repetition of ideas the similar with previous paragraph.
Write research question as question forms.

Commented [A2]: Not important

Commented [A3]: Add reference

Commented [A4]: Add reference

Commented [A5]: Add reference

For adolescent mothers, factors such as marital status, low socio-economic status, and lack of social support may contribute to higher rates of prenatal depression. Cultural factors, including societal views on single parenting, also play a role.

Finally, healthcare factors, including the quality of medical care and access to healthcare facilities, are significant predictors of prenatal depression. Factors such as distance to health facilities, inadequate staffing, and drug supplies can impact healthcare-seeking behavior among pregnant women, potentially contributing to prenatal depression.

3. Research Methodology

3.1 Research Design

The study utilized a hospital-based case-control study design to identify factors associated with prenatal depression. The choice of this design was driven by its effectiveness in investigating rare outcomes that might be overlooked in random sampling. While a population-based study would be ideal, the hospital-based design was chosen for its convenience in recruiting adolescent pregnant mothers, both cases and controls, who seek care at the antenatal clinic. To minimize potential differences in health-seeking behavior between hospital-based and population-based controls, controls were recruited from the same antenatal clinic as the cases.

Commented [A6]: You said above hospital based contradicting ideas.

3.2 Study Site

The study was conducted at the Wajir County Referral Hospital in Wajir County, Kenya. Serving the entire county, it is located near Wajir police station within Wajir East Constituency. The hospital, occupying about an acre, provides comprehensive outpatient and inpatient services, covering preventive, curative, and rehabilitative care. With six inpatient wards, it has a total bed capacity of 83, experiencing an average annual inpatient turnover of 8341 (Wajir Hospital Statistics, 2016). The annual outpatient workload is around 6,700, catering to both adults and children.

The hospital offers Essential Maternal Child Health (MCH) services, including antenatal clinic services (ANC), family planning, child welfare, immunization, prevention of mother-to-child transmission of HIV (PMTCT), and health education. ANC clinic services encompass health education, promotion of healthy behavior, physical examination of pregnant women, management of pregnancy-related complications, treatment of concurrent illnesses, PMTCT, reproductive health consultations, and preventive measures such as iron and folic acid

supplementation, deworming, tetanus toxoid immunization, distribution of long-lasting insecticidal bed nets (LLIN), and intermittent preventive treatment of malaria (IPTp).

3.3 Study Population

The study population will comprise of adolescent pregnant women from the age 12 years to 19 years routinely attending the antenatal clinic at the Wajir County Referral Hospital during the data collection period of three months. Pregnant adolescent mothers are regarded as emancipated minors in Kenya and can legally give consent. Cases and controls will be selected from this population.

Commented [A7]: Clearly define population for case and for control

3.4 Sample Size Determination

Sample size calculation reduces the cost and time for research studies while at the same time ensuring representativeness. Since this was a controlled study and the main variable for analysis is the Odds Ratio (OR), 159 cases and 115 controls. The sample size was estimated by Methods in Observational Epidemiology 2nd Edition as illustrated by Kelsey et al. 1996.

Commented [A8]: Pls use sample size determination formula for case control study. What is OR for you? OR is used to compare between the two group

3.5 Data Collection, Processing and Analysis

The study recruited participants through two hospital-based research assistants who underwent training on study content, EPDS scale administration, ethical considerations, and questionnaire filling. Data collection occurred in the morning, aligning with the typical ANC clinic hours (9 am to 1 pm). Pregnancy confirmation involved checking the ANC booklet for evidence like a positive PDT test or ultrasound. Research assistants explained eligibility, study purpose, and procedures before obtaining consent. Participating mothers, upon voluntary consent, underwent EPDS administration to determine depression status. Subsequently, a predetermined questionnaire was administered to both cases and controls, with a maximum of two controls for each case.

For data processing and analysis, information collected via electronic tablets was checked for accuracy and double entries. The dataset underwent cleaning and analysis using Stata software version 13.0. Descriptive statistics, such as means and medians, summarized continuous variables, while proportions and percentages provided descriptive data for categorical variables. Odds ratios were determined, using a reference to describe and explain the relationship between variables. Non-significant variables in the case-control study were excluded from the model if their removal didn't cause over a 30% change in the regression coefficients of the remaining variables (Mirieri, 2019). A case-control two-way interaction was fitted in the final model, with

significance assessed. The Hosmer-Lemeshow goodness of fit tested how well the model fitted the case-control data, with a p-value > 0.05 indicating a good fit.

Commented [A9]: This is used to check the goodness of fit of the model for logistic regression. But, you didn't say anything about logistic model

4. Findings

4.1 Respondents Demographics Characteristics




Figure 1 Age Distribution Between Cases of the Respondent and Control Graph

The figure above shows that majority of the respondents are aged between 19-21 years 29%, 15-18 years 24%, and 22-24 years 17%. This implies that most pregnant women are above the consent age of 18.'

4.2 Prevalence of Prenatal Depression among Adolescent Mothers

The overall Prevalence was

$$Prevalence = \frac{\text{Total Number of depressed women}}{\text{Total number of screened women}} \times 100\%$$

$$\frac{53}{159} \times 100\%$$

$$33.3\%$$

4.3 Maternal Factors Associated Prenatal Depression among teenage Pregnant Mothers

Table 1 The Association Pregnancy and Effects with Prenatal depression

Exposure pregnancy complication	Outcome (Dependent variables)			OR
	Cases (Depression)	Non-Cases (No Depression)	Total	
Yes	40	36	76	Reference
No	13	70	83	4.7
Total	53	106	159	

From the above, the OR is 6. This means that an adolescent woman with complications are 6 times more likely to have prenatal depression as compared to an adolescent with no obstetric complication.

4.4 Association of Health Care Factors with Prenatal Depression amongst teenage Pregnant Mothers

This section seeks to answer the fourth research question: What are the health care systems associated with burden of prenatal depression among adolescent mothers attending the antenatal clinic at Wajir County referral hospital, Kenya?

The results are shown in the table below:

Table 2 Association of Health Care Factors with Prenatal Depression amongst teenage Pregnant Mothers

Health Care Factors Causing Prenatal Depression amongst teenage Pregnant Mothers					
Factor	SA	A	N	D	SD
There is poor government provision and utilization of antenatal care services	89(56%)	41	14	12	3
Increasing number of adolescent depression and mental health burden	73(46%)	63	14	8	1
Staff are not friendly when it comes to handling adolescent pregnant women	58(36%)	55	35	9	2
Facility is far from her home so is it a challenge to access basic health services	59(37%)	53	32	14	1
Poor security is on the way demoralizes search for hospital-based care	46	59(37%)	35	14	5
Insufficient personnel to sensitize on emotional, psychosocial, health, and educational problems in the lives of vulnerable pregnant adolescents	78	60(47%)	9	10	2

From the Table 2 it can be established that majority of the respondents agree that there is poor government provision and utilization of antenatal care services (strongly agree (56%) and agree (26%). Most of the respondents also agree that there is an increasing number of adolescent depression and mental health burden in the community. The respondents agree that medical staff are not friendly when it comes to handling adolescent pregnant women. Majority of the respondents strongly agree that the health facility is far from their homes making it a challenge to access basic health services. In the community most of the respondents attribute poor security on

the way demoralized search for hospital-based care also they agree that insufficient personnel to sensitize on emotional, psychosocial, health, and educational problems in the lives of vulnerable pregnant adolescents in Wajir County hospital.

5. Discussion, Conclusion, and Recommendation

5.1 Discussion

Objective i: Prevalence of Prenatal Depression The study revealed that 33.3% of adolescent mothers attending the antenatal care clinic at Wajir County Referral Hospital experienced prenatal depression. This highlights the substantial issue, underscoring the necessity for targeted interventions and support systems for this vulnerable population.

Objective ii: Socio-cultural Factors Socio-cultural factors significantly contribute to prenatal depression among adolescent mothers, with traditional norms, social stigma, and lack of social support playing pivotal roles. The findings emphasize the importance of culturally sensitive interventions and community-based programs to address and alleviate these factors.

Objective iii: Maternal Factors Maternal factors, including age, educational level, marital status, and financial status, emerged as strong predictors of prenatal depression. The study recommends targeted interventions focusing on improving education, economic empowerment, and social support structures for adolescent mothers to mitigate the impact of these factors.

Objective iv: Health Care Factors Health care-related factors were identified as influencers of prenatal depression prevalence among adolescent mothers. Barriers to accessing quality antenatal care services, lack of mental health screening, and limited awareness were noted contributors. The study suggests integrating mental health screening into routine antenatal care and enhancing healthcare accessibility as crucial measures to address these factors.

5.2 Conclusions

- I. The study highlighted a notable prevalence of prenatal depression among adolescent mothers in the antenatal care clinic at Wajir County Referral Hospital. It underscores the importance of routine screening for prenatal depression to enable early identification and intervention.
- II. Critical socio-cultural factors contributing to prenatal depression include social stigma, cultural norms, and a lack of support networks. The study emphasizes the necessity of

community-based awareness campaigns and support initiatives to address these challenges effectively.

- III. Maternal factors, such as age, educational background, and socioeconomic status, were identified as determinants of prenatal depression. The study recommends targeted interventions and educational programs within the antenatal care setting, specifically tailored for vulnerable groups.
- IV. Health care factors, particularly the quality of antenatal care services and the integration of mental health support were found to significantly influence prenatal depression. The study suggests enhancing mental health training for healthcare providers and advocating for improved resources in the antenatal care setting to address these factors adequately.

5.3 Recommendations

In response to the findings on the prevalence of prenatal depression among adolescent mothers in antenatal care at Wajir County Referral Hospital, the following recommendations are proposed to address socio-cultural, maternal, and healthcare factors influencing mental well-being:

- i. Implement routine screening protocols for prenatal depression during antenatal care visits. Integrate mental health assessments into existing ANC services to identify and address symptoms early on.
- ii. Develop community-based awareness campaigns to combat the social stigma surrounding adolescent pregnancy. Establish support networks within the community to enhance the emotional well-being of adolescent mothers.
- iii. Initiate targeted interventions for younger mothers and those with lower educational backgrounds. Implement educational programs within the ANC setting to empower adolescent mothers with coping strategies and resources.
- iv. Enhance mental health training for healthcare providers involved in antenatal care. Advocate for the integration of mental health services within the ANC setting, ensuring that adequate resources and support are available for adolescent mothers.

References

- Abuidhail, J., & Abujilban, S. (2014). Characteristics of Jordanian depressed pregnant women: A comparison study: Antenatal depression. *J. Psychiatr. Ment. Health Nurs.* 21, , 573-579.
- Adewuya, A., Ola, B., Aloba, O., Dada, A., & Fasoto, O. (2017). Prevalence and correlates of depression in late pregnancy among Nigerian women. *Depress. Anxiety* 24, 15-21.
- Ajinkya, S. J. (2013a). Depression during pregnancy: Prevalence and obstetric risk factors among pregnant women attending a tertiary care hospital in Navi Mumbai. *Ind. Psychiatry J.* 22, 37-40.
- Ali, N., Azam, I., Ali, B., Tabbusum, G., & Moin, S. (2012). Frequency and Associated Factors for Anxiety and Depression in Pregnant Women: A Hospital-Based Cross-sectional study. *Scientific World Journal* 2012.
- Amorim, P. (2000). Mini International Neuropsychiatric Interview (MINI) . *Rev. Bras. Psiquiatr* 22, 106-115.
- Babu, G., Murthy, G., Singh, N., Nath, A., Rathnaiah, M., Saldanha, N., . . . Kinra, S. (2018). Socio demographic and Medical Risk Factors Associated with Antepartum Depression. *Front. Public Health* 6.
- Bawahab, J., Alahmadi, J., & Ibrahim, A. (2017). Prevalence and determinants of antenatal depression among women attending primary health care centers in Western Saudi Arabia. *Saudi Med. J.* 38, 1237-1242.
- Beck, A. T., Ward, C., Mendelson, M., Mock, J., & Erbaugh, J. (1961). An inventory for Measuring Depression. *Arch. Gen. Psychiatry* 4, , 561-571.
- Bennet, H. A., Einarson, A., Taddio, A., Koren, G., & Einarson, T. R. (2004a). Prevalence of depression during pregnancy: systematic review. *Obstet. Gynecol.* 103, , 698-709.
- Bhat, N. A., Hassam, R., Shafiq, M., & Sheikh, S. (2015). Sociodemographic factors: A major predictor of anxiety and depression among pregnant women., *Delhi psychiatry Journal* 2015; 18(1).
- Biaggi, A., Conroy, S., Pawlby, S., & Pariante, C. (2016a). Identifying the women at risk of antenatal anxiety and depression: Asystemic review. *J. Affect. Disord.* 191, 62-77.
- Biratu, A., & Haile, D. (2015). Prevalence of antenatal depression and associated factors among pregnant women in Addis Ababa, Ethiopia: a cross-sectional study. *Reprod. Health* 12, 99.

- Bodecs, T., Szilagyai, E., Cholnoky, P., Sandor, J., Gonda, X., Rihmer, Z., & Horvath, B. (2013). Prevalence and psychosocial background of anxiety and depression emerging during the first trimester of pregnancy: data from a Hungarian population-based sample. *Psychiatric Danud*, 25, 352-358.
- Bonari, L., Pinto, N., Ahn, E., Einarson, A., Steiner, M., & Koren, G. (2004). Perinatal risks of untreated depression during pregnancy. *J. Psychiatry* 49, 726-735.
- Brittain, K., Myler, L., Koen, N., Koopowitz, S., A., D. K., Barnett, W., . . . Stein, D. (2015). Risk Factors for antenatal Depression and associations with infant Birth Outcomes: Results from a South African Birth Cohort Study. *Paediatr.Perinat. Epidemiol*, 29, 504-514.
- Castro e Cousto, T., Martins Brancaglioni, M. Y., Nogueira Cardoso, M., Bergo Protzner, A., Duarte Garcia, F., Nicolato, R., . . . Correa, H. (2015). What is the best tool for screening antenatal depression. *J. Affect. Disord*. 178, 120-17.
- Chorwe-Sungani, G., & Chipps, J. (2017, Mar 24). A systematic Review of screening Instruments for Depression for use in antenatal services in low resource settings. *BMC Psychiatry*, p. 1.
- CIDP. (2018). *Wajir County Referral and Teaching Hospital Statistics*. Wajir: Unpublished data.
- Coll, C., de, V., da Silveira, M., Bassani, D. G., Netsi, E., Wehmeister, F. C., . . . Stein, A. (2017). Antenatal depressive symptoms among pregnant women: Evidence from a Southern Brazilian population-based cohort study. *J. Affect. Disord*. 209, 140-146.
- Collin, P., Insel, T., Chockalingam, A., Daar, A., & Maddox, Y. (2013). Grand Challenges in Global Mental Health: Integration in Research, Policy, and Practice. . *PLOS Med*. 10 e1001434.
- Cox, J., Holden, J., & Sagovsky, R. (1987). Detection of postnatal Depression. Development of the 10-item Edinburgh Postnatal Depression Scale. *Br. J. Psychiatry* 150, 782-786.
- Dadi, F. A., Miller, E. R., Bisetegn, T., & Mwanri, L. (2020, February 04). Global Burden of antenatal Depression and its association with adverse birth outcomes: an umbrella review. *BMC Public Health* 20, Article number 173 (2020).
- Dias, C., & Figueiredo, B. (2015). Breastfeeding and depression: a systemic review of the literature. *J. Affect. Disord*. 171, 142-154.