

Effect of long-term continuous Tea cultivation in water quality index under Deep, fine loamy, very deep, coarse loamy and very deep, fine loamy well drained soil

Abstract

For this investigation, three different types of soil were chosen: very deep, coarse loamy, well-drained soil; deep, fine loamy, well-drained soil; and deep, coarse loamy, well-drained soil. Ten water samples were taken from the ring well, ponds, drains, and existing shallow tube wells (hand and mark tube wells) for each category of soil types. The minimal data set, which include all variables, was chosen on the basis of the significant correlation discovered by correlation analysis. Very deep, coarse loamy, well-drained soil had the lowest water quality index (10.34), whereas very deep, fine loamy, well-drained soil had the highest (15.38).

Keywords: water quality index, agricultural activity, aquatic biodiversity, urbanization

Introduction:

The impact of agriculture on surface and ground water quality as well as aquatic biodiversity is a growing concern for scientists, consumers, and even the agricultural community (Katak et al., 1994). Tea cultivation is a major agricultural activity in Assam, N.E India that provide livelihood to a large number of people. However, the tea business also uses a lot of pesticides and fertilizers, which end up being washed into streams and other freshwater systems by surface runoff. This harms the aquatic biodiversity and water quality. In Malaysia, it has been discovered that increased agricultural activities, notably the growing of tea, causes nitrates to seep into surface and ground water. In tea gardens, a lot of agrochemicals are used to increase its productivity. Consequently, contamination of the soil and water becomes obvious. Thus, it is crucial to keep an eye on the sources of drinking water in the tea gardens region. (Joydev Dutta, 2020)

One of the best techniques for informing concerned individuals and decision-makers about the condition of the water is the water quality index. The World Health Organization estimates that water is the primary cause of around 80% of human ailments. It is not possible to restore the quality of groundwater once contamination has occurred by removing the pollutants

from their source [9-11]. As a result, it becomes crucial to constantly check the quality of groundwater and devise strategies and tactics to safeguard it. Consequently, it turns into a crucial factor in the evaluation and maintenance of groundwater (Ramakrishnaiah et al., 2008). A grade that reflects the combined influence of several water quality measures is known as the Water Quality Index. It is estimated concerning groundwater's appropriateness for human consumption. By normalizing values to arbitrary rating curves, it is a dimensionless number that integrates several water quality parameters into a single number (Miller et al., 1986). WQI factors are subject to change based on local preferences and the water body's defined usage. Dissolved oxygen (DO), pH, chemical and biological oxygen demands (BOD and COD), temperature, bacteria, and nutrients (nitrogen and phosphorus) are a few of the variables. These parameters have various ranges of measurement and units of expression. These parameters have various ranges of measurement and units of expression. Because of population growth, industrialization, and urbanization, there is an enormous increase in the demand for fresh water, which is measured by the Water Quality Index (WQI), which synthesizes complicated scientific data about these variables into a single number. Additionally, a major issue facing the tea sector is the degradation of water quality brought on by the prudent use of pesticides and agrochemicals in the gardens.

Material and Method:

For this investigation, three different types of soil were chosen: very deep, coarse loamy, well-drained soil; deep, fine loamy, well-drained soil; and deep, coarse loamy, well-drained soil. Ten water samples were taken from the ring well, ponds, drains, and existing shallow tube wells (hand and mark tube wells) for each category of soil types. The substantial association found by correlation analysis was used to pick the minimum data set, which represents all of the variables (Andrews et al., 2002a). For every statistically significant variable, standardized Principal Component Analysis (PCA) was carried out using SPSS-15 software, and component-wise MDS were chosen in the following stage. It was considered that variables with high factor loading and principal components with high eigen values better described the qualities of the system. The principal components that best describe the system were chosen based on the breaking point, which is where the steady reduction in eigen values appears to uniform towards the right part of

the graphical principal components. We will only look at PCs that explained at least 10% of the variation in the data (Wander and Bollero, 1999) and those with eigenvalues >1 (Brejda et al., 2000). Only highly weighted factors were kept for MDS inside each PC. Absolute values within 10% of the highest factor loading were considered highly weighted factor loadings. Multivariate correlation matrices were also utilized to assess the strength of the correlations between the variables in order to eliminate duplication and rule out erroneous groupings among the highly weighted variables inside PCs. Only one well-correlated variable was taken into account for the MDS since other variables considered superfluous. The remaining ones were removed from the dataset. Each highly weighted variable was considered significant and kept in the MDS if there was no correlation between them. Following the identification of the MDS indicators, a linear scoring approach was applied to each observation of the MDS indicators (Andrews et al., 2002a). The order of the indicators was determined by categorizing a greater value as "good" or "bad" for soil function. Each observation was divided by the greatest observed value for "more is better" indications, resulting in a score of 1 for the highest observed value. In the case of "less is better" indicators, each observation (in the denominator) was divided by the lowest observed value (in the numerator) to give the lowest observed value a score of 1. Following transformation, the PCA findings were used to weight each observation's MDS variables. A specific percentage (%) of the variation in the entire data set was explained by each PC.

$$WQI = \sum_{i=1}^n W_i.S_i$$

where W_i is the weighted factor obtained from the PCA and S_i is the subscripted variable's score. Here, it was assumed that higher index scores corresponded to either improved soil function or better soil quality (Chaudhury, 2005).

Result and Discussion:

The correlation matrix among the observed variables was examined in order to evaluate the compatibility of correlation values with the hypothesized factor structure. Four out of the thirteen pairs of water properties for the deep, fine loamy, well-drained soils in the Jorhat district showed significant correlations ($P < 0.01$). There were notable positive associations found for pH and HCO_3^{2-} (0.755**), TDC (1.00**), and Ca (0.897**) in EC. Ca showed significant negative

correlations with SO_4^{2-} (-0.836**). Using the Cattell scree test, the 13 water quality attributes that were taken into consideration for the principal component analysis were classified into components (Table 1) for the purpose of grouping water quality indicators. The components were plotted as the X-axis and the associated eigen values as the Y-axis (Fig. 1). The first three main components were kept for interpretation since they had eigen values >1 and explained 72.20% of the variance in the whole data set. The six components described more than 90% of the variation in EC, TDS, and Ca; more than 80% in F; more than 70% in pH, Mg, CO_3^{2-} , HCO_3^{2-} , NO_3^- , SO_4^{2-} and more than 60% in PO_4^{3-} ; more than 30% in Cl, and more than 10% in As, according to the communities for the soil parameters. 31.76 percent of the variation was explained by the first principal component. It loaded positively for TDS (0.942), Ca (0.967), and EC (0.942). Based on the absolute values within 10% of the highest factor loading, these soil parameters were selected. The second major component demonstrated positive loading for pH (0.813), CO_3^{2-} (0.848), and HCO_3^{2-} (0.8850), accounting for 24.81 percent of the variance. The third principal component exhibited a positive loading for F (0.904) and explained 15.63% of the variance. Just 3 PCs with an eigen value greater than one were identified by a PCA of 13 variables, which accounted for 72.20% of the variance in the total data set. Separate runs of a correlation matrix were conducted for the highly weighted variables under various PCs (Table 2). F was kept for MDS in PC2, pH, and PC3, while EC and TDS were retained in PC1. Four variables in total—EC, TDS, pH, and F—were selected as MDS. The MDS variables were transformed by using scoring functions. The WQI was calculated by using weighing factors for each scored MDS variable according to the formula:

$$\text{WQI} = \Sigma(0.439 \text{ EC} + 0.439 \text{ TDS} + 0.344 \text{ pH} + 0.216 \text{ F}) = 14.44$$

If all four MDS were considered to be responsible for contributing ideal (100%) water quality index for deep, fine loamy, well drained soil then the relative soil quality explained would be 42.47% for TDS, 23.9% for EC, 22.23% for pH and 11.37% F for MWD (Fig. 2).

Table 1. Rotated Component loadings and communalities of water properties in deep, fine loamy, well drained soil of Jorhat district, Assam

| Water properties | Component | | | Communalities |
|------------------|-----------|-------|--------|---------------|
| | PC1 | PC2 | PC3 | |
| pH | 0.100 | 0.813 | -0.270 | 0.744 |

| | | | | |
|-------------------------------|--------|--------|--------|-------|
| EC | 0.942 | 0.178 | 0.114 | 0.932 |
| TDS | 0.942 | 0.178 | 0.114 | 0.932 |
| Ca | 0.967 | -0.047 | -0.054 | 0.939 |
| Mg | 0.733 | -0.376 | -0.345 | 0.798 |
| F- | 0.049 | 0.068 | 0.904 | 0.824 |
| Cl- | -0.267 | -0.371 | -0.396 | 0.366 |
| CO ₃ ²⁻ | -0.024 | 0.848 | 0.193 | 0.757 |
| HCO ₃ ⁻ | 0.022 | 0.850 | 0.109 | 0.734 |
| PO ₄ ³⁻ | 0.230 | -0.264 | 0.708 | 0.624 |
| NO ₃ ⁻ | -0.072 | -0.740 | 0.459 | 0.764 |
| SO ₄ ²⁻ | -0.779 | 0.339 | 0.255 | 0.787 |
| As | 0.365 | 0.202 | 0.108 | 0.186 |
| Eigan values | 4.130 | 3.225 | 2.032 | Total |
| % variance | 31.76 | 24.81 | 15.63 | 72.20 |

Table .2. Correlation matrix for highly weighted variables of water analysis under PC's with deep, fine loamy, well drained soil of Jorhat district, Assam

| PC1 | EC | TDS | Ca |
|-------------------------------|---------|-------------------------------|-------------------------------|
| EC | 1 | 1.000** | 0.897** |
| TDS | 1.000** | 1 | 0.897** |
| Ca | 0.897** | 0.897** | 1 |
| Correlation sum | 2.897 | 2.897 | 2.794 |
| PC2 | pH | CO ₃ ²⁻ | HCO ₃ ⁻ |
| pH | 1 | 0.516 | 0.755** |
| CO ₃ ²⁻ | 0.516 | 1 | 0.585* |
| HCO ₃ ⁻ | 0.755** | 0.585* | 1 |
| Correlation sum | 2.271 | 2.101 | 2.24 |
| PC3 | F | | |
| F- | 1 | | |

**Correlation is significant at the 0.01 level

*Correlation is significant at the 0.05 level




FIG: 1: SCREE PLOT FOR SELECTING PRINCIPAL COMPONENTS UNDER DEEP, FINE LOAMY, WELL DRAINED SOIL




FIG: 2: WQI UNDER DEEP, FINE LOAMY, WELL DRAINED SOIL

Water quality under very deep, fine loamy, well drained soils

The correlation matrix among the observed variables was examined in order to evaluate the compatibility of correlation values with the proposed factor structure. Four out of the thirteen pairs of water properties for the deep, fine loamy, well-drained soils in the Jorhat district showed a significant connection ($P < 0.01$). Significant positive correlations were observed for pH with

CO_3^{2-} and HCO_3^{2-} (0.0.826** and 0.849**) and EC with TDC (1.00**) and Ca (0.913**); CO_3^{2-} with SO_4^{2-} (0.746**).

The 13 water quality attributes taken into account in the main component analysis were divided into components (Table 3) for the purpose of grouping the soil quality indicators. The components were plotted as the X-axis and the associated eigen values as the Y-axis (Fig. 3) using the Cattelscree test. Retaining the first three principal components for interpretation, they accounted for 74.91 percent of the total variance in the entire data set and had eigen values >1. Communalities for the soil properties indicate that the six components explained >90% of the variance in pH, EC, TDS and Ca; >80% in Cl^- and CO_3^{2-} , >70% in HCO_3^- , PO_4^{3-} , SO_4^{2-} , >60% in As, > 50% in F and NO_3^- and >20% in Mg. The greatest negative loading was observed for pH (-0.815) and HCO_3^- (-0.815). Based on the absolute values within 10% of the highest factor loading, these soil parameters were chosen. Positive loading for SO_4^{2-} (0.779), As (0.756), and 18.48 percent of the variation was explained by the second principal component. The third principal component had a positive loading for HCO_3^- (0.656) and explained 14.99% of the variation. Only three PCs with eigen values > 1 were retrieved by a PCA of 23 variables, which accounted for 74.19 percent of the variance in the total data set. A correlation matrix for the highly weighted variables under different PCs was run separately (Table 3). In PC1, Ca; in PC2, pH and SO_4^{2-} ; in PC3, HCO_3^- were chosen as MDS. The MDS variables were transformed by using scoring functions. The WQI was calculated by using weighing factors for each scored MDS variable according to the formula:

$$\text{WQI} = \Sigma(0.437 \text{ Ca} + 0.372 \text{ pH} + 0.372 \text{ SO}_4^{2-} + 0.192 \text{ HCO}_3^-) = 10.34$$

If all four MDS were considered to be responsible for contributing ideal (100%) water quality index for continuous tea cultivation, then the relative water quality explained would be 32.94% for pH, 28.22% for Ca, 23.47% for SO_4^{2-} and 15.35% for HCO_3^- for MWD (Fig. 6).

Table 3. Rotated component loadings and communalities of water properties in very deep, fine loamy, well drained soil of Jorhat district, Assam

| Soil properties | Component | | | Communalities |
|-----------------|-----------|--------|--------|---------------|
| | PC1 | PC2 | PC3 | |
| pH | -0.815 | 0.066 | 0.484 | 0.902 |
| EC | 0.874 | -0.019 | 0.382 | 0.910 |
| TDS | 0.874 | -0.019 | 0.383 | 0.910 |
| Ca | 0.848 | 0.109 | 0.429 | 0.915 |
| Mg | 0.239 | -0.462 | 0.146 | 0.292 |
| F- | -0.507 | -0.314 | -0.485 | 0.590 |
| Cl- | 0.877 | 0.321 | -0.145 | 0.894 |
| CO32- | -0.815 | 0.312 | 0.226 | 0.813 |
| HCO3- | -0.694 | 0.070 | -0.656 | 0.791 |
| PO43- | 0.289 | 0.464 | 0.552 | 0.730 |
| NO3- | 0.152 | 0.688 | 0.251 | 0.559 |
| SO42- | -0.373 | 0.779 | 0.175 | 0.776 |
| As | 0.026 | 0.756 | -0.285 | 0.654 |
| Eigan values | 5.387 | 2.402 | 1.950 | Total |
| % variance | 41.44 | 18.48 | 14.99 | 74.91 |

Table 4. Correlation matrix for highly weighted variables of water analysis under PC's with very deep, fine loamy, well drained soil of Jorhat district, Assam

| | | | | | | |
|-----------------|-----------------|-------------------------------|---------|---------|---------|---------|
| PC1 | Cl ⁻ | CO ₃ ²⁻ | Ca | TDS | EC | pH |
| Cl ⁻ | 1 | -0.643* | 0.712* | 0.664* | 0.664* | -0.690* |
| CO32- | -0.643* | 1 | -0.596* | -0.553* | -0.553* | 0.826** |
| Ca | 0.712* | -0.596* | 1 | 0.913** | 0.913** | -0.467 |
| TDS | 0.664* | -0.553* | 0.913** | 1 | 1.000** | -0.542 |
| EC | 0.664* | -0.553* | 0.913** | 1.000** | 1 | -0.542 |
| pH | -0.690* | 0.826** | -0.467 | -0.542 | -0.542 | 1 |
| Correlation sum | 1.707 | -0.519 | 2.475 | 2.482 | 2.482 | -0.415 |
| PC2 | SO42- | As | | | | |
| SO42- | 1 | 0.327 | | | | |
| PC3 | HCO3- | | | | | |
| HCO3- | 1 | | | | | |

**Correlation is significant at the 0.01 level

*Correlation is significant at the 0.05 level




FIG: 3: SCREE PLOT FOR SELECTING PRINCIPAL COMPONENTS UNDER VERY DEEP, FINE LOAMY, WELL DRAINED SOIL




FIG: 4: WQI UNDER VERY DEEP, FINE LOAMY, WELL DRAINED SOIL

Water quality under very deep, coarse loamy, well drained soils

The correlation matrix among the observed variables was examined in order to evaluate the consistency of correlation values with the proposed factor structure. Four out of the thirteen pairs of water properties for the very deep, coarse loamy, well-drained soils in the Jorhat district showed a significant connection ($P < 0.01$). pH and CO_3^{2-} and HCO_3^{2-} (0.781** and 0.786**), EC and TDC (1.00**) and Ca (0.876**), and TDS and Ca (0.876) showed significant positive correlation. Mg and PO_4^{3-} showed significant negative correlations (-0.732**).

Grouping of Water Quality Indicators

Using the Cattelscree test, the 13 water quality parameters taken into account for the Principal component analysis were divided into components (Table 5). The components were plotted as the X-axis and the associated eigen values as the Y-axis (Fig. 5). The first three main components were kept for interpretation since they had eigen values greater than 1 and explained 75.15% of the variance in the whole data set. The six components accounted for over 90% of the variation in EC, TDS, Ca, Mg, and HCO_3^- ; >80% in pH and PO_4^{3-} ; >70% in CO_3^{2-} ; >60% in SO_4^{2-} ; >50% in NO_3^- , As; and >40% in F and Cl, according to the communities for the soil parameters. 32.82 percent of the variation was explained by the first principal component. For Ca, it demonstrated positive loading (0.939). Based on the absolute values within 10% of the highest factor loading, these soil parameters were chosen. The second principal component, which showed positive loading for SO_4^{2-} (0.679) and pH (0.674), explained 27.93 percent of the variation. The third main component, which had the maximum negative loading (-0.724) for magnesium and a positive loading (0.656) for HCO_3^- explained 14.39% of the variation. Only three PCs with eigen values > 1 were retrieved using a PCA of 23 variables, which accounted for 75.15% of the variance in the total data set. Separate runs of a correlation matrix under various PC settings were conducted for the highly weighted variables (Table 6). In PC1, Ca; in PC2, pH and SO_4^{2-} ; in PC3, HCO_3^- were chosen as MDS. The MDS variables were transformed by using scoring functions. The WQI was calculated by using weighing factors for each scored MDS variable according to the formula:

$$\text{WQI} = \Sigma(0.437 \text{ Ca} + 0.372 \text{ pH} + 0.372 \text{ SO}_4^{2-} + 0.192 \text{ HCO}_3^-) = 10.34$$

If all four MDS were considered to be responsible for contributing ideal (100%) water quality index for continuous tea cultivation, then the relative water quality explained would be 32.94% for pH, 28.22% for Ca, 23.47% for SO_4^{2-} and 15.35% for HCO_3^- for MWD (Fig. 6).

Table 5. Rotated component loadings and communalities of water properties in very deep, coarse loamy, well drained soil of Jorhat district, Assam

| Soil properties | Component | | | Communalities |
|-------------------------------|-----------|--------|--------|---------------|
| | PC1 | PC2 | PC3 | |
| pH | -0.431 | 0.674 | 0.412 | 0.809 |
| EC | 0.776 | 0.392 | 0.421 | 0.934 |
| TDS | 0.776 | 0.392 | 0.421 | 0.934 |
| Ca | 0.939 | 0.059 | 0.209 | 0.930 |
| Mg | 0.521 | 0.347 | -0.724 | 0.917 |
| F ⁻ | 0.068 | 0.529 | -0.426 | 0.466 |
| Cl ⁻ | 0.171 | -0.683 | -0.003 | 0.496 |
| CO ₃ ²⁻ | -0.686 | 0.522 | 0.096 | 0.753 |
| HCO ₃ ⁻ | -0.594 | 0.546 | 0.656 | 0.938 |
| PO ₄ ³⁻ | -0.191 | -0.422 | 0.412 | 0.881 |
| NO ₃ ⁻ | 0.718 | -0.034 | 0.052 | 0.519 |
| SO ₄ ²⁻ | 0.412 | 0.679 | 0.063 | 0.634 |
| As | 0.379 | -0.539 | 0.353 | 0.559 |
| Eigen values | 4.267 | 3.631 | 1.871 | total |
| % variance | 32.82 | 27.93 | 14.39 | 75.15 |

Table 6. Correlation matrix for highly weighted variables under PC's with very deep, course loamy, well drained soil of Jorhat district, Assam

| | | | |
|-------------------------------|---------|-------------------------------|-------------------------------|
| PC1 | Ca | | |
| Ca | 1 | | |
| PC2 | pH | Cl ⁻ | SO ₄ ²⁻ |
| pH | 1 | -0.690* | 0.410 |
| Cl ⁻ | -0.690* | 1 | -0.143 |
| SO ₄ ²⁻ | 0.410 | -0.143 | 1 |
| Correlation sum | 0.72 | 0.167 | 1.267 |
| PC3 | Mg | HCO ₃ ⁻ | |
| Mg | 1 | -0.109 | |
| HCO ₃ ⁻ | -0.109 | 1 | |

**Correlation is significant at the 0.01 level

*Correlation is significant at the 0.05 level




Fig: 5: SCREE PLOT FOR SELECTING PRINCIPAL COMPONENTS UNDER DEEP, COARSE LOAMY, WELL DRAINED SOIL




Fig:6.: WQI UNDER DEEP, COARSE LOAMY, WELL DRAINED SOIL

Discussion:

In deep, fine-loamy, well-drained soil, the water quality index was found to be 14.44. Since there is a strong correlation between Ca and Ec, EC and TDS were kept for MDS in PC1, while Ca was removed. Because CO₃²⁻ and HCO₃⁻ were linked with pH, they were removed from MDS in PC2, while pH was kept in place. F⁻ was kept for PC3 in the MDS. Ultimately, for MDS, four variables—EC, TDS, pH, and F⁻—were kept. We can conclude that TDS is the primary indicator of water quality if all four MDS are thought to be important for contributing to the ideal (100%) water quality index. The leaching of ions from agricultural fields and the soils and rocks of tea

gardens could be the cause of the high TDS value. TDS shows how salinity behaves in water. According to Dutta *et al.* (2010), it is typically the primary component that restricts or determines the use of ground water for any purpose. According to Dutta *et al.* (2010), in small tea estates in the Sonitpur district, almost 13% of the water samples had TDS levels over the allowable limit of 500 ppm. TDS content is usually the main factor, which limits or determines the use of ground water for any purpose. Dutta *et al.* (2010) found that more than 13 per cent of the water samples were exhibited TDS values outside the permissible limit 500 ppm in small tea gardens of Sonitpur district. In PC1, Ca was retained for MDS. In PC2, pH and SO_4^{2-} were retained and Cl^- was dropped from MDS as it showed the lowest correlation sum. In PC3, HCO_3^- . Finally, Ca, pH, SO_4^{2-} and HCO_3^{2-} were retained for MDS. Finally four parameters viz., Ca^{2+} , pH, SO_4^{2-} were retained for MDS. The water quality index in deep, fine loamy, well drained soil was found to be 10.34. If all four MDS were considered to be responsible for contributing ideal (100%) water quality index, the we could conclude that pH was the key indicator of water quality. The guideline value of pH in drinking water set by WHO (World Health Organisation) is 6.5 to 8.5. Dutta *et al.*, 2013 reported that pH was an important factor of of water quality in the tea growing areas of Sonitpur district, Assam.

Conclusion:

From this study it was found that the water quality index under very deep, coarse loamy, well drained soil was found to be lowest (10.34) and the highest (15.38) was in very deep, fine loamy, well drained soil.

Reference

1. Andrews, S.S.; Mitchell, J.P.; Mancinelli, R.; Larlen, D.L.; Hartz, T.K.; Horwarth, W. R.; Pettygrove, G.S.; Scow, K.M. and Munk, D.S. (2002a). On farm assessment of soil quality in California's Central Valley. *Agron. J.* **94**: 12–23.
 2. Brejda, J.J.; Moorman, T.B.; Karlen, D.L. and Dao, T.H. (2000). Identification of regional soil quality factors and indicators. I. Central and Southern High Plains. *Soil Sci. Soc. Am. J.* **64**: 2115–2124
-

3. **C. R. Ramakrishnaiah**, C. Sadashivaiah, and G. Ranganna (2008). Assessment of Water Quality Index for the Groundwater in Tumkur Taluk, Karnataka State, India, Volume 6.

4. Chaudhury, J.; Mandal, U.K.; Sharma, K.L.; Ghosh, H. and Mandal, B. (2005). Assessing Soil Quality Under Long-Term Rice-Based Cropping System. *Commun. Soil Sci. Plant Anal.* **36**(9-10): 1141-1161.
5. Dutta, J.; Chetia, M.; Sarmah Baruah, J.P and Misra, A.K. (2010). Monitoring of Fluoride Concentration in Ground Water of Small Tea Gardens in Sunitpur District, Assam, India *Arch. Appl. Sci. Res.* **2**(5): 226-238.
6. Miller, W.P.; Martens, D.C. and Zelazny, L.W. (1985). Effects of manure amendment on soil chemical properties and hydrous oxides. *Soil Sci. Soc. Am. J.* **49**: 856-861.
7. Wander, M.M. and Bollero, G.A. (1999). Soil quality assessment of tillage impacts on Illinois. *Soil Sci. Soc. Am. J.* **63**: 961-971.
8. Joydev Dutta (2020). Ground Water Contamination in Tea Gardens Area: A Case Study in Gohpur Sub-Division, Biswanath, (Assam), India, IJCRT | Volume 8, Issue 11, ISSN: 2320-2882
9. Kumar, R., Paliyal, S. S., Sharma, S. K., & Sharma, S. (2023). Soil Quality Index for Different Cropping Systems in Northwestern Himalaya Region of India. *International Journal of Plant & Soil Science*, 35(24), 131-146. <https://doi.org/10.9734/ijpss/2023/v35i244305>
10. Choudhary, R., Naga, I. R., Jatara, A. M., Goyal, M., & Choudhary, D. (2024). Survey and Characterization of Ground Water Quality and its Effect on Soil Properties in Malpura Block of Tonk District Rajasthan, India. *Journal of Experimental Agriculture International*, 46(6), 293-302. <https://doi.org/10.9734/jeai/2024/v46i62480>
11. de Paul Obade V, Lal R. A standardized soil quality index for diverse field conditions. *Science of the total environment*. 2016 Jan 15;541:424-34.

UNDER PEER REVIEW