

Original Research Article

Response of Nitrogen Management on Growth and Yield of Groundnut

(Arachis hypogea L.)

Abstract

A field experiment was conducted during *Zaid* season of 2023 at Crop Research Farm Department of Agronomy, Naini Agriculture Institute, Sam Higginbottom University of Agriculture Sciences and Technology, To determine “Response of Nitrogen Management on Growth and Yield of Groundnut” The result revealed that treatment 6 [75% Nitrogen through poultry manure + 25% Nitrogen through urea] recorded significantly higher plant height (42.28 cm), maximum number of nodules/plant (21.06), higher plant dry weight (28.44 g), maximum number of pod/plant (28.31), higher number of kernels/pod (2.87), higher seed index (41.50 g), higher seed yield (3.30 t/ha), higher haulm yield (8.04 t/ha) and higher harvest index (29.95%).

Keywords: Groundnut, Organic manure, nitrogen, Growth, and yield.

1. INTRODUCTION

Groundnut (*Arachis hypogaea* L.) is a ~~self-self~~-pollinated, annual, herbaceous, ~~autotetraploid~~ autotetraploid legume with $2n = 40$ chromosomes belonging to the family leguminosae (Fabaceae). Groundnut is recognized as a golden or miracle bean due to its high nutritive value and various usage viz., for feed, oil, and soy food products. It is rich in protein (38-42%) and contains 18- 22 percent edible oil. It is known by various names such as “wondernut”, “Poor man’s Cashew nut”, “Peanut” “Monkey nut”, “Earth nut” or “King of Oilseeds” (**Tasnimet al., 2022**). It is the thirteenth most important food crop of the world; the fourth most important source of edible oil and the third most important source of vegetable protein. This crop has its own-importance due to the high edible oil content and nutritional value of kernel as human food, and haulm as rich feed for animals. Peanut is a very important source of oil (40-45%), protein (26%), carbohydrates (25%), minerals (Phosphorus,

calcium, and iron), and vitamins (vitamin B complexes like thiamine, riboflavin, niacin, and vitamin E) in addition to a higher proportion of unsaturated fatty acids, including essential fatty acids like linolenic and linoleic acids (Tasnimet *et al.*, 2022).

Groundnut ranks first in the area and second in production after soybean and is grown in almost all parts of the country over a wide range of agro-climatic conditions. Globally, groundnuts cover an area of 28.89 million hectares with the production of 54.41 million tons with the productivity of 1.88 tonnes/ha (USDA, 2024). In India Groundnut is grown over an area of about 5.75 million hectares with a production of 10.11 million tons and productivity of 1.7 t/ha (GOI 2022). The total area coverage under groundnut in Uttar Pradesh is 1.21 million hectares with a production of 1.24 million tones and the productivity of 1.02 t/ha (GOI, 2022).

The country is currently dealing with an edible oil issue as well as rising edible oil consumption due to population growth. Therefore, farmers should implement appropriate and improved groundnut production strategies to solve these obstacles. The primary cause of the low peanut production is the crop's restricted *kharif* season cultivation, which is subject to monsoon variations and foliar disease epidemics. Crops experience varying degrees of soil dehydration throughout deficiency monsoon, and the issue is made harder during above-average rainfall seasons by waterlogging. The productivity of summer groundnut is considerably higher than the *kharif* groundnut due to favorable conditions such as high temperature, more sunshine hours, assured irrigation under control conditions, and comparatively low incidence of disease and pests reduction in the use of chemical fertilizer and increased use of organic manure.

Poultry manure is advisable as the global environment pollution can be controlled considerably. The application of organic manure helps in mitigating multiple nutrient deficiencies and at the same time provides a better environment for growth and development by improving in physical, chemical, and biological properties of soil. In India, poultry farming is increasing.

The use of organic sources such as vermicompost is generally helpful for improving soil aggregation, structure, and fertility improving the moisture holding capacity and increasing crop yield.

Nitrogen is a critical and limiting element for the growth and development of most of the plants. Nitrogen is also an integral part of chlorophyll, which is the primary absorber of light energy needed for photosynthesis.

Therefore, judicious and combined use of organic and inorganic sources of plant nutrients plays an important role in the economizing the use of fertilizers under the increasing cost of chemical fertilizer ~~out~~ to find out the effect of organic and inorganic sources of nutrients on growth, yield, and economics of summer groundnut.

Keeping ~~in view of~~ given the above fact, the experiment was conducted to find out the “Response of Nitrogen Management on Growth and Yield of Groundnut”

2. MATERIALS AND METHODS:

The experiment was conducted during *Zaid* season 2023 at Crop Research Farm, Department of Agronomy, SHUATS, Prayagraj (U.P). The soil of the experimental field was sandy loam in texture, with soil (pH 7.8), low level of organic carbon (0.62%), available N (225 Kg/ha), P (38.2 kg/ha), K (240.7 kg/ha) and zinc (2.32 mg/kg). The experiment was laid out in RBD with 09 treatments each replicated thrice. The treatment combinations are T1- 100% Nitrogen through urea, T2- 75% Nitrogen through vermicompost + 25% Nitrogen through urea, T3- 50% Nitrogen through vermicompost + 50% Nitrogen through urea, T4- 25% Nitrogen through vermicompost + 75% Nitrogen through urea, T5- 100% Nitrogen through vermicompost, T6- 75% Nitrogen through poultry manure + 25% Nitrogen through urea, T7- 50% Nitrogen through poultry manure + 50 % Nitrogen through urea, T8-

25% Nitrogen through poultry manure + 75% Nitrogen through urea, T9- 100% Nitrogen through poultry manure. Data recorded on different aspects of [the](#) crop, viz., growth, yield attributes & yield were subjected to statistically ~~analysed~~[analyzed](#) by analysis of variance method as described by (Gomez and Gomez, 1976).

3. RESULT AND DISCUSSION

3.1 Growth Attributes

3.1.1 Plant height (cm)

The data revealed that significant and higher plant height (42.28 cm) was recorded in treatment 6 (75% Nitrogen through poultry manure + 25% Nitrogen through urea). However, treatment 2 (75% Nitrogen through vermicompost + 25% Nitrogen through urea) was found statistically at par with treatment 6 (75% Nitrogen through poultry manure + 25% Nitrogen through urea). Significant and higher plant height was observed with the application of 75% Nitrogen through poultry manure, [which](#) might be due to poultry manure ~~increased~~[increasing](#) the nutrient supply to the plants and ~~played~~[playing](#) a vital role in better root proliferation, higher cell division, higher biological nitrogen

fixation, and nutrient availability uptake, which would have resulted in increased high plant height similar to that reported by **Patel et al. (2022)**. Further, the higher plant height ~~was~~ observed with the application 25% Nitrogen through urea might be due to when urea was utilized via foliar application an increase in macro and micronutrient content and enhancing photosynthesis above it takes place and consequently affects the height of the plant. These results are similar to that reported by **Kharetwalet al. (2021)**.

3.1.2 Number of nodules/plant

The data revealed that a significant and maximum number of nodules/plant (21.06) was recorded in treatment 6 (75% Nitrogen through poultry manure + 25% Nitrogen through urea). However, treatment 2 (75% Nitrogen through vermicompost + 25% Nitrogen through urea) ~~which~~ was found statistically at par with treatment 6 (75% Nitrogen through poultry manure + 25% Nitrogen through urea). ~~Significant~~ A significant and maximum number of nodules/plant was observed with the application of 75% nitrogen through poultry manure, which might be due to the basal application of poultry manure increased nodules, supplied all essential nutrients, growth hormones, and enzymes to plant, which favour rapid cell division and ultimately increased nodules. ~~The~~ Similar result ~~were~~ was reported by **Mishra and Chaturvedi-Chaturvedi (2023)**. Further, the higher number of nodules/plants ~~was~~ observed with the application of 25% Nitrogen through urea might be due to the influence of nutrients to produce larger cells with thinner cell walls and its contribution ~~in~~ to cell division and cell elongation which improved vegetative growth and ultimately increased root nodules/plant. ~~Similar~~ A similar result was reported by **Vala et al. (2017)**.

Comment [BK1]: Add the results

3.1.3 Plant dry weight (g)

Results revealed that significant and higher dry weight (28.44 g) was recorded in treatment 6 (75% Nitrogen through poultry manure + 25% Nitrogen through urea). However, treatment 2 (75% Nitrogen through vermicompost + 25% Nitrogen through urea) ~~which~~ was found statistically at par with treatment 6 (75% Nitrogen through poultry manure + 25% Nitrogen through urea). Significant and maximum plant dry weight was observed with the application of 75% Nitrogen through poultry manure, which might be due to the application of poultry manure increased total dry matter, higher production of the new call, and increased photosynthesis ~~resulted~~ resulting in maximum plant dry weight. Similar reports were also in agreement by **Patel et al. (2022)**. Further significant maximum plant dry weight was 25% Nitrogen through urea increased may be due to nitrogen at higher levels

might have accelerated photosynthetic activity by increasing the source size (plant height and branch), thereby providing the developing bud with more photosynthates, thereby greater dry matter production (Palsandeet *et al.* 2019)

3.1.4 Crop Growth Rate (g/m²/day)

The data recorded during 60-80 DAS, intervals highest crop growth rate (18.46 g/m²/day) was observed in treatment 6 [75% Nitrogen through poultry manure + 25% Nitrogen through urea]. However, there was found no significant difference was found among all the treatments.

3.1.5 Relative Growth Rate (g/g/day)

The data revealed that during 60-80DAS, intervals higher relative growth rate (0.0298 g/g/day) was recorded in treatment 5 [100% Nitrogen through vermicompost] However there was found no significant difference was found among all the treatments.

3.2 Yield and Yield Parameters:

3.2.1 Number of pods/plant

The data recorded that a significant and maximum number of pod/plant (28.31) was recorded in treatment 6 [75% Nitrogen through poultry manure + 25% Nitrogen through urea]. However, treatment 2 [75% Nitrogen through vermicompost + 25% Nitrogen through urea] were found to be statistically at par with the treatment 6 [75% Nitrogen through poultry manure + 25% Nitrogen through urea]. The significant and maximum number of pods/plants with the application of 75% Nitrogen through poultry manure might be due to poultry manure supply of almost all plant essential nutrients by translocation of photosynthates accumulated under the influence of the source of organic nutrients. These results are similar to that reported by Patel *et al.* (2022). Further, the significant and maximum number of pods/ plants was with the application of 25% Nitrogen through urea may be due to the supply of nutrients from inorganic nutrient sources and prolonged availability of nutrients to the growing plant, which results in tissue differentiation from somatic to reproductive meristematic activity and increase in the development of floral primordia, resulting in higher number of pods/plant. These results were corroborated by Patel *et al.* (2022).

3.2.2 Number of kernels/pod

The data recorded that Significantly maximum Number of kernels/pod (2.87) were recorded in treatment 6 [75% Nitrogen through poultry manure + 25% Nitrogen through urea]. However, ~~the~~ treatment 2 [75% Nitrogen through vermicompost + 25% Nitrogen through urea] was found statically at par with treatment 6 [75% Nitrogen through poultry manure + 25% Nitrogen through urea]. ~~Significant~~ ~~The significant~~ and maximum number of kernels/pod obtained with the application of 75% Nitrogen through poultry manure might be due to ~~the~~ greater availability of metabolites (photosynthates) and nutrients to developing reproductive structures seems to have ~~resulted in increase in~~ ~~increased the~~ yield attributing characters particularly number of kernels/pod. ~~Similar A similar~~ result was reported by **Mishra and Chaturvedi (2023)**. Further, ~~the~~ significant and maximum number of kernels/pod ~~was~~ with the application of 25% Nitrogen through urea might be due to improvement in ~~the~~ nutritional environment which might have favourably influenced carbohydrate metabolism which in turn increased the uptake of nutrients and ultimately resulted in increased kernel weight and shelling percent. The findings agreed with those of **Vala et al. (2017)**.

3.2.3 Seed index (g)

Treatment 6 [75% Nitrogen through poultry manure + 25% Nitrogen through urea] recorded ~~the~~ highest seed index (41.50). ~~Though However~~ there is no significant difference found among all the treatments.

3.2.4 Seed yield (t/ha)

The data recorded Significant and higher seed yield (3.30 t/ha) was recorded in treatment 6 [75% Nitrogen through poultry manure + 25% Nitrogen through urea]. However, treatment 2 [75% Nitrogen through vermicompost + 25% Nitrogen through urea] ~~were was~~ found to be statistically at par with ~~the~~ treatment 6 [75% Nitrogen through poultry manure + 25% Nitrogen through urea]. Significant and higher seed yield was recorded with the application of RDN and poultry manure might be due to ~~an~~ adequate supply of nutrients through organic and inorganic sources enhanced the biological nitrogen fixation which might facilitates better root proliferation and increased ~~d~~ the seed yield. Similar results were reported by **Gowsalya et al. (2023)**.

3.2.5 Haulm yield (t/ha)

Significant and higher haulm yield (8.04 t/ha) was recorded in treatment 6 [75% Nitrogen through poultry manure + 25% Nitrogen through urea]. However, treatment 2 [75% Nitrogen through vermicompost + 25% Nitrogen through urea] ~~which~~ was found to be statistically at par with ~~the~~ treatment 6 [75% Nitrogen through poultry manure + 25% Nitrogen through urea]. Significant and

higher haulm yield was recorded with the application of poultry manure might be due to the better yield performance by increasing activities of N-fixing bacteria and increased rate of humification, Humic acid present in poultry manure may ~~be~~ enhanced the availability of added micronutrients in soil, thus improved yield attributes and haulm yield. The present findings are within the ~~close~~ proximity of **Devi et al. (2021)**. Further higher haulm yield ~~was~~ with the application of nitrogen might be due to proper fertilization coupled with increased net photosynthates on the one hand and greater mobilization of photosynthates towards reproductive structures on the other hand which might have increased the haulm yield. The present findings are within the ~~close~~ proximity of **Patel et al. (2022)**.

3.2.6 Harvest ~~index~~ **Index** (%)

~~Significant~~ **A significant** and higher harvest index (30.01%) was recorded in treatment 6 [75% Nitrogen through poultry manure + 25% Nitrogen through urea]. However, treatment 2 [75% Nitrogen through vermicompost + 25% Nitrogen through urea] ~~which~~ was found to be statistically at par with ~~the~~ treatment 6 [75% Nitrogen through poultry manure + 25% Nitrogen through urea]. ~~Significant~~ **A significant** and higher harvest index was observed with ~~the~~ application of poultry manure and RDF might be due to the ~~indicates~~ better partitioning of photosynthetic substance to economic yield, ~~an~~ appreciably high harvest index shows the efficiency of converting biological yield into economic yield. The present findings are within the ~~close~~ proximity of **Arifet al. (2014)**.

Table 1: Response of nitrogen management on growth attributes of groundnut.

| S.No. | Treatment combinations | Plant height (cm) (80 DAS) | Number of nodules/plant | Plant dry weight (g) | CRG (g/m ² /day) | RGR (g/g/day) |
|-------|------------------------------------------------------------------|----------------------------|-------------------------|----------------------|-----------------------------|---------------|
| 1 | 100% Nitrogen through urea | 35.33 | 16.46 | 23.32 | 16.35 | 0.0273 |
| 2 | 75% Nitrogen through vermicompost + 25% Nitrogen through urea | 40.04 | 19.93 | 27.1 | 17.58 | 0.0246 |
| 3 | 50% Nitrogen through vermicompost + 50% Nitrogen through urea | 36.41 | 16.86 | 23.51 | 16.27 | 0.0269 |
| 4 | 25% Nitrogen through vermicompost + 75% Nitrogen through urea | 32.28 | 15.53 | 22.68 | 16.5 | 0.0287 |
| 5 | 100% Nitrogen through vermicompost | 30.46 | 14.73 | 21.47 | 16.08 | 0.0298 |
| 6 | 75% Nitrogen through poultry manure + 25% Nitrogen through urea | 42.28 | 21.06 | 28.44 | 18.46 | 0.0246 |
| 7 | 50% Nitrogen through poultry manure + 50 % Nitrogen through urea | 36.76 | 17.73 | 24.44 | 15.08 | 0.0232 |
| 8 | 25% Nitrogen through poultry manure + 75% Nitrogen through urea | 34.72 | 16.13 | 23.06 | 16.52 | 0.0281 |
| 9 | 100% Nitrogen through poultry manure | 30.98 | 15 | 22.01 | 15.96 | 0.0286 |
| | F-test | S | S | S | NS | NS |
| | SEm(±) | 0.75 | 0.55 | 0.55 | 1.43 | 0.0020 |

CD (P=0.05)

2.26

1.64

1.66

-


-

Table 2: Response of nitrogen management on yield and yield attributes of Groundnut.

| Number of S.No. | Treatment combination | Number of kernels/pod | Seed index (g) | Seed Yield (t/ha) | Haulm Yield (t/ha) | Harvest Index (%) |
|-----------------|------------------------------------------------------------------|-----------------------|----------------|-------------------|--------------------|-------------------|
| 1 | 100% Nitrogen through urea | 1.98 | 40.17 | 1.97 | 5.79 | 25.42 |
| 2 | 75% Nitrogen through vermicompost + 25% Nitrogen through urea | 2.81 | 41.21 | 3.16 | 7.41 | 29.80 |
| 3 | 50% Nitrogen through vermicompost + 50% Nitrogen through urea | 1.87 | 40.58 | 1.97 | 6.09 | 24.40 |
| 4 | 25% Nitrogen through vermicompost + 75% Nitrogen through urea | 1.77 | 39.87 | 1.72 | 5.53 | 23.73 |
| 5 | 100% Nitrogen through vermicompost | 1.47 | 38.71 | 1.31 | 4.89 | 21.14 |
| 6 | 75% Nitrogen through poultry manure + 25% Nitrogen through urea | 2.87 | 41.50 | 3.30 | 8.04 | 29.95 |
| 7 | 50% Nitrogen through poultry manure + 50 % Nitrogen through urea | 2.37 | 40.97 | 2.58 | 6.48 | 28.42 |
| 8 | 25% Nitrogen through poultry manure + 75% Nitrogen through urea | 1.60 | 39.68 | 1.52 | 5.53 | 21.59 |
| 9 | 100% Nitrogen through poultry manure | 1.55 | 39.20 | 1.42 | 5.12 | 21.63 |

| | | | | | |
|------------|------|------|------|------|------|
| Ftest | S | NS | S | S | S |
| SEm(±) | 0.09 | 0.73 | 0.06 | 0.26 | 0.58 |
| CD(p=0.05) | 0.27 | 2.20 | 0.19 | 0.78 | 1.74 |

UNDER PEER REVIEW



CONCLUSION

It is concluded that (treatment 6), [the](#) application of 75% Nitrogen through poultry manure and 25% Nitrogen through urearecorded [the](#) highest growth and seed yield.

REFERENCES

Agriculture, Government of India, New Delhi; c2022. Available from:
<https://eands.dacnet.nic.in>.

Arif, M., Tasneem, M., Bashir, F., Yaseen, G. and Iqbal, R.M. (2014). effect of integrated use of organic manures and inorganic fertilizers on yield and yield components of rice. *Journal of agriculture research* **52**(2): 197-206.

Devi, R., Singh, V., Tiwari, D., Sudhakar, m. and Rakesh, M. (2021). Influence of organic manures and moybdenum on growth and yield of groundnut. *An international journal* **13**(3): 402-406.

GOI. Agricultural Statistics at a Glance. Agricultural Statistics Division, Directorate of Economics and Statistics, Department of Agriculture and Cooperation, Ministry of Gomez KA, Gomez AA. (1976) Statistical procedures for agriculture research, 2nd edition, John Wiley and Son, New York. 680p.

Gowasalya, M., Murugan, G., Suseendran, K., Stalin, P. and Perumal, M.S. (2023). Effect of Integrated Application of Organic Manure and Inorganic Fertilizer on Yield, Quality and Economics of Machine Sown Groundnut. *Journal of Emerging Technology and Innovative Research* **10**(2): 2349-5162.

Kharetwal, N., Ghatak, R.G., Thaokar, A., Gawali, K., Sarda, A. and Nagmote, A. (2021) effect of different nutrient management practices on growth, yield and economics in chickpea. *The pharma innovation journal* **10**(9): 993-996.

Mishra, K.S. and Chaturvedi, P.K. (2023). effect of integrated nutrient management on growth and yield of green gram. *The pharma innovation journal* **12**(1): 671-673.

Palsande, V.N., Nagral, M.R., Kasture, M.C., Gokhale, N.B., Dhekale, J.S. and Salvi, V.G. (2019) Growth, yield and quality of kharif groundnut as affected by different levels of, potassium and zinc in lateritic soils of konkan. *Journal of pharmacognosy and phytochemistry* **8**(5): 790-794.

Patel, R., Patel, P., Patel, H., Patel, H. and Patel, T. (2022). Nitrogen management through organic sources and biofertilizers in summer groundnut. *The pharma innovation journal***11**(9): 1342-1347.

Patel, T., Patel, R.A. and Patel, P. (2022) Effect of irrigation scheduling and nitrogen management on growth and yield of summer groundnut. *The Pharma Innovation Journal* **11**(9): 1623-1627.

Patel, T., Patel, R.A. and Patel, P. (2022). Effect of irrigation scheduling and nitrogen management on growth and yield of summer groundnut. *The Pharma Innovation Journal***11**(9): 1623-1627.

Patil, D.B., Pawar, P.P., Wadile, S.C. and Patil, H.M. (2017) Effect of integrated nutrient management on growth and yield of kharif groundnut. *International journal of horticulture and science***1**(1): 21-23.

Tasnim, S., Ivy, N.A., Raihan, M.S., Akter, N., Islam, M.Z., Jahan, N., and Khan, A.U. (2022). Evaluation of Genetic Variation in Segregating Population of Pea (*Pisum sativum* L.). *SAARC Journal of Agriculture***20**(1): 29-39. Doi: <https://doi.org/10.3329/sja.v20i1.60545>

Tasnim, S., Poly, N.Y., Jahan, N., and Khan, A.U. (2022). Relationship of Quantitative Traits in Different Morphological Characters of Pea (*Pisum Sativum* L.). *Journal of Multidisciplinary Applied Natural Science***2**(2): 103-114. Doi: <https://doi.org/10.47352/jmans.2774-3047.119>.

USDA. (2023). United states Department of Agriculture. World Agricultural Production. Vala, F.G., Vaghasis, P.M., Zala, K.P. and Buha D.B. (2017). Effect of integrated nutrient management on productivity of summer groundnut. *International journal of current microbiology and applied sciences* **6**(10): 1951-1957.