

Common Fixed-Point Theorem for Expansive Mappings in Dualistic Partial Metric Spaces

Abstract

O'Neill [14] introduces the concept of dualistic partial metric space. In this study, we prove some common fixed-point theorems for dualistic expanding mappings defined on a dualistic partial metric space. Some famous conclusions of [5] and [17] are extended and generalized by our result. Additionally, we offer an example that demonstrates the value of these dualistic expanding mappings.

2010 Mathematics Subject Classification: 47H10

Keywords: common fixed point, dualistic partial metric, dualistic expanding mappings.

1. Introduction

“Because metric fixed-point theory has so many applications in applied mathematics and the sciences, it is becoming more and more important in mathematics. The conventional understanding of a metric space has been generalized in several ways. A partial metric space, which Matthews developed and examined, is one such generalization” [7]. He verified the exact correspondence between the so-called weightable quasi-metric spaces and partial metric spaces. Partial metrics have certain generalizations. One major modification to Matthews' formulation of the partial metric, for instance, was suggested by O'Neill [14] and involved moving its range from $[0, \infty)$ to $(-\infty, \infty)$. A pair (\mathcal{U}, d^*) where \mathcal{U} is a nonempty set and d^* is a dualistic partial metric on \mathcal{U} is referred to as a dualistic partial metric space. According to [14], “the partial metrics in the O'Neill sense shall be called dualistic partial metrics. In this way, O'Neill established several links between partial metrics and the topological aspects of domain theory”. The study of Banach's contraction principle served as the foundation for contractive conditions. These conditions

have been used in many generalized metric spaces for fixed point theorems. After that, expansive conditions were added [17], and expansive mappings were used to provide new fixed-point solutions. For different contractive or expansive mappings, several fixed-point findings have still been studied utilizing the concepts of metric space and partial metric space. View [3], [4], [5], [6], [8] [15], [16], and [19] for additional information.

The existence of fixed points for a particular class of mappings known as "expansive mappings" is addressed by the fixed-point theorem for expansive mappings, a finding in fixed point theory. According to the theorem, there is always at least one fixed point in an expansive mapping defined on a nonempty bounded metric space. A concept of non-contraction is reflected by expansive mappings, which lead points to spread apart during the transformation. Put otherwise, the theorem asserts that a mapping can have invariant points even in the absence of a contraction in the distances between its points. The idea behind it is to expand upon the conventional fixed-point theorem for contraction mappings, which asserts that every contraction mapping has a single fixed point.

Theorem 1.1 Let \mathcal{U} be a bounded metric space that is not empty, and let f be an expansive mapping from \mathcal{U} to \mathcal{U} . Then, f has a fixed point.

A number of diverse scenarios have been expanded upon using this theorem [15, 26, 28]. Applications of the expansive mappings fixed point theorem can be found in dynamical systems, differential equations, and nonlinear analysis. The idea of expansive mappings makes sense in these domains. Expansive mappings, for instance, are employed in nonlinear analysis to examine the behavior of dynamical systems [27]. Expanding mappings are used in differential and integral equations to examine the stability of solutions [29]. Expansive mappings are widely utilized in dynamical systems to examine the existence of chaotic attractors [27]. This technique is applied in many different domains, such as the study of dynamics in mathematical biology and economics.

In a dualistic partial metric space, the purpose of this study is to show several common fixed-point theorems under different dualistic expansive mappings. Some famous conclusions of [5] and [17] are extended and generalized by our result. Furthermore, we validate our findings using an example.

2. Preliminaries

To make this work self-contained, we review key definitions and foundational concepts in mathematics.

Definition 2.1. (see[7]) Let \mathcal{U} be a non-empty set. A partial metric on \mathcal{U} is a function $d: \mathcal{U} \times \mathcal{U} \rightarrow [0, \infty)$ complying with following axioms, for all $x, y, z \in \mathcal{U}$

$$\begin{aligned} (d_1) x = y &\Leftrightarrow d(x, y) = d(x, x) = d(y, y), \\ (d_2) d(x, x) &\leq d(x, y), \\ (d_3) d(x, y) &= d(y, x), \\ (d_4) d(x, y) &\leq d(x, z) + d(z, y) - d(z, z). \end{aligned}$$

The pair (\mathcal{U}, d) is called a partial metric space.

Definition 2.2. (see [14]) Let \mathcal{U} be a non-empty set. A dualistic partial metric on \mathcal{U} is a function $d^*: \mathcal{U} \times \mathcal{U} \rightarrow (-\infty, \infty)$ satisfying the following axioms, for all $x, y, z \in \mathcal{U}$

$$\begin{aligned} (d_1^*) x = y &\Leftrightarrow d^*(x, y) = d^*(x, x) = d^*(y, y), \\ (d_2^*) d^*(x, x) &\leq d^*(x, y), \\ (d_3^*) d^*(x, y) &= d^*(y, x), \\ (d_4^*) d^*(x, z) + d^*(y, y) &\leq d^*(x, y) + d^*(y, z). \end{aligned}$$

The pair (\mathcal{U}, d^*) is called a dualistic partial metric space.

Remark 2.3 Noting that each partial metric is a dualistic partial metric, but the converse is false. Indeed, define d^* on $(-\infty, \infty)$ as $d^*(x, y) = \max\{x, y\}, \forall x, y \in (-\infty, \infty)$. Obviously, d^* is a dualistic partial metric on $(-\infty, \infty)$. Since $d^*(x, y) < 0 \notin [0, \infty), \forall x, y \in (-\infty, 0)$ and then d^* is not a partial metric on $(-\infty, \infty)$. This confirms our remark.

Example 2.4(see [10, 14])

- (1) Define $d_\rho^*: \mathcal{U} \times \mathcal{U} \rightarrow (-\infty, \infty)$ by $d_\rho^*(x, y) = \rho(x, y) + b$, where ρ is a metric on a nonempty set \mathcal{U} and $b \in (-\infty, \infty)$ is arbitrary constant, then it is easy to check that d_ρ^* verifies axioms $(d_1^*) - (d_4^*)$ and hence (\mathcal{U}, d^*) is a dualistic partial metric space.
- (2) Let d be a partial metric defined on a non-empty set \mathcal{U} . The function $d^*: \mathcal{U} \times \mathcal{U} \rightarrow (-\infty, \infty)$ defined by $d^*(x, y) = d(x, y) - d(x, x) - d(y, y)$ satisfies the axioms $(d_1^*) - (d_4^*)$ and so it defines a dualistic partial metric on \mathcal{U} . Note that $d^*(x, y)$ may have negative values.

(3) Let $\mathcal{U} = (-\infty, \infty)$. Defined*: $\mathcal{U} \times \mathcal{U} \rightarrow (-\infty, \infty)$ by $d^*(x, y) = |x - y|$ if $x \neq y$ and $d^*(x, y) = -\beta$ if $x = y$ and $\beta > 0$. We can easily see that d^* is a dualistic partial metric on \mathcal{U} .

O'Neill [14] established that each dualistic partial metric d^* on \mathcal{U} generates a T_0 topology $\tau(d^*)$ on \mathcal{U} having a base, the family of d^* -balls $\{\mathcal{B}_{d^*}(x, \epsilon) \mid x \in \mathcal{U}, \epsilon > 0\}$, where

$$(2.1) \mathcal{B}_{d^*}(x, \epsilon) = \{y \in \mathcal{U} \mid d^*(x, y) < d^*(x, x) + \epsilon\}.$$

If (\mathcal{U}, d^*) is a dualistic partial metric space, then the function $\rho_{d^*} : \mathcal{U} \times \mathcal{U} \rightarrow [0, \infty)$ defined by

$$(2.2) \rho_{d^*}(x, y) = d^*(x, y) - d^*(x, x),$$

defines a quasi-metric on \mathcal{A} such that $\tau(d^*) = \tau(\rho_{d^*})$ and

$$(2.3) \rho_{d^*}^m(x, y) = \max\{\rho_{d^*}(x, y), \rho_{d^*}(y, x)\},$$

defines a metric on \mathcal{U} .

Definition 2.5 (see [13]) Let (\mathcal{U}, d^*) be a dualistic partial metric space.

1. A sequence $\{x_n\}$ in \mathcal{U} is said to converge or to be convergent if there is a $x \in \mathcal{U}$ such that $\lim_{n \rightarrow \infty} d^*(x_n, x) = d^*(x, x)$. x is called the limit of $\{x_n\}$ and we write $x_n \rightarrow x$.
2. A sequence $\{x_n\}$ in \mathcal{U} is said to be Cauchy sequence if $\lim_{n, m \rightarrow \infty} d^*(x_n, x_m)$ exists and is finite.
3. A dualistic partial metric space $\mathcal{U} = (\mathcal{U}, d^*)$ is said to be complete if every Cauchy sequence $\{x_n\}$ in \mathcal{U} converges, with respect to $\tau(d^*)$, to a point $x \in \mathcal{U}$ such that $d^*(x, x) = \lim_{n, m \rightarrow \infty} d^*(x_n, x_m)$.

Remark 2.6 For a sequence, convergence with respect to metric space may not imply convergence with respect to dualistic partial metric space. Indeed, if we take $\beta = 1$ and $\{x_n = \frac{1-n}{n} : n \geq 1\}_{n \in \mathbb{N}} \subset \mathcal{U}$ as in Example 2.4(3). Mention that $\lim_{n \rightarrow \infty} \rho(x_n, -1) = -1$ and therefore, $x_n \rightarrow -1$ with respect top. On the other hand, we make a conclusion that $x_n \not\rightarrow -1$ with respect to d^* because

$$\lim_{n \rightarrow \infty} d^*(x_n, -1) = \lim_{n \rightarrow \infty} d^*|x_n - (-1)| = \lim_{n \rightarrow \infty} \left| \frac{1-n}{n} + 1 \right| = 0$$

and $d^*(-1, -1) = -1$.

Lemma 2.7(see [13])Let (\mathcal{U}, d^*) be a dualistic partial metric space.

- (1) Every Cauchy sequence in $(\mathcal{U}, \rho_{d^*}^m)$ is also a Cauchy sequence in (\mathcal{U}, d^*) .
- (2) A dualistic partial metric (\mathcal{U}, d^*) is complete if and only if the induced metric space $(\mathcal{U}, \rho_{d^*}^m)$ is complete.
- (3) A sequence $\{x_n\}$ in \mathcal{U} converges to a point $x \in \mathcal{U}$ with respect to $\tau(\rho_{d^*}^m)$ if and only if $d^*(x, x) = \lim_{n \rightarrow \infty} d^*(x_n, x) = \lim_{n \rightarrow \infty} d^*(x_n, x_m)$.

Then \mathcal{P} is a parametric metric on \mathcal{U} and the pair $(\mathcal{U}, \mathcal{P})$ is a parametric metric space. Let \mathcal{U} be a set. A point $z \in \mathcal{U}$ is a point of coincidence of a pair of self-maps $f, g: \mathcal{U} \rightarrow \mathcal{U}$ and $\theta \in \mathcal{U}$ is its coincidence point if $f\theta = g\theta = z$. Mappings f and g are weakly compatible if $f g \theta = g f \theta$ for each of their coincidence points θ [20, 23, 24] and occasionally weakly compatible if the same holds for some coincidence point [22]. The set of fixed points of a self-map $f: \mathcal{U} \rightarrow \mathcal{U}$ will be denoted as $\mathfrak{F}(f)$. The mapping f is said to possess property (P) if $\mathfrak{F}(f^n) = \mathfrak{F}(f)$ for each $n \in \mathbb{N}$ (see [20, 25]). A pair of self-maps $f, g: \mathcal{U} \rightarrow \mathcal{U}$ is said to have property (Q) if $\mathfrak{F}(f^n) \cap \mathfrak{F}(g^n) = \mathfrak{F}(f) \cap \mathfrak{F}(g)$ holds for each $n \in \mathbb{N}$ (see [20]).

Definition 2.8(see [30])Let (\mathcal{U}, d^*) be a dualistic partial metric space and $f: \mathcal{U} \rightarrow \mathcal{U}$. Then f is called a dualistic expanding mapping, if for every $x, y \in \mathcal{U}$, there exists a number $\lambda > 1$ such that

$$|d^*(fx, fy)| \geq \lambda |d^*(x, y)|.$$

3. Main Result

Our main result as follows.

Theorem 3.1 Let (\mathcal{U}, d^*) be a complete dualistic partial metric space and $f, g: \mathcal{U} \rightarrow \mathcal{U}$ be two maps such that $f(\mathcal{U}) \supset g(\mathcal{U})$ and that at least one of these subspaces is complete. Suppose that there exist real numbers α, β, γ satisfying $\beta, \gamma \geq 0$ and $\alpha > 1, \gamma < 1$ such that

$$(3.1) |d^*(fx, fy)| \geq \alpha |d^*(gx, gy)| + \beta |d^*(gx, fx)| + \gamma |d^*(gy, fy)|,$$

$\forall x, y \in \mathcal{U}$. Then f and g have a unique point of coincidence. If, moreover, the pair (f, g) is (occasionally) weakly compatible, then f and g have a unique common fixed point.

Proof Take arbitrary $x_0 \in \mathfrak{D}$. Construct sequences $\{x_n\}$ and $\{y_n\}$ such that $y_n = gx_n = fx_{n+1}$ for $n = 0, 1, 2, \dots$. Applying (3.1), we obtain

$$\begin{aligned} |d^*(y_n, y_{n-1})| &= |d^*(fx_{n+1}, fx_n)| \\ &\geq \alpha |d(gx_{n+1}, gx_n)| + \beta |d^*(gx_{n+1}, fx_{n+1})| + \gamma |d^*(gx_n, fx_n)| \\ &= \alpha |d(y_{n+1}, y_n)| + \beta |d^*(y_{n+1}, y_n)| + \gamma |d^*(y_n, y_{n-1})| \\ &= (\alpha + \beta) |d^*(y_{n+1}, y_n)| + \gamma |d^*(y_n, y_{n-1})| \end{aligned}$$

Hence

$$(3.2) \quad (1 - \gamma) |d^*(y_{n-1}, y_n)| \geq (\alpha + \beta) |d^*(y_n, y_{n+1})|$$

Consequently,

$$(3.3) \quad |d^*(y_n, y_{n+1})| \leq \mu |d^*(y_{n-1}, y_n)|$$

where $\mu = \frac{1-\gamma}{\alpha+\beta} < 1$ for all $n \in \mathbb{N} \cup \{0\}$ and $t > 0$. Repeating (3.3) r -times, we get

$$(3.6) \quad |d^*(y_n, y_{n+1})| \leq \mu^n |d^*(y_0, y_1)|.$$

Now,

$$\begin{aligned} |d^*(y_n, y_n)| &= |d^*(fx_{n+1}, fx_{n+1})| \\ &\geq \alpha |d(gx_{n+1}, gx_{n+1})| + \beta |d^*(gx_{n+1}, fx_{n+1})| + \gamma |d^*(gx_{n+1}, fx_{n+1})| \\ &= \alpha |d(y_{n+1}, y_{n+1})| + \beta |d^*(y_{n+1}, y_n)| + \gamma |d^*(y_{n+1}, y_n)| \\ &= (\beta + \gamma) |d^*(y_{n+1}, y_n)| + \alpha |d(y_{n+1}, y_{n+1})| \end{aligned}$$

The last inequality gives

$$(3.7) \quad |d(y_{n+1}, y_{n+1})| \leq \frac{1}{\alpha} |d^*(y_n, y_n)| - \frac{(\beta+\gamma)}{\alpha} |d^*(y_{n+1}, y_n)|.$$

Due to inequality (3.6), we have

$$(3.8) \quad |d(y_{n+1}, y_{n+1})| \leq \frac{1}{\alpha} |d^*(y_n, y_n)| - \frac{(\beta+\gamma)}{\alpha} \mu^n |d^*(y_0, y_1)|.$$

Similarly,

$$(3.9) \quad |d^*(y_n, y_n)| \leq \frac{1}{\alpha} |d^*(y_{n-1}, y_{n-1})| - \frac{(\beta+\gamma)}{\alpha} \mu^{n-1} |d^*(y_0, y_1)|.$$

The inequality (3.6) implies that

$$\begin{aligned}
|d(y_{n+1}, y_{n+1})| &\leq \frac{1}{\alpha} \left\{ \frac{1}{\alpha} |d^*(y_{n-1}, y_{n-1})| - \frac{(\beta + \gamma)}{\alpha} \mu^{n-1} |d^*(y_0, y_1)| \right\} \\
&\quad - \frac{(\beta + \gamma)}{\alpha} \mu^n |d^*(y_0, y_1)| \\
&= \frac{1}{\alpha^2} |d^*(y_{n-1}, y_{n-1})| - \frac{(\beta + \gamma)}{\alpha^2} \mu^{n-1} |d^*(y_0, y_1)| \\
&\quad - \frac{(\beta + \gamma)}{\alpha} \mu^n |d^*(y_0, y_1)| \\
&= \frac{1}{\alpha^2} |d^*(y_{n-1}, y_{n-1})| - (\beta + \gamma) \left[\frac{\mu^{n-1}}{\alpha^2} + \frac{\mu^n}{\alpha} \right] |d^*(y_0, y_1)| \\
&\leq \frac{1}{\alpha^3} |d^*(y_{n-2}, y_{n-2})| - (\beta + \gamma) \left[\frac{\mu^{n-2}}{\alpha^3} + \frac{\mu^{n-1}}{\alpha^2} + \frac{\mu^n}{\alpha} \right] |d^*(y_0, y_1)|.
\end{aligned}$$

Continuing further, we get

$$\begin{aligned}
(3.10) |d(y_{n+1}, y_{n+1})| &\leq \frac{1}{\alpha^{n+1}} |d^*(y_0, y_0)| - (\beta + \gamma) \left[\frac{1}{\alpha^{n+1}} + \frac{\mu}{\alpha^n} + \dots + \frac{\mu^n}{\alpha} \right] |d^*(y_0, y_1)| \\
&\leq \delta^{n+1} |d^*(y_0, y_0)| - (\beta + \gamma) [\delta^{n+1} + \delta^n \mu + \dots + \delta \mu^n] |d^*(y_0, y_1)| \\
&= \delta^{n+1} |d^*(y_0, y_0)| - (\beta + \gamma) \rho^{n+1} |d^*(y_0, y_1)| \\
&\leq \delta^{n+1} |d^*(y_0, y_0)| + \rho^{n+1} |d^*(y_0, y_1)|,
\end{aligned}$$

where $\delta = \frac{1}{\alpha}$ and $\rho^{n+1} = \delta^{n+1} + \delta^n \mu + \dots + \delta \mu^n$.

We deduce from (2.2) that

$$\begin{aligned}
(3.11) \rho_{d^*}(y_n, y_{n+1}) &\leq |d^*(y_n, y_{n+1})| - d^*(y_n, y_n) \\
&\leq |d^*(y_n, y_{n+1})| + d^*(y_n, y_n) \\
&\leq \mu^n |d^*(y_0, y_1)| + \delta^n |d^*(y_0, y_0)| + \rho^n |d^*(y_0, y_1)| \\
&= (\mu^n + \rho^n) |d^*(x_0, x_1)| + \delta^n |d^*(x_0, x_0)|.
\end{aligned}$$

Now, for $m > n$, we have

$$\begin{aligned}
\rho_{d^*}(y_n, y_m) &\leq \rho_{d^*}(y_n, y_{n+1}) + \rho_{d^*}(y_{n+1}, y_{n+2}) + \cdots + \rho_{d^*}(y_{m-1}, y_m) \\
&\leq (\mu^n + \rho^n)|d^*(y_0, y_1)| + \delta^n|d^*(y_0, y_0)| \\
&+ (\mu^{n+1} + \rho^{n+1})|d^*(y_0, y_1)| + \delta^{n+1}|d^*(y_0, y_0)| + \cdots \\
&+ (\mu^{m-1} + \rho^{m-1})|d^*(y_0, y_1)| + \delta^{m-1}|d^*(y_0, y_0)| \\
&= (\mu^n + \mu^{n+1} + \cdots + \mu^{m-1})|d^*(y_0, y_1)| \\
&+ (\rho^n + \rho^{n+1} + \cdots + \rho^{m-1})|d^*(y_0, y_1)| \\
&+ (\delta^n + \delta^{n+1} + \cdots + \delta^{m-1})|d^*(y_0, y_0)| \\
&\leq (\mu^n + \mu^{n+1} + \cdots + \mu^{m-1} + \cdots)|d^*(y_0, y_1)| \\
&+ (\rho^n + \rho^{n+1} + \cdots + \rho^{m-1} + \cdots)|d^*(y_0, y_1)| \\
&+ (\delta^n + \delta^{n+1} + \cdots + \delta^{m-1} + \cdots)|d^*(y_0, y_0)| \\
&= \frac{\mu^n}{1-\mu}|d^*(y_0, y_1)| + \frac{\rho^n}{1-\rho}|d^*(y_0, y_1)| + \frac{\delta^n}{1-\delta}|d^*(y_0, y_0)|.
\end{aligned}$$

Hence

$$(3.12) \quad \rho_{d^*}(y_n, y_m) \leq \frac{\mu^n}{1-\mu}|d^*(y_0, y_1)| + \frac{\rho^n}{1-\rho}|d^*(y_0, y_1)| + \frac{\delta^n}{1-\delta}|d^*(y_0, y_0)|.$$

As $m, n \rightarrow \infty$, $\rho_{d^*}^m(y_n, y_m) = \max\{\rho_{d^*}(y_n, y_m), \rho_{d^*}(y_m, y_n)\} \rightarrow 0$, thus, $\{y_n\}$ is a Cauchy sequence in $(\mathcal{U}, \rho_{d^*}^m)$. Since (\mathcal{U}, d^*) is a complete dualistic partial metric space, by Lemma 2.1(2), $(\mathcal{U}, \rho_{d^*}^m)$ is a complete metric space. Thus, there exists $z \in (\mathcal{U}, \rho_{d^*}^m)$ such that $y_n = g x_n = f x_{n+1} \rightarrow z$ as $n \rightarrow \infty$, that is $\lim_{n \rightarrow \infty} \rho_{d^*}(y_n, z) = 0$ and by Lemma 2.1 (3), we know that

$$(3.13) \quad d^*(z, z) = \lim_{n \rightarrow \infty} d^*(y_n, z) = \lim_{n \rightarrow \infty} d^*(y_n, y_m).$$

Since, $\lim_{n \rightarrow \infty} \rho_{d^*}(y_n, z) = 0$, by (2.2) and (3.27), we have

$$(3.14) \quad d^*(z, z) = \lim_{n \rightarrow \infty} d^*(y_n, z) = \lim_{n \rightarrow \infty} d^*(y_n, y_m) = 0.$$

This shows that $\{y_n\}$ is a Cauchy sequence in (\mathcal{U}) . Suppose that $f(\mathcal{U})$ is complete. Then, there exists $\theta \in \mathcal{U}$ such that $fx_n \rightarrow f\theta = z$. Let us prove that $f\theta = g\theta$. Then, using (3.1), we get

$$\begin{aligned} |d^*(y_n, f\theta)| &= |d^*(fx_{n+1}, f\theta)| \\ &\geq \alpha |d(gx_{n+1}, g\theta)| + \beta |d^*(gx_{n+1}, fx_{n+1})| + \gamma |d^*(g\theta, f\theta)| \end{aligned}$$

which implies that as $n \rightarrow +\infty$,

$$0 \geq (\alpha + \gamma) |d(f\theta, g\theta)|$$

Hence $f\theta = g\theta$. Thus, $f\theta = g\theta = z$ is a point of coincidence for (f, g) . Suppose that there is another point of coincidence $f\theta_1 = g\theta_1 = z_1$. Then

$$\begin{aligned} |d^*(z, z_1)| &= |d^*(f\theta, f\theta_1)| \\ &\geq \alpha |d(g\theta, g\theta_1)| + \beta |d^*(g\theta, f\theta)| + \gamma |d^*(g\theta_1, f\theta_1)| \end{aligned}$$

implying (since $\alpha > 1$) that $|d^*(z, z_1)| = 0 = |d^*(z, z)| = |d^*(z_1, z_1)|$. Consequently, $z = z_1$. Thus, the point of coincidence is unique. If the pair (f, g) is weakly compatible, applying [21, Proposition 1.12] we conclude that f and g have a unique common fixed point. If (f, g) is occasionally weakly compatible, the same conclusion follows from [22, Lemma 1.6]. This completes the proof.

Setting $\beta = 0 = \gamma$ in Theorem 3.1, we can obtain the following result.

Corollary 3.2 Let (\mathcal{U}, d^*) be a complete dualistic partial metric space and $f, g: \mathcal{U} \rightarrow \mathcal{U}$ be two maps such that $f(\mathcal{U}) \supset g(\mathcal{U})$ and that at least one of these subspaces is complete. Suppose that there exists a real number $\alpha > 1$ such that

$$(3.15) \quad |d^*(fx, fy)| \geq \alpha |d(gx, gy)|,$$

$\forall x, y \in \mathcal{U}$. Then f and g have a unique point of coincidence. If, moreover, the pair (f, g) is (occasionally) weakly compatible, then f and g have a unique common fixed point.

Corollary 3.3 Let (\mathcal{U}, d^*) be a complete dualistic partial metric space and $f: \mathcal{U} \rightarrow \mathcal{U}$ be a surjection. Suppose that there exists a real number $\alpha > 1$ such that

$$(3.16) \quad |d^*(fx, fy)| \geq \alpha |d(x, y)|,$$

$\forall x, y \in \mathcal{U}$. Then f has a unique fixed point.

Proof From Corollary 3.3, it follows that f has a fixed point z in \mathcal{U} by setting $g = i_{\mathcal{U}}$.

Uniqueness. Suppose that $z \neq z^*$ is also another fixed point of f , then from condition (3.16), we obtain

$$|d^*(z, z^*)| = |d^*(fz, fz^*)| \geq \alpha |d(z, z^*)|$$

which implies $d^*(z, z^*) = 0 = d^*(z, z) = d^*(z^*, z^*)$, that is $z = z^*$. This completes the proof.

Corollary 3.4 Let (\mathcal{U}, d^*) be a complete dualistic partial metric space and $f: \mathcal{U} \rightarrow \mathcal{U}$ be a surjection. Suppose that there exists a positive integer n and a real number $\alpha > 1$ such that

$$(3.17) \quad |d^*(f^n x, f^n y)| \geq \alpha |d(x, y)|,$$

$\forall x, y \in \mathcal{U}$. Then f has a unique fixed point.

Proof From Corollary 3.3, f^n has a fixed point z . But $f^n(fz) = f(f^n z) = fz$, So fz is also a fixed point of f^n . Hence $fz = z$, z is a fixed point of f . Since the fixed point of f is also fixed point of f^n , the fixed point of f is unique.

Corollary 3.5 (Corollary 2.1 of Huang et al. [5]) Let (\mathcal{U}, d) be a complete partial metric space and $f: \mathcal{U} \rightarrow \mathcal{U}$ be a surjection. Suppose that there exists $\lambda > 1$ such that

$$(3.18) \quad d(fx, fy) \geq \lambda d(x, y),$$

$\forall x, y \in \mathcal{U}$, then f has a unique fixed point.

Proof Since the restriction of a dualistic partial metric d^* to $[0, \infty)$, $d^*|_{[0, \infty)} = d$ is a partial metric, so arguments follow the same lines as in the proof of Theorem 3.1.

Corollary 3.6 (Theorem 2.1 of Huang et al. [5]) Let (\mathcal{U}, d) be a complete partial metric space and $f: \mathcal{U} \rightarrow \mathcal{U}$ be a surjection. Suppose that there exist real numbers α, β, γ satisfying $\beta, \gamma \geq 0$ and $\alpha > 1$ such that

$$(3.19) \quad d(fx, fy) \geq \alpha d(x, y) + \beta d(x, fx) + \gamma d(y, fy),$$

$\forall x, y \in \mathcal{U}$, then f has a unique fixed point.

Proof Set $d^*|_{[0, \infty)} = d$ and arguments follow the same lines as in the proof of Theorem 3.1.

Observations 3.7

1. Usually the range of a dualistic partial metric d^* is $(-\infty, \infty)$ but if we replace $(-\infty, \infty)$ by $[0, \infty)$, then d^* is identical to a partial metric d and hence Theorem 3.1 is applicable in the setting of partial metric space.
2. If we set $d(x, x) = 0$ in Corollary 3.5 and Corollary 3.6, we retrieve corresponding theorems in metric spaces.
3. Our main result extends and generalizes some well-known results of Huang et al. [5] and Wang et al. [17].

Example 3.8 Define $d^*: (-\infty, 0] \times (-\infty, 0] \rightarrow (-\infty, \infty)$ by $d^*(x, y) = \max\{x, y\}$. It is easy to check that $((-\infty, 0], d^*)$ is a complete dualistic partial metric space. Define $f, g: (-\infty, 0] \rightarrow (-\infty, 0]$ as $fx = 6x, gx = 3x, \forall x \in (-\infty, 0]$. Further, for all $x, y \in (-\infty, 0]$ with $x \geq y$, and $\alpha = 2$, we have

$$\begin{aligned} |d^*(fx, fy)| &= |\max\{6x, 6y\}| = |6x| \\ &\geq 2|\max\{3x, 3y\}| \\ &= 2|d^*(x, y)|. \end{aligned}$$

Clearly, (3.15) is satisfied and f is a self-surjection on $(-\infty, 0]$. In the view of Corollary 3.1, f and g has a unique common fixed point in $(-\infty, 0]$, indeed $f0 = g0 = 0$. Also

$$|d^*(fx, fy)| = |\max\{6x, 6y\}| = |6x| > \alpha|x| = \alpha|\max\{x, y\}| = \alpha|d^*(x, y)|,$$

for $1 < \alpha < 6$ and $\forall x, y \in (-\infty, 0]$ with $x \geq y$. Thus, f is a dualistic expanding surjective self-mapping on $(-\infty, 0]$. From, Corollary 3.3, $f0 = 0 \in (-\infty, 0]$ is unique.

Authors' Contributions

All authors contributed equally and significantly in writing this article. All authors read and approved the final manuscript.

Acknowledgment

The authors thank the anonymous referees for their remarkable comments, suggestions, and ideas that helped to improve this paper.

Conflicts of Interest

The authors declare that they have no conflicts of interest.

Disclaimer (Artificial intelligence)

Option 1:

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

Option 2:

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc have been used during writing or editing of manuscripts. This explanation will include the name, version, model, and source of the generative AI technology and as well as all input prompts provided to the generative AI technology

Details of the AI usage are given below:

- 1.
- 2.
- 3.

References

- [1]. I. Altun, F. Sola and H. Simsek, Generalized contractions on partial metric spaces, *Topology Appl.*, **157(18)** (2010), 2778-2785.
- [2]. M. Arshad, M. Nazam and I. Beg, Fixed point theorems in ordered dualistic partial metric spaces, *Korean J. Math.*, **24(2)** (2016), 169-179.
- [3]. P. Z. Daffer and H. Kaneko, On expansive mappings, *Math. Japonica*, **37**(1992), 733-735.
- [4]. R.D. Daheriya, R. Jain and M. Ughade, Some fixed-point theorem for expansive type mapping in dislocated metric space, *ISRN Mathematical Analysis*, 2012, 1-5, Article ID 376832, doi:10.5402/2012/376832.
- [5]. X. Huang, C. Zhu and X. Wen, Fixed point theorems for expanding mappings in partial metric spaces, *An. St. Univ. Ovidius Constanta*, **20(1)**, 2012, 213-224.
- [6]. R. Jain, R. D. Daheriya and M. Ughade, Fixed point, coincidence point and common fixed-point theorems under various expansive conditions in parametric metric spaces and parametric b-metric spaces, *Gazi University Journal of Science*, **29(1)** (2016), 95-107.
- [7]. S. G. Matthews, Partial Metric Topology, *Ann. New York Acad. Sci.* **728**(1994), 183-197.

- [8]. S. K. Mohanta and R. Maitra Coincidence points and common fixed points for expansive type mappings in cone b-metric spaces, *Applied Mathematics E-Notes*, **14**(2014), 200-208.
- [9]. M. Nazam, M. Arshad and M. Abbas, Some fixed-point results for dualistic rational contractions, *Appl. Gen. Topol.*, **17** (2016), 199-209.
- [10]. M. Nazam, M. C. Park and H. Mahmood, On a fixed-point theorem with application to functional equations, *Open Math.* **17** (2019), 1724–1736.
- [11]. M. Nazam, M. Arshad and C. Park, Fixed point theorems for improved α -Geraghty contractions in partial metric spaces, *J. Nonlinear Sci. Appl.*, **9**(2016), 4436–4449.
- [12]. M. Nazam and M. Arshad, Some fixed-point results in ordered dualistic partial metric spaces, *Trans. A. Razmadze Math. Institut.*, **172**(2018), 498–509.
- [13]. S. Oltra and O. Valero, Banach's fixed point theorem for partial metric spaces, *Rend. Ist. Mat. Univ. Trieste*, **36** (2004), 17-26.
- [14]. S. J. O'Neill: Partial metric, valuations and domain theory, *Ann. New York Acad. Sci.*, **806**(1996), 304-315.
- [15]. W. Shatanawi and F. Awawdeh, Some fixed and coincidence point theorems for expansive maps in cone metric spaces, *Fixed Point Theory and Applications*, **19**(2012), 1-10.
- [16]. N. Tas, and N.Y. Ozgur, On parametric S-metric spaces and fixed-point type theorems for expansive mappings, *Journal of mathematics*, (2016), Article ID 4746732, 1-6.
- [17]. S. Z. Wang, B.Y. Li, Z. M. Gao and K. Iseki, Some fixed-point theorems for expansion mappings, *Math. Japonica*, **29** (1984), 631-636.
- [18]. X. Wen and X. J. Huang: Common fixed-point theorem under contractions in partial metric spaces, *J. Comput. Anal. Appl.*, **13**(3) (2011), 583-589.
- [19]. Yan Han and Shaoyuan Xu, Some new theorems of expanding mappings without continuity in cone metric spaces, *Fixed Point Theory and Applications*, **3**(2013), 1-9.
- [20]. G.S. Jeong, B.E. Rhoades: Maps for which $F(T) = F(T^n)$, *Fixed Point Theory Appl.*, **6** (2005), 87–131.
- [21]. G. Jungck, S. Radenovic, S. Radojevic, V. Rakocevic: Common fixed-point theorems for weakly compatible pairs on cone metric spaces, *Fixed Point Theory Appl.*, 13 pages, doi:10.1155/2009/643840.
- [22]. G. Jungck, B.E. Rhoades: Fixed point theorems for occasionally weakly compatible mappings, *Fixed Point Theory*, **7** (2006), 287–296.
- [23]. G. Jungck: Commuting mappings and fixed points, *Amer. Math. Monthly*, **83** (1976), 261–263. 15
- [24]. G. Jungck: Compatible maps and common fixed points, *Internat. J. Math. Math. Sci.*, **9** (1986), 771–779.
- [25]. Z. Kadelburg, S. Radenovic, V. Rakocevi : Remarks on “Quasicontraction on a cone metric space”, *Appl. Math. Lett.*, **22** (2009), 1674– 1679.
- [26]. D. Wardowski: Fixed points of a new type of contractive mappings in complete metric spaces, *Fixed Point Theory and Applications*, Article number: **94**, (2012).
- [27]. M. Pireddu: Fixed points and chaotic dynamics for expansive contractive maps in Euclidean spaces, with some applications. Ph.D. Thesis. (2009), <https://arxiv.org/abs/0910.3832>.
- [28]. J. Gornicki: Fixed point theorems for F-expanding mappings, *Fixed Point Theory and Applications*, Article number: **9**, 2016.
- [29]. H. Aydi, H. K. Nashine, B. Samet, H. Yazidi: Coincidence and common fixed point results in partially ordered cone metric spaces and applications to integral equations. *Nonlinear Analysis*, **74**, (2010), 6814–6825.
- [30]. M. Ughade, S. D. Kushwaha, P. L. Sanodia, Fixed Point Theorems for Dualistic Expanding Mapping in Dualistic Partial Metric Spaces, *Jnanabha*, **51**(1) (2021), 42-48.