

Syntheses and X-ray structure of *N*-(Benzothiazol-2-yl)-3-chlorobenzamide

ABSTRACT

The reaction of 3-chloro-benzoyl chloride and 2-aminobenzothiazole in the presence of potassium thiocyanate yielded a yellowish solid $C_{15}H_{10}ClN_3OS_2$ (1). Reaction of 1 with cobalt chloride yield a yellowish solid $C_{14}H_{11}ClN_2O_2S$ (2) which gives crystal suitable for X-ray analyses. The compounds were characterized by elemental analyses FTIR spectroscopy and melting point measurement. Compound 2 crystallizes in the triclinic space group $P\bar{1}$ with the following parameters: $a = 7.0299(2) \text{ \AA}$ $b = 7.0466(2) \text{ \AA}$ $c = 14.5490(2) \text{ \AA}$ $\alpha = 103.844(3)^\circ$ $\beta = 92.929(2)^\circ$ $\gamma = 107.291(3)^\circ$ $V = 662.33(4) \text{ \AA}^3$ $Z = 2$ $R_1 = 0.030$ $wR_2 = 0.034$. The asymmetric unit of (2) consists of one molecule in which the 3-chloro phenyl moiety and the benzothiazole-2-yl moiety form a dihedral angle of $6.417(4)^\circ$. An intramolecular O—H...O hydrogen bond occurs. In the crystal O—H...N and N—H...O hydrogen bonds link the molecules forming layers parallel to the *ac* plane.

Key words: Cobalt(II)3-chloro-benzoyl chloride2-aminobenzothiazole dethiocyanation FTIR, X-ray diffraction.

1. INTRODUCTION

Thiourea derivatives can be synthesized with different methods [1-2] and uses in different applications such as the extraction of certain transition metals in acidic medium [3-5]. Multiple biological activities such as antibacterial [56] antifungal [6] antituberculosis [7] anticancer [8] antiviral [9] anticonvulsant [10] antioxidant [11] insecticidal [12] and herbicidal [13-14]. Benzoylthiourea derivatives are widely and successfully used in coordination chemistry. Indeed these compounds have C=O and C=S functions which can serve as donors in the chelation of metal ions [15-18]. Thiourea derivatives often have versatile behavior and are good intermediates in organic synthesis. In the presence of metal ions formation of heterocycles including the carbon atom of the C=S group or the oxygen atom of the C=O group can occur. In our recent work we reported triazolic and oxadiazolic compounds from 1-isonicotinoyl-4-phenylthiosemicarbazide in the presence of Fe^{2+} or Mn^{2+} ions respectively [19]. A heterocyclic compound in which the sulfur atom of the C=S group is included in the ring is reported in the literature [20]. On continuing our work in the synthesis coordination complexes from thiourea derivative the ligand (13-Benzothiazol-2-yl)-1-(3-chlorobenzoyl)thiourea was reacted with cobalt chloride (II). This reaction led to the dethiocyanation of the organic molecule to give *N*-(Benzothiazol-2-yl)-3-chlorobenzamide. In this article we report the synthesis and structural characterization by a spectroscopic study and by x-ray diffraction of the compounds.

2. METHODOLOGY

2.1 Starting materials and Instrumentation

“Potassium thiocyanate, 3-chloro-benzoyl chloride and 2-aminobenzothiazole were purchased from Sigma-Aldrich and used as received without further purification. All solvents used were of reagent grade. Elemental analyses of C H N S and Cl were recorded on a VxRio EL Instrument. Infrared spectra were obtained on a FTIR Spectrum Two of Perkin Elmer spectrometer in the 4000-400 cm^{-1} region” [21]. The compounds were synthesized following the procedure reported by Odame et al. [21 22] with slight modification.

2.2 Synthesis of 3-(13-Benzothiazol-2-yl)-1-(3-chlorobenzoyl)thiourea

The thiourea derivative was obtained by the reaction of potassium thiocyanate (1.9388 g 20 mmol) with 3-chloro-benzoyl chloride (3.3492 g 20 mmol) in 25 mL of acetone and heating under reflux for 2 h to yield the 3-chlorobenzoyl isothiocyanate. To the above solution was added a solution of 2-aminobenzothiazole (3 g 20 mmol) in 25 mL of acetone. The resulting mixture was heated overnight. The solvent was removed by evaporation and the crude product was recrystallized in methanol. Yield : 75.7 % . MP : 141-142 °C. Anal. calcd. for $\text{C}_{15}\text{H}_{10}\text{ClN}_3\text{OS}_2$: Calculated % C 51.79; % H 2.90; % N 12.08; % S 18.44; % Cl 10.19. Found C 51.81; % H 2.92; % N 12.03; % S 18.41; % Cl 10.16. FTIR: ($\nu \text{ cm}^{-1}$) : 3290 (N-H) 3055 (N-H) 1675 (C=O) 1592 (C=C) 1546 (C=C) 1451 (C-N) 1439 (C-N).


2.3 Synthesis of N-(Benzothiazol-2-yl)-3-chlorobenzamide

3-(13-Benzothiazol-2-yl)-1-(3-chlorobenzoyl)thiourea (0.3068 g 1 mmol) was dissolved in 20 mL of methanol and a solution of $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$ (0.2369 g 1 mmol) in 5 mL of methanol was added. The mixture was refluxed for 2 hours and filtered while hot. Slow evaporation of the resulting solution yield crystal suitable for X-ray analyses after a week. Yield : 80.3 % . MP : 155 °C. Anal. calcd. for $\text{C}_{14}\text{H}_{11}\text{ClN}_2\text{O}_2\text{S}$: . Calculated % C 54.81; % H 3.61; % N 9.13; % S 10.45; % Cl 11.56. Found % C 54.79; % H 3.58; % N 9.10; % S 10.39; % Cl 11.53. FTIR: ($\nu \text{ cm}^{-1}$) : 3280 (N-H amide) 3039 (C-H aromatic) 1702 (C=O) 1590 (C=C aromatic) 1450 (C-N amide) 1441 (C-N thiazole) 725 (C-Cl).

2.4 X-ray data collection structure determination and refinement

Crystals suitable for X-diffraction of the reported compound were grown by slow evaporation of MeOH solution of the complex. Details of the X-rays crystal structure solution and refinement are given in Table 1. Diffraction data were collected using a SuperNova Rigaku Oxford Diffraction diffractometer single diffractometer with graphite monochromatized $\text{MoK}\alpha$ radiation ($\lambda = 0.71073 \text{ \AA}$). “All data were corrected for Lorentz and polarization effects. No

absorption correction was applied. Complex scattering factors were taken from the program package *SHELXTL* [23]. The structures were solved by direct methods which revealed the position of all non-hydrogen atoms. All the structures were refined on F^2 by a full-matrix least-squares procedure using anisotropic displacement parameters for all non-hydrogen atoms [24]. The hydrogen atoms were geometrically optimized and refined as riding model by AFIX instructions. Molecular graphics were generated using *ORTEP-3* [25].



Scheme 1. General scheme for the synthesis of the compounds


Table 1. Crystal data and structure refinement details for (2).

Chemical formula	$\text{C}_{14}\text{H}_9\text{ClN}_2\text{OS} \cdot \text{H}_2\text{O}$
M_r	306.76
Crystal system space group	Triclinic <i>P</i> $\bar{1}$
Temperature (K)	100
a (Å)	7.0299 (2)
b (Å)	7.0466 (2)
c (Å)	14.5490 (4)
α	103.843 (2)

β	92.929 (2)
γ ($^\circ$)	107.291 (2)
V (\AA^3)	662.32 (3)
Z	2
Radiation type	Mo $K\alpha$
μ (mm^{-1})	0.45
Crystal size (mm)	0.22 \times 0.11 \times 0.07
$T_{\min} T_{\max}$	0.4991.000
Measured reflections	12431
Independent reflections	3437
Reflections [$ I > 2\sigma(I)$]	2964
R_{int}	0.030
R [$ I > 2\sigma(I)$]	0.034
wR_2	0.087
Goodness-of-Fit	1.06
Data/parameters/restraint	3437/184/0
$\Delta\rho_{\max} \Delta\rho_{\min}$ ($e \text{\AA}^{-3}$)	0.38–0.33

3. Results and discussion

In the present study the compound 3-(13-Benzothiazol-2-yl)-1-(3-chlorobenzoyl)thiourea (**1**) was synthesized by the reaction of 3-chlorobenzoyl isothiocyanate generated in situ and 2-aminobenzothiazole (Scheme 1). *N*-3-(13-Benzothiazol-2-yl)-1-(3-chlorobenzoyl)thiourea (**1**) was used to synthesize coordination complex owing to the presence of multiple donor sites such as O N and S atoms. (Benzothiazol-2-yl)-3-chlorobenzamide (**2**) was obtained during our attempt to synthesize complex using Co(II) salt. It seems that the metal cation act as catalyst for the dethiocyanation reaction. This phenomena was observed in analogous reactions. In fact in our previous work in attempt to synthesize Mn(II) and Fe(II) complexes using 1-isonicotinoyl-4-phenylthiosemicarbazidewe have isolated two news organic ligands 4-phenyl-3-(pyridin-4-yl)-1*H*-124-triazole-5(4*H*)-thione and *N*-phenyl-5-(pyridin-4-yl)-134-oxadiazol-2-amine hydrochloride respectively [19]. The reaction sequences depicted in the Scheme 1 were followed to obtain (**1**) and (**2**) respectively. A mechanism for the dethiocyanation and the formation of (**2**) is proposed in Scheme 2.



Scheme 2. Proposed mechanism for the formation of (2)

3.1 General study

The infrared spectrum of compound (1) exhibited characteristic bands. The bands of medium intensities appearing at 3290 cm^{-1} and 3055 cm^{-1} are attributed to N–H stretching vibrations of the amide and the thioamide groups[26]. The bands pointed at 1675 cm^{-1} is attributed to the C=O linked to the phenyl ring. Bands due to the aromatic units are pointed in the range $1592\text{-}1546\text{ cm}^{-1}$ [27]. The bands pointed at 1451 cm^{-1} and 1439 cm^{-1} are attributed to the C–N stretching vibrations[28]. Upon reaction of (1) and Co(II) salt compound (2) was recovered. On the infrared spectrum of compound (2) the band pointed at 3280 cm^{-1} is indicative of the presence of N–H of the amide group. The band due to the C=O moiety is shifted to high frequencies and is pointed at 1702 cm^{-1} . The presence of aromatic rings is confirmed by the bands which appear in the range $1590\text{-}1540\text{ cm}^{-1}$. Bands at 1450 cm^{-1} and 1441 cm^{-1} are attributed to the C–N stretching vibrations. Band due to C–Cl is pointed at 730 cm^{-1} [29].

3.2 Description of the structure of (2)

Compound (**2**) was obtained by mixing methanolic solution of Cobalt(II) chloride with methanol solution of 3-(13-Benzothiazol-2-yl)-1-(3-chlorobenzoyl)thiourea. After the reaction gradual evaporation of the filtered solution yields crystals of (**2**) suitable for X-ray diffraction. The title compound $C_{14}H_{11}ClN_2O_2S$ (**2**) crystallizes in the triclinic space group *P* $\bar{1}$. The crystallographic data selected bond lengths and bond angles for the crystal structure are provided in Table 2. The ORTEP diagram for compounds is presented in Fig. 1. The molecule consists of a 3-chlorophenyl moiety linked through a carbonyl unit to a 2-aminobenzothiazole moiety yielding the monohydrated *N*-(Benzothiazol-2-yl)-3-chlorobenzamide derivative. The benzothiazole-2-yl moiety is quite planar (rms deviations of fitted atoms of 0.0099 (3) Å) and form a dihedral angle of 6.417 (4)° with the 3-chlorophenyl moiety. The bond distance of O1–C8 is 1.2305 (17) Å which is consistent with a carbonyl whilst the S1–C1 bond distance value of 1.7504 (14) Å is consistent with double bond character. These value are comparable to those found for *N*-(benzothiazol-2-yl)benzamide [1.750 (1) Å for S–C and 1.222 (2) Å for O1–C2] [22]. The C7–N2 and C8–N2 bond lengths are 1.3867 (18) and 1.3657 (18) Å are shorter than the normal single C–N bond length indicating the sp^2 hybridization of the nitrogen atom [22–30]. The bond angle value of 122.67(12)° for C8–N2–C7 confirms the sp^2 hybridization of N2. The bond angle of C1–S1–C7 is 88.22 (7)° whilst the bond angle of S1–C7–N3 is 117.13 (11)°. The benzothiazolyl moiety is *cis* with respect to the carbonyl oxygen atom across the C8–N2 bond. The independent water molecule donates one intramolecular hydrogen bond to the carbonyl atom of the ligand [O2–H2B...O1] (Fig. 2, Table 3). Molecular layers running almost parallel to the *ac* plane are formed by intermolecular hydrogen bonds involving the second H atom of the independent water molecule group and the N atom of the benzothiazolyl moiety as acceptor [O2–H2C...N3ⁱ: $i = -x+1 -y+1 -z+1$] and NH groups and the oxygen atom of the water molecule as acceptor [N2–H2...O2ⁱⁱ: $ii = xy+1 z$] (Fig. 3, Table 3).




Figure 1. ORTEP plot (30% probability ellipsoids) showing the structure of (2).

Table 2. Selected geometric parameters (Å °) for (2).

S1—C1	1.7359 (14)	N3—C7	1.3018 (18)
S1—C7	1.7504 (14)	N2—C7	1.3867 (18)
O1—C8	1.2305 (17)	N2—C8	1.3657 (18)
N3—C6	1.3940 (18)	C13—C12	1.382 (2)
C1—S1—C7	88.22 (7)	N3—C7—N2	120.34 (12)
C7—N3—C6	109.67 (12)	O1—C8—N2	120.84 (13)
C8—N2—C7	122.67 (12)	O1—C8—C9	120.11 (12)
N3—C7—S1	117.13 (11)	N2—C8—C9	119.05 (12)

Table 3. Hydrogen-bonding geometry (Å °) of compound (2).

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
O2—H2B...O1	0.87	2.00	2.8542(15)	165.0
O2—H2C...N3 ⁱ	0.87	2.19	2.9472(16)	145.1
N2—H2...O2 ⁱⁱ	0.88	1.99	2.8541(16)	167.1

Symmetry codes: (i) $-x+1 -y+1 -z+1$; (ii) $xy+1 z$.




Figure 2. Intra and intermolecular hydrogen contacts showing the dimer in the crystal.




Figure 3. View along the *b* axis of the crystal packing of compound (2)

4. Conclusion

Compound (1) synthesized from the reaction between 2-aminothiazole and 3-chlorobenzoyl isothiocyanate yields compound (2) when it reacts with Co(II). It appears that the action of the metal is determinate in the dethiocyanation reaction. The structures of compounds (1) and (2) were confirmed by elemental analysis and spectroscopic techniques. The molecular structure of compound (2) is determined by X-ray diffraction.

SUPPLEMENTARY MATERIALS

CCDC-2340894 contains the supplementary crystallographic data for this paper. These data can be obtained free of charge via <https://www.ccdc.cam.ac.uk/structures/> or by contacting The Cambridge Crystallographic Data Centre 12 Union Road Cambridge CB2 1EZ UK.

REFERENCES

1. Aydın F, Aykaç D, Ünver H, İskeleli NO. Synthesis spectral properties and structure of new novel 3,3'-dibenzoyl-1,1'-(propan-1,3-diyl)-bisthiourea. *J. Chem. Crystallogr.* 2012; 42(4):381–387.
Available:<https://doi.org/10.1007/s10870-011-0258-5>
2. Samb I, Gaye N, Sylla-Gueye R, Thiam EI, Gaye M, Retailleau P. Crystal structure of *N,N*-[(ethane-1,2-diyl)bis(azanediylcarbonothioyl)]bis(benzamide). *Acta Crystallogr.* 2019:E75(5):642–645.
Available:<https://doi.org/10.1107/S205698901900495X>
3. Kuchekar SR, Bhumkar SD, Aher HR, Han SH. separation of platinum(IV) from pharmaceuticals using p-methylphenyl thiourea by solvent extraction: separation from palladium(II) nickel (II). *Anal. Chem. Lett.* 2019;9(6):775–788.
Available:<https://doi.org/10.1080/22297928.2020.1712236>
4. Uçak ŞŞ, Aydın A. A novel thiourea derivative for preconcentration of copper(II) nickel(II) cadmium(II) lead(II) and iron(II) from seawater samples for flame atomic absorption spectrophotometry. *Mar. Pollut. Bull.* 2022;180:113787.
Available:<https://doi.org/10.1016/j.marpolbul.2022.113787>
5. Huang X, Jin K, Zhang R, Gong Y, Zeng J, Zhang R, Liu Y, Xue J. Selective solvent extraction of Cu(II) from aqueous solutions using an acyl-based thiourea: Extraction study and DFT analysis of reaction mechanism. *Hydrometallurgy* 2024;223:106226.
Available:<https://doi.org/10.1016/j.hydromet.2023.106226>
6. Antypenko L, Meyer F, Kholodniak O, Sadykova Z, Jirásková T, Troianova A, Buhaiova V, Cao S, Kovalenko S, Gabe L-A, Steffens KG. Novel acyl thiourea derivatives: synthesis antifungal activity gene toxicity drug-like and molecular docking screening. *Arch. Pharm.* 2019;352(2):1800275.
Available:<https://doi.org/10.1002/ardp.201800275>
7. Doğan ŞD, Gündüz MG, Doğan H, Krishna VS, Lherbet C, Sriram D. Design and synthesis of thiourea-based derivatives as mycobacterium tuberculosis growth and enoyl acyl carrier protein reductase (InhA) inhibitors. *Eur. J. Med. Chem.* 2020;199:112402.
Available:<https://doi.org/10.1016/j.ejmech.2020.112402>
8. Arafa WAA, Ghoneim AA, Mourad AK. N-Naphthoyl Thiourea derivatives: an efficient ultrasonic-assisted synthesis reaction and in vitro anticancer evaluations. *ACS Omega* 2022;7(7):6210–6222.
Available:<https://doi.org/10.1021/acsomega.1c06718>

9. Krishna Reddy RC, Rasheed S, Subba Rao D, Adam S, Venkata Rami Reddy Y, Raju C. N. New urea and thiourea derivatives of piperazine doped with febuxostat: synthesis and evaluation of anti-TMV and antimicrobial activities. *Sci. World J.* 2013; 2013:682603.
Available:<https://doi.org/10.1155/2013/682603>
10. Thakur AS, Deshmukh R, Jha AK, Kumar PS. Molecular docking study and anticonvulsant activity of synthesized 4-((4,6-dimethyl-6*H*-1,3-thiazin-2-yl)phenylsulfonyl)urea/thiourea derivatives. *J. King Saud Univ. Sci.* 2018;30(3):330–336.
Available:<https://doi.org/10.1016/j.jksus.2016.12.006>
11. Naz S, Zahoor M, Umar MN, Alghamdi S, Sahibzada MUK, UIBari W. Synthesis characterization and pharmacological evaluation of thiourea derivatives *Open Chem.* 2020;18(1):764–777.
Available:<https://doi.org/10.1515/chem-2020-0139>
12. Han Z-Y, Wu W-Y, Chen F-L, Guan X-L, Fu X-H, Wan R. Design synthesis crystal structure and insecticidal evaluation of novel arylpyrazole derivatives containing cyhalothroyl thiourea moiety. *Phosphorus Sulfur Silicon Relat. Elem.* 2017;192(8):911–918.
Available:<https://doi.org/10.1080/10426507.2017.1292272>
13. Arslan H, Duran N, Borekci G, Koray Ozer C, Akbay C. Antimicrobial activity of some thiourea derivatives and their nickel and copper complexes. *Molecules* 2009;14(1):519–527.
Available:<https://doi.org/10.3390/molecules14010519>
14. Cunha S, Macedo F, Costa GAN, Rodrigues MT, Verde RBV, de Souza Neta LC, Vencato I, Lariucci C, Sá FP. Antimicrobial activity and structural study of disubstituted thiourea derivatives. *Monatsh. Chem.* 2007;138(5):511–516.
Available:<https://doi.org/10.1007/s00706-007-0600-y>
15. Binzet G, Külcü N, Flörke U, Arslan H. Synthesis, and characterization of Cu(II) and Ni(II) complexes of some 4-bromo-*N*-(di(alkyl/aryl)carbamoithiroyl) benzamide derivatives. *J. Coord. Chem.* 2009;62(21):3454–3462.
Available:<https://doi.org/10.1080/00958970903082200>
16. Kavak G, Özbey S, Binzet G, Külcü N. Synthesis and single crystal structure analysis of three novel benzoylthiourea derivatives. *Turk. J. Chem.* 2009;33(6):857-868.
Available:<https://doi.org/10.3906/kim-0901-1>
17. Vivas-Reyes R, Espinosa-Fuentes E, Forigua J, Arias A, Gaitán R, Arguello E. Theoretical study of a series of *N*-(*N*-propyl)-*N'*-(para-*R*-benzoyl)-thioureas with trans[Pt(Py)₂Cl₂] through chemistry reactivity descriptors based on density functional theory. *J. Mol. Struct.: THEOCHEM* 2008;862(1):92–97.

Available:<https://doi.org/10.1016/j.theochem.2008.04.036>

18. Binzet G, Kavak G, Külcü N, Özbey S, Flörke U, Arslan H. Synthesis and Characterization of novel thiourea derivatives and their nickel and copper complexes. *J. Chem.* 2013;2013:536562.
Available:<https://doi.org/10.1155/2013/536562>
19. Faye N, Mbow B, Gaye AA, Ndoye C, Diop M, Excoffier G, Gaye M. Syntheses & Antioxidant activity of 1-isonicotinoyl-4-phenylthiosemicarbazide and crystal structures of *N*-phenyl-5-(pyridin-4-yl)-1,3,4-oxadiazol-2-amine hydrochloride and 4-Phenyl-3-(pyridin-4-yl)-1*H*-1,2,4-triazole-5(4*H*)-thione derived from 1-isonicotinoyl-4-phenylthiosemicarbazide. *Earthline J. Chem. Sci.* 2022;9(2):189–208.
Available:<https://doi.org/10.34198/ejcs.9223.189208>
20. Kalhor M, Salehifar M, Nikokar I. Synthesis characterization and antibacterial activities of some novel *N,N*-disubstituted thiourea 2-amino thiazole and imidazole-2-thione derivatives. *Med. Chem. Res.* 2014;23(6):2947–2954.
Available:<https://doi.org/10.1007/s00044-013-0883-y>
21. Odame F, Hosten E, Krause J, Isaacs M, Hoppe H, Khanye SD, Sayed Y, Frost C, Lobb K, Tshentu Z. Synthesis characterization and biological activity of some dithiourea derivatives. *Acta Chim. Slovenica* 2020;67(3):764–777.
Available:<https://doi.org/10.17344/acsi.2019.5689>
22. Odame F, Woodcock G, Hosten EC, Lobb K, Tshentu ZR. A novel gold(I)-mediated intramolecular transamidation of benzoyl thiourea derivatives to form benzamides via dethiocyanation. *J. Organomet. Chem.* 2020;922:121359.
Available:<https://doi.org/10.1016/j.jorganchem.2020.121359>
23. Sheldrick GM. SHELXT—Integrated space-group and crystal-structure determination. *Acta Crystallogr.* 2015;A71(1):3–8.
Available:<https://doi.org/10.1107/S2053273314026370>
24. Sheldrick GM. Crystal structure refinement with SHELXL. *Acta Crystallogr.* 2015;C71(1):3–8.
Available:<https://doi.org/10.1107/S2053229614024218>
25. Farrugia LJ. WinGX and ORTEP for Windows: an update. *J. Appl. Crystallogr.* 2012;45(4):849–854.
Available:<https://doi.org/10.1107/S0021889812029111>
26. González DLN, Saeed A, Shabir G, Flörke U, Erben MF. Conformational and crystal structure of acyl thiourea compounds: The case of the simple (2,2-dimethyl-propionyl) thiourea derivative. *J. Mol. Struct.* 2020;1215:128227.
Available:<https://doi.org/10.1016/j.molstruc.2020.128227>

27. Moradi P, Kikhavani T, Abbasi Tyula Y. A new samarium complex of 13-bis(pyridin-3-ylmethyl)thiourea on boehmite nanoparticles as a practical and recyclable nanocatalyst for the selective synthesis of tetrazoles. *Sci. Rep.* 2023;13(1):5902.
Available:<https://doi.org/10.1038/s41598-023-33109-y>
28. Alcívar León CD, Echeverría GA, Piro OE, Ulic SE, Jios JL, Luna Tapia CA, Mera Guzmán MF. New thiourea and urea derivatives containing trifluoromethyl- and bis-trifluoromethyl-4*H*-chromen-3-yl substituents. *Mol. Phys.* 2019;117(3):368–381.
Available:<https://doi.org/10.1080/00268976.2018.1514132>
29. Gumus I, Solmaz U, Binzet G, Keskin E, Arslan B, Arslan H. Supramolecular self-assembly of new thiourea derivatives directed by intermolecular hydrogen bonds and weak interactions: crystal structures and Hirshfeld surface analysis. *Res. Chem. Intermed.* 2019;45(2):169–198.
Available:<https://doi.org/10.1007/s11164-018-3596-5>
30. Avşar G, Arslan H, Haupt H-J, Külcü N. Crystal Structure of cis-bis(*N,N*-dimethyl-*N'*-benzoylthiourea) palladium(II). *Turk. J. Chem.* 2003;27(3):281–286.
Available:<https://journals.tubitak.gov.tr/chem/vol27/iss3/1>