

## Original Research Article

# Factors Associated with The Incident of Hypertension in Outpatient Patients at The Kramat Jati District Health Center, East Jakarta, 2024

### ABSTRACT

**Aims:** This research aims to determine the factors that most cause the incidence of hypertension in outpatients at the Kramat Jati District Health Center, East Jakarta in 2024.

**Study design:** *Cross-Sectional Design*

**Place and Duration of Study:** This research was carried out at the Non-Communicable Diseases Polyclinic or Hypertension Polyclinic, Kramat Jati District Health Center, East Jakarta. Data collection took place for 2 weeks in February 2024.

**Methodology:** The selected population is outpatients who have hypertension and are active as controls and non-hypertensive outpatients who visit the Kramat Jati District Health Center, East Jakarta. The sample determined was outpatients at the Kramat Jati District Health Center, East Jakarta. Samples were collected using non-random accidental sampling technique and 101 research samples were obtained. Data Processing and Analysis using SPSS (Statistical Package for the Social Sciences) 29th version was used to view research results using univariate and bivariate analysis. Univariate analysis.

**Results:** The results obtained showed that 81 respondents (80.2%) experienced hypertension. The dominant characteristics of hypertensive patients were found to be women (65.3%), aged 56-65 years (28.7%), patients who had a family history of hypertension (52.5%), patients who consumed  $\geq 5$  grams of sodium (70.3%), patients who did physical activity (76.2%), and patients who did not smoke (68.3%). The factors of age (p-value = 0.0001), history of hypertension (p-value = 0.046), sodium diet (p-value = 0.0001), and smoking (p-value = 0.039) are significantly related and most dominantly influence the incidence of hypertension. The factors of gender (p-value = 0.410) and physical activity (p-value = 0.454) were not significantly related to increasing blood pressure.

**Conclusion:** There is a significant relationship between factors that cannot be changed, namely age and family history of hypertension, with the incidence of hypertension at the Kramat Jati District Health Center, East Jakarta, and there is also a significant relationship between factors that can be changed, namely dietary sodium patterns and smoking activities, and the incidence of hypertension in Kramat Jati District Health Center, East Jakarta.

*Keywords: Hypertension, Risk Factors, Relationship, Patients*

### 1. INTRODUCTION

High blood pressure is the result of measuring a person's blood pressure with a systolic value of 140 mmHg or more or a diastolic value exceeding 90 mmHg [1]. Blood pressure increases gradually over time, such as consuming excess sodium, insufficient potassium intake, overweight and obesity, drinking alcohol, and lack of physical activity [2]. Damage to the cardiovascular system such as stroke, myocardial infarction, heart failure, and even death is

closely related to the incidence of hypertension experienced by the elderly. In the elderly, the adverse impacts that occur are strongly influenced by changes in organ systems such as arterial stiffness, mechanical, neurohormonal, and autonomic hemodynamic changes, dysregulation, and decreased function of the kidneys [3]. Hypertension is classified into two based on its causes, some can be classified by etiology and others cannot be classified. classified according to etiology [4]. The prevalence of hypertension is high in low and middle-income countries, cases of hypertension are more common in men, namely around 31.9%, compared to women, which is around 30.1%. Hypertension in men is highest in Eastern European and Central Asian countries at 39.0%, in women cases of hypertension are highest in sub-Saharan African countries at 36.3% [5].

According to the World Health Organization (WHO), around the world there are 1.13 billion people who suffer from high blood pressure and it is estimated that this will increase to 1.5 billion in 2025 [6] The hypertensive population in the Asian continent, such as in Malaysia, is 33.1%, in China is 31.5% and in India it is 28.5% [7] The increase in hypertension cases has doubled from 1990-2019 in the population aged 30-79 years, which initially numbered 317 million in men to 652 million. while women initially numbered 331 million to 626 million in 2019 [8] Risesdas research in Indonesia said the incidence of hypertension with a minimum age of 18 years in the Bangka Belitung region was 30.9%, in the Papua region it was 16.8%, in Bali province, there were 19, 9% of cases of hypertension [9]. According to a survey conducted in 2018, it was found that 2,886 cases, or 10.45% of the total population living in East Jakarta experienced high blood pressure [10].

The hypertension factor does not stand alone, hypertension can occur due to the habit of consuming alcohol, smoking habits, consuming salt, consuming red meat, physical activity, and obesity, it can also be due to heredity, age, and gender differences. Someone who does not consume red meat has a 77% lower chance of developing hypertension, similarly, someone who does not consume alcohol has a 70% lower chance of developing hypertension and those who have a BMI  $\geq 25$  have a 3.05 times greater chance of developing hypertension [11]. Other factors, such as genetics or an adverse intrauterine environment such as gestational hypertension or pre-eclampsia have a small but strong association with high blood pressure levels in adulthood [12].

Many factors can influence the occurrence of hypertension, plus the high incidence of hypertension has made the Indonesian government have a new goal, namely reducing the incidence of hypertension in Indonesia through the BPJS program known as the Chronic Disease Management Program or PROLANIS. This series of management focuses on non-communicable phenomena such as hypertension [18].

Based on the explanation and data above, it was found that many factors support the occurrence of increased blood pressure. So the author is interested in carrying out research related to factors related to the incidence of hypertension in outpatients at the Kramat Jati District Health Center, East Jakarta in 2024

### **Research Problem**

The formulation of the problem based on the background obtained is "What factors are related to the incidence of hypertension in outpatients at the Kramat Jati District Health Center, East Jakarta in 2024"

### **Research Purposes**

To find out the factors that most cause the incidence of hypertension in outpatients at the Kramat Jati District Health Center, East Jakarta in 2024

## **2. MATERIAL AND METHODS**

### **2.1. Research Design**

The research was obtained using cross-sectional analytical techniques using simultaneous measurements to determine the dynamics of the relationship between the dependent variable, namely the incidence of hypertension, and the independent variables, namely genetic factors

that cannot be changed, such as age, gender and family history of hypertension, as well as lifestyle factors that can be changed, such as salt consumption, physical activity, smoking.

## **2.2. Location and Time of Research**

This research was carried out at the Non-Communicable Diseases Polyclinic or Hypertension Polyclinic, Kramat Jati District Health Center, East Jakarta. Data collection took place for 2 weeks in February 2024.

## **2.3. Research Subjects**

### **2.3.1 Population**

The selected population was outpatients who had hypertension and were active as controls and non-hypertensive outpatients who visited the Kramat Jati District Health Center, East Jakarta. The population was found to be 136 people.

### **2.3.2. Research Sample**

The sample was selected from the population who were willing to participate and met the inclusion criteria. Inclusion and exclusion criteria are as follows:

#### **Inclusion Criteria**

1. Outpatients at the Non-Communicable Diseases Polyclinic or Hypertension Polyclinic, Kramat Jati District Health Center, East Jakarta.
2. Adults to the elderly, namely patients aged  $\geq 26$  years to  $\geq 65$  years.
3. The patient is willing to be a sample in the research.
4. Compos mentis awareness, cooperative ability to listen and respond.

#### **Exclusion Criteria**

1. Do not agree or do not fill out the informed consent form.
2. Not an outpatient at the Kramat Jati District Health Center, East Jakarta

### **2.3.3. Sampling Technique**

The sample determined was outpatients at the Kramat Jati District Health Center, East Jakarta. Samples were collected using a non-random accidental sampling technique where the sampling technique was planned by the researcher and not random.

## **2.4. Data Collection**

Primary data was directly obtained from research subjects using a questionnaire sheet to determine the characteristics and lifestyle of outpatients which influence the incidence of hypertension.

## **2.5. Data Processing and Analysis**

SPSS (Statistical Package for the Social Sciences) 29th version was used to view research results using univariate and bivariate analysis. Univariate analysis in this study looked at the frequency of each existing independent variable, while bivariate analysis was used to see whether or not there was a relationship between the incidence of hypertension and the independent variables.

## **3. RESULTS AND DISCUSSION**

### **3.1. Research Results**

This research uses univariate analysis to see the frequency of each variable and bivariate analysis to describe whether or not there is a relationship between the two variables.

#### **3.1.1. Univariate Analysis**

This analysis is also called descriptive statistics. The purpose of this analysis is to describe the conditions of the phenomenon under study. The following are the results of the univariate analysis for each variable:

##### **3.1.1.1. Age**

**Table 1. Frequency Distribution of Respondents Based on Age**

<b>Age</b>	<b>Frequency</b>	<b>Percentage (%)</b>
26-35	5	5.0
36-45	12	11.9

46-55	27	26.7
56-65	29	28.7
>65	28	27.7
<b>Total</b>	<b>101</b>	<b>100.0</b>

Based on table 1 above, this research shows that the age variable consists of 5 or 5% of respondents aged 26-35 years, 12 or 11.9% of respondents aged 36-45 years, 27 or 26.7% of respondents aged 46-55 years, 29 or 28.7% of respondents were aged 56-65 years and 28 or 27.7% of respondents were aged >65 years.

### 3.1.1.2. Gender

**Table 2. Frequency Distribution of Respondents Based on Gender**

Gender	Frequency	Percentage (%)
Female	66	65.3
Male	35	34.7
<b>Total</b>	<b>101</b>	<b>100.0</b>

Based on table 2 above, this research shows that the respondent gender variable is known to consist of 66 or 65.3% female respondents and 35 or 34.7% male respondents.

### 3.1.1.3. Family history of hypertension

**Table 3. Frequency Distribution Based on Family History of Hypertension**

History of Hypertension	Frequency	Percentage (%)
No	48	47.5
Yes	53	52.5
<b>total</b>	<b>101</b>	<b>100.0</b>

Based on table 3 above, this research shows that the variable history of hypertension in the respondent's family is known to consist of 48 or 47.5% of respondents who have no history and 53 or 52.5% of respondents who have a history.

### 3.1.1.4. Diet Sodium

**Table 4. Frequency Distribution of Respondents Based on Sodium Diet**

Diet Sodium	Frequency	Percentage (%)
<5 gram	30	29.7
≥5 gram	71	70.3
<b>Total</b>	<b>101</b>	<b>100.0</b>

Based on table 4 above, this research shows that the sodium diet variable shows that respondents consist of 30 or 29.7% of respondents who consume <5 grams of salt and 71 or 70.3% of respondents who consume ≥5 grams of salt.

### 3.1.1.5. Physical Activity

**Table 5. Frequency Distribution of Respondents Based on Physical Activity**

Physical Activity	Frequency	Percentage (%)
No	24	23.8
Yes	77	76.2
<b>Total</b>	<b>101</b>	<b>100.0</b>

Based on table 5 above, this research shows that the physical activity variable shows that respondents consist of 24 or 23.8% of respondents who do not do physical activity and 77 or 76.2% of respondents who do physical activity.

### 3.1.1.6. Smoking History

**Table 6. Frequency Distribution of Respondents Based on Smoking History**

Smoking	Frequency	Percentage (%)
No	69	68.3
Yes	32	31.7
<b>Total</b>	<b>101</b>	<b>100.0</b>

Based on Table 6 above, this research shows that the respondent smoking history variable is known to consist of 69 or 68.3% of respondents who are not active smokers and 32 or 31.7% of respondents who are active smokers.

### 3.1.1.7. Hypertension Status

**Table 7. Frequency Distribution of Respondents Based on the Incident of Hypertension**

Hypertension Status	Frequency	Percentage (%)
Not Hypertension	20	19.8
Hypertension	81	80.2
<b>Total</b>	<b>101</b>	<b>100.0</b>

Based on Table 7 above, this study shows that the hypertension variable is known to respondents consisting of 20 or 19.8% of respondents who do not experience hypertension and 81 or 80.2% of respondents who experience hypertension.

## 3.1.2. Bivariate Analysis

This analysis is used to find out whether there is a relationship between the two variables. The following are the results of the research analysis

### 3.1.2.1. Relationship between Age and the Incident of Hypertension

**Table 8. Relationship between Age and the Incident of Hypertension**

Age	Hypertension		Total	Asymptotic significance (2- sided)	Odds Ratio
	No	Yes			
26-35	5	0	5	0.0001	10.571
36-45	5	7	12		
46-55	5	22	27		
56-65	3	26	29		
>65	2	26	28		

Based on Table 8 of the Chi-Square test for the age variable, it is known that the test used is for the 5x2 table, using Pearson Chi-Square, namely 0.0001 ( $<0.05$ ), which means there is a significant relationship between the incidence of hypertension and each age category. Inferential tests obtained on the age variable on the incidence of hypertension showed that the Odds Ratio value was 10.571, which means that respondents aged >45 years had a 10.5 times greater risk of developing hypertension.

### 3.1.2.2. Relationship between gender and the incidence of hypertension

**Table 9. Relationship between gender and the incidence of hypertension**

Gender	Hypertension		Total	Asymptotic significance (2- sided)	Odds Ratio
	No	Yes			
Female	11	55	66	0.410	0.578

Based on table 9 of the ChiSquare test for the gender variable, it is known that the test used was for a 2x2 table and no Expected value was found <5, so using Continuity Correction the value was 0.410 (>0.05), which means there is no significant relationship between the incidence of hypertension and respondents who are male or female. In the gender variable regarding the incidence of hypertension, it is known that the Odds Ratio value is 0.578, which means that female respondents have a higher risk of developing hypertension, namely 0.57 times compared to men.

**3.1.2.3. Relationship between family history of hypertension and the incidence of hypertension**

**Table 10. Relationship between family history of hypertension and the incidence of hypertension**

History of Hypertension	Hypertension		Total	Asymptotic significance (2- sided)	Odds Ratio
	No	Yes			
No	14	34	48	0.046	3.225
Yes	6	47	53		

Based on table 10 of the ChiSquare test for the family history of hypertension variable, it is known that the test used was for a 2x2 table and no Expected value was found <5, so using Continuity Correction the value was 0.046 (<0.05), which means there is a significant relationship between the incidence of hypertension with respondents who have a family history of hypertension. In the family history of hypertension variable, it is known that the Odds Ratio value is 3.225, which means that respondents with a family history of hypertension are at 3.2 times greater risk of developing hypertension.

**3.1.2.4. Relationship between dietary sodium patterns and the incidence of hypertension**

**Table 11. Relationship between dietary sodium patterns and the incidence of hypertension**

Dietary Sodium Patterns	Hypertension		Total	Asymptotic significance (2- sided)	Odds Ratio
	No	Yes			
< 5 gram	13	17	30	0,0001	6.992
≥ 5 gram	7	64	71		

Based on Table 11 of the Chi-Square test for the sodium diet variable, it is known that the test used was for a 2x2 table and no Expected value <5 was found, so using Continuity Correction the value was 0.0001 (<0.05), which means there is a significant relationship between events. hypertension with respondents who consumed sodium in amounts <5 grams or consumed sodium in amounts ≥5 grams. The inferential test on the sodium dietary pattern variable on the incidence of hypertension shows that the Odds Ratio value is 6.992, which means that respondents who consume sodium in amounts ≥5 grams/day are 6.9 times more likely to suffer from hypertension.

**3.1.2.5. Relationship between physical activity and hypertension**

**Table 12. Relationship between physical activity and hypertension**

Physical Activity	Hypertension		Total	Asymptotic significance (2- sided)	Odds Ratio
	No	Yes			
No	4	20	24	0,454	0,763
Yes	16	61	77		

Berdasarkan tabel 3.12 dari uji ChiSquare variabel aktivitas fisik diketahui bahwa uji yang digunakan adalah untuk tabel 2x2 dan ditemukan nilai Expepected <5, maka menggunakan Fisher Exact Test yaitu 0,454 (>0,05), yang artinya tidak terdapat hubungan signifikan antara kejadian hipertensi dengan responden yang melakukan aktivitas fisik. Pada variabel ini diketahui nilai Odds Ratio sebesar 0,763 yang berarti responden yang beraktivitas mempunyai potensi 0,76 kali lebih besar mengalami hipertensi.

### 3.1.2.6. Relationship between smoking activities and the incidence of hypertension

**Table 13. Relationship between smoking activities and the incidence of hypertension**

Smoking Activities	Hypertension		Total	Asymptotic significance (2- sided)	Odds Ratio
	No	Yes			
No	18	51	69	0,039	5,294
Yes	2	30	32		

Based on table 13 of the ChiSquare test for the smoking activity variable, it is known that the test used was for the 2x2 table and no Expepected value <5 was found, so we used Continuity Correction, namely 0.039 (<0.05), which means there is a significant relationship between the incidence of hypertension and the respondent who ever smoked actively or never smoked actively. In the smoking history variable on the incidence of hypertension, it is known that the Odds Ratio value is 5.294, which means that respondents who are exposed to cigarette smoke have a 5.2 times higher risk of developing hypertension.

## 3.2. Discussion

### 3.2.1. Relationship between Age and the Incident of Hypertension

In this study, it was found that 5% of subjects were aged 26 to 35 years, 11.9% of subjects were aged 36 to 45 years, 26.7% of subjects were aged 46 to 55 years, 28.7% of subjects were aged 56 to 65 years and 27.7% of subjects were over 65 years old. There is a significant relationship between the incidence of hypertension and each age category with the result of Asymptotic significance (2-sided), namely 0.0001 (<0.05). Research on the incidence of hypertension found that subjects over 45 years of age had a 10.5 times greater risk than respondents under 45 years of age. This research is in line with Cohen et al's research that the 40-44-year age group has the highest risk of developing hypertension because the human body becomes less responsive to stimuli as age increases, causing reduced sympathetic regulation in older individuals [20]. The results of Rahmawati et al's research are inconsistent, showing that those aged 20 to 40 years are more likely to experience hypertension due to unhealthy lifestyle habits such as not liking activities, smoking continuously, malnutrition, stress, and abnormalities in imaging results in the heart and brain which increase cardiovascular events in the elderly [21]. Age is calculated from the time a person is born until their birthday, where the older a person is, the more mature their thoughts and behavior become [22]. As age increases, there is also a process of dysfunction where there is a decrease in the regulation of repair mechanisms, thereby increasing the possibility of disease and causing an increase in the death rate [23].

### **3.2.2. Relationship between gender and the incidence of hypertension**

In this study, it was found that 65.3% of the subjects were female and 34.7% were male. According to the relationship, a value of 0.410 ( $>0.05$ ) was obtained, which means there is no significant relationship between the incidence of hypertension and respondents who are male or female. In the gender variable, it was found that female respondents had a higher risk of developing hypertension, namely 0.57 times compared to men.

Research by Mouhtadi et al agrees with this research where it is stated that hypertension attacks women the most because women leave the house less often and do not exercise as much, so they are at risk of developing hypertension [24]. Kusumawaty et al.'s research also agrees with the current study which shows that the incidence of hypertension in women is higher, namely from 92 samples found 54 women had hypertension and 38 men had hypertension. Kusumatay et al said that gender is closely related to the occurrence of hypertension because women suffer from hypertension during menopause due to a decrease in the hormone estrogen so they cannot maintain levels of High-Density Lipoprotein whose function is a protective factor in preventing atherosclerosis and blood vessel damage [25]. Gender is a division of subjects that is distinguished from the structure of chromosomes, namely the structure of the cell nucleus, where genes are carriers of inherited traits. Men and women can be differentiated based on two different chromosomes, women have both X sex chromosomes, for men the two chromosomes are different, namely the X chromosome and the Y chromosome. Female genetic carriers are the X chromosome type and the Y chromosome in men [26].

### **3.2.3. Relationship between family history of hypertension and the incidence of hypertension**

This study shows that the variable history of hypertension in the family of respondents is known to consist of 48 or 47.5% of respondents who have no history and 53 or 52.5% of respondents who have a history. According to the relationship, there is a significant relationship between the incidence of hypertension and respondents who have a family history of hypertension. In this study, it was also found that respondents with a family history of hypertension were 3.2 times more likely to develop hypertension. Li et al's research agrees with this research where it was found that individuals with a family history of hypertension had an odds ratio of 4.103, which means that people with a family history of hypertension are 4.1 times more likely to develop hypertension [27].

Liu et al's research also agrees with this research which shows that patients with a family history of hypertension are 2-4 times more likely to develop hypertension. A study by Liu et al found that the relationship between the risk of hypertension based on a family history of hypertension was stronger in women than in men and this relationship appeared to occur more frequently in the first generation [28]. According to research by Rodríguez-Moran et al, it shows that a family history of hypertension inherited from the mother is associated with hyperinsulinemia (OR 1.5), which means a 1.5-fold increase in risk of developing hypertension, high blood pressure, and hypertension (OR 4.0), which means a 4-fold increase in risk. high levels of hypertension, hypertriglyceridemia (OR 1.6) which means a 1.6 times higher risk of hypertension, and low high-density lipoprotein cholesterol (OR 1.3) which means a 1.3 times higher risk of developing hypertension. These researchers concluded the hypothesis that cardiovascular risk phenotypes are inherited from the mother via mitochondria is still unclear and the mechanisms underlying gender-related differences require further research [29].

Maternal factors such as food intake, central adiposity, and placental insufficiency during pregnancy contribute significantly to the programming of disease phenotypes in the offspring. Fetal growth is determined by the interaction between the fetal genome and its environment, which is determined by the mother's environment as well as the physiology of the mother and placenta. Many underlying assumptions have a big impact on overcoming metabolic and cardiovascular diseases. So further research is still needed to determine the evolution,

etiology, and associated risk factors [30]. The genome also has an influence on the occurrence of hypertension, when one gene co-segregates with one trait in a family, this results in the gene being linked to that trait and there is likely a causal relationship. 31 There is a serine proteinase encoded by the gene which functions to inhibit the activity of tissue type I plasminogen activator. The encoded protein forms the bulk of lipoprotein(a) and then through a proteolytic process is cleaved and produces fragments that attach to atherosclerotic lesions and cause thrombogenesis. According to research by Antonicelli et al, Lp (a) and peroxidative stress may be involved as cofactors in essential hypertension [32]. A family history of hereditary hypertension is a condition of hypertension that originates in the family, if it is not inherited from the family then it cannot be said to be a family history. Genetic factors do not completely influence the history of hypertension, but other factors support it. High blood pressure is often lowered but we can avoid it by controlling blood pressure regularly and maintaining a healthy pattern [33].

#### **3.2.4. Relationship between dietary sodium patterns and the Incidence of hypertension**

In this study, it was found that 29.7% of subjects consumed <5 grams of sodium per day, and it was also found that 70.3% of subjects consumed ≥5 grams of sodium per day. According to the relationship, there is a relationship between subjects who consume sodium in amounts <5 grams or ≥5 grams and the incidence of hypertension with the result of Asymptotic significance (2-sided) namely 0.0001 (<0.05). In this study, it was found that subjects who consumed ≥5 grams of sodium per day had a 6.9 times greater chance of developing hypertension than subjects who consumed <5 grams of sodium per day. Pertiwi et al's research agree with this research, where hypertensive patients who consumed high amounts of salt in healthcare facilities experienced a large increase in systolic blood pressure at each visit, p-value = 0.038 (<0.05) [34].

Farapti et al's research disagrees because it shows there is no correlation between salt intake and blood pressure, but a significant correlation was found between Na/K ratio levels, not with sodium and potassium levels separately. Blood pressure increases 3.99 times for every 1 unit increase in the Na/K ratio in the urine [35]. Wahyu et al's research shows that people with hypertension tend to reduce sodium intake to prevent disease complications. This eating pattern will influence the relationship between sodium intake and the incidence of hypertension. This relationship may be influenced by body weight status or the presence of metabolic disorders. In this study, most of the respondents with low fat intake were overweight and obese [36].

A person who consumes more salt than is excreted through the kidneys increases blood vessel volume and cardiac output. Due to an increase in vascular volume and cardiac output, the body's response is an increase in blood pressure. Salt not only affects vascular mechanisms but also activates neural, endocrine, or paracrine responses [37]. Consuming excessive amounts of salt such as >10 grams is found in people who often consume instant foods, anchovies, cuttlefish, shrimp, and noodles with soy sauce [38]. According to research Shafrina et al. The average amount of salt consumption in Southeast Asia varies, such as in Indonesia, from 1.94 g/day to >5 g/day [37]. High sodium intake is associated with increased blood pressure levels due to changes in vascular resistance, sodium consumption can trigger water retention and cause arteries to experience high flow, changes in plasma sodium levels not only impact small resistance arteries but can also affect function and structure large elastic arteries. Other side effects of consuming excessive amounts of sodium are microvascular endothelial inflammation, anatomical remodeling, and functional abnormalities [39].

#### **3.2.5. Relationship between physical activity and the incidence of hypertension**

In this study, it was found that 23.8% of subjects did not exercise either regularly or not at all, and also 76.2% of subjects exercised either regularly or irregularly. According to the relationship, there is no significant difference in the possibility of hypertension between subjects who do or do not do physical activity with the result of Asymptotic significance (2-

sided) namely 0.454 ( $>0.05$ ). Subjects who are active have a 0.76 times risk of developing hypertension. Changes in blood pressure during exercise are the difference between peak and initial systolic blood pressure of at least 60 mmHg in men and at least 50 mmHg in women during exercise testing, or it is also said that systolic blood pressure in men exceeds 210 mmHg and in women exceeds 190 mmHg. This research according to Caselli et al is in line with this research because subjects who do a physical activity can increase systolic blood pressure [40]. The normal value of blood pressure during maximum exercise testing is 220/85 mmHg in male athletes and 200/80 mmHg in female athletes. The increase in blood pressure increases for a short period, especially during heavy lifting activities, blood pressure values usually increase to 480/350 mmHg [41].

Purba et al.'s research states that there is a relationship between the incidence of hypertension and physical activity. The OR value was 3.619, which means that people who do light exercise have a three times greater risk of developing hypertension compared to people who do moderate and heavy activities [42].

Light physical activity or a sedentary lifestyle is an activity that spends time sitting or standing with little body movement, not exercising, and using vehicles. Heavy physical activity is someone who is active and works hard for a long period. Physical activity affects blood pressure, heart rate is usually higher in people who are not active. High blood pressure is caused by hardening of the heart muscle which functions to pump, thereby placing a burden on the artery walls and peripheral resistance. Not doing physical activity can also cause obesity which affects blood pressure [42].

### **3.2.6. Relationship between smoking activities and the incidence of hypertension**

In this study, it was found that 68.3% of subjects were not active smokers and 31.7% of subjects were active smokers. According to the relationship, there is a significant difference between patients who are active and inactive smokers with the incidence of hypertension with the result of Asymptotic significance (2-sided) namely 0.039 ( $<0.05$ ). According to this study, subjects who were exposed to cigarette smoke were 5.2 times more likely to develop hypertension. Misnaniarti et al's research agrees with this research where the risk of hypertension is two times greater in smokers. There is also a relationship between the two with logistic regression showing that smoking has an increased risk of developing hypertension by 20% (OR = 1.97; 95% CI = 1.01-1.41;  $p = 0.034$ ) [43]

Octavian et al's research also states that there is a significant relationship between smoking and blood pressure ( $p=0.003$ ). This is proven by an increase in blood pressure measurements from  $140 \pm 7/99 \pm 3$  mmHg to  $151 \pm 5/108 \pm 2$  mmHg after 10 minutes of inhaling cigarette smoke [44]

The nicotine in cigarettes affects a person's blood pressure. This may be caused by atherosclerosis which appears due to plaque, other effects occur in the release of the hormones epinephrine and norepinephrine or the effect of CO on red blood cell proliferation [44]. Long-term tobacco use affects the adaptability of blood vessels through endothelial damage and chronic inflammation [45]

## **4. CONCLUSION**

1. The incidence of hypertension in the Kramat Jati District Health Center, East Jakarta in 2024 will be 81 people.
2. some factors cannot be changed in outpatients at the Kramat Jati District Health Center, East Jakarta, namely age 56-65 years, female, and have a family history of hypertension.
3. some factors can be changed in outpatients at the Kramat Jati District Health Center, East Jakarta, namely consuming  $\geq 5$  grams of sodium per day, doing physical activity, and not smoking.

4. There is a significant relationship between factors that cannot be changed, namely age and family history of hypertension, with the incidence of hypertension at the Kramat Jati District Health Center, East Jakarta.
5. There is a significant relationship between factors that can be changed, namely dietary sodium patterns and smoking activities, with the incidence of hypertension at the Kramat Jati District Health Center, East Jakarta.

## REFERENCES

- [1] Lukito AA, Harmeiwaty E, Umami V. Association of Indonesian Hypertension Doctors Editor. 2018.
- [2] Oparil S, Acelajado MC, Bakris GL, Berlowitz DR, Cífková R, Dominiczak AF, et al. Hypertension. Vol. 4, Nature Reviews Disease Primers. Nature Publishing Group; 2018. <https://doi.org/10.1038/nrdp.2018.14> PMID: 29565029
- [3] Oliveros E, Patel H, Kyung S, Fugar S, Goldberg A, Madan N, et al. Hypertension in older adults: Assessment, management, and challenges. Vol. 43, Clinical Cardiology. John Wiley and Sons Inc.; 2020. p. 99–107. <https://doi.org/10.1002/clc.23303> PMID: 31825114
- [4] Medicine F. HYPERTENSION Frits Reinier Wantian Suling.
- [5] Mills KT, Stefanescu A, He J. The global epidemiology of hypertension. Vol. 16, Nature Reviews Nephrology. Nature Research; 2020. p. 223–37. <https://doi.org/10.1038/s41581-019-0244-2> PMID: 32024986
- [6] Chan GC, Teo BW, Tay JC, Chen CH, Cheng HM, Wang TD, et al. Hypertension in a multi-ethnic Asian population of Singapore. Vol. 23, Journal of Clinical Hypertension. John Wiley and Sons Inc; 2021. p. 522–8. <https://doi.org/10.1111/jch.14140> PMID: 33340436
- [7] Liew SJ, Lee JT, Tan CS, Koh CHG, Van Dam R, Müller-Riemenschneider F. Sociodemographic factors with hypertension prevalence, awareness, treatment and control in a multi-ethnic Asian population: A cross-sectional study. *BMJ Open*. 2019 May 1;9(5). <https://doi.org/10.1136/bmjopen-2018-025869> PMID: 31110091
- [8] Zhou B, Carrillo-Larco RM, Danaei G, Riley LM, Paciorek CJ, Stevens GA, et al. Worldwide trends in hypertension prevalence and progress in treatment and control from 1990 to 2019: a pooled analysis of 1201 population-representative studies with 104 million participants. *The Lancet*. 2021 Sep 11;398(10304):957–80. [https://doi.org/10.1016/S0140-6736\(21\)01330-1](https://doi.org/10.1016/S0140-6736(21)01330-1) PMID: 34450083
- [9] Study P, Society K, Medicine F. LITERATURE REVIEW OF HYPERTENSION AND ITS MANAGEMENT By dr. I Made Sutarga, M. Kes. 2017.
- [10] Indonesian Ministry of Health. Results of Basic Health Research (Riskesdas) 2018. Health Research and Development Agency, Ministry of the Republic of Indonesia. 2018;
- [11] Ondimu DO, Kikvi GM, Otieno WN. Risk factors for hypertension among young adults (18-35) years attending in tenwek mission hospital, Bomet county, Kenya in 2018. *Pan African Medical Journal*. 2019;33. <https://doi.org/10.11604/pamj.2019.33.210.18407> PMID: 31692887
- [12] Poulter NR, Prabhakaran D, Caulfield M. Hypertension. In: *The Lancet*. Lancet Publishing Group; 2015. p. 801–12. [https://doi.org/10.1016/S0140-6736\(14\)61468-9](https://doi.org/10.1016/S0140-6736(14)61468-9) PMID: 25832858
- [13] Roerecke M, Kaczorowski J, Tobe SW, Gmel G, Hasan OSM, Rehm J. The effect of a reduction in alcohol consumption on blood pressure: a systematic review and meta-analysis. *Lancet Public Health*. 2017 Feb 1;2(2):e108–20. [https://doi.org/10.1016/S2468-2667\(17\)30003-8](https://doi.org/10.1016/S2468-2667(17)30003-8) PMID: 29253389
- [14] James PA, Oparil S, Carter BL, Cushman WC, Dennison-Himmelfarb C, Handler J, et al. 2014 Evidence-based guideline for the management of high blood pressure in adults:

- Report from the panel members appointed to the Eighth Joint National Committee (JNC 8). Vol. 311, JAMA. American Medical Association; 2014. p. 507–20. <https://doi.org/10.1001/jama.2013.284427> PMID: 24352797
- [15] Mancía G, Fagard R, Narkiewicz K, Redón J, Zanchetti A, Böhm M, et al. 2013 ESH/ESC Guidelines for the management of arterial hypertension. *J Hypertens*. 2013 Jul;31(7):1281–357. <https://doi.org/10.1097/01.hjh.0000431740.32696.cc>
- [16] James PA, Oparil S, Carter BL, Cushman WC, Dennison-Himmelfarb C, Handler J, et al. 2014 Evidence-Based Guideline for the Management of High Blood Pressure in Adults. *JAMA*. 2014 Feb 5;311(5):507. <https://doi.org/10.1001/jama.2013.284427>
- [17] Iskedjian M, Einarson TR, MacKeigan LD, Shear N, Addis A, Mittmann N, et al. Relationship between daily dose frequency and adherence to antihypertensive pharmacotherapy: Evidence from a meta-analysis. *Clin Ther*. 2002 Feb;24(2):302–16. [https://doi.org/10.1016/S0149-2918\(02\)85026-3](https://doi.org/10.1016/S0149-2918(02)85026-3)
- [18] Rachmawati S, Prihastuti-Puspitasari H, Zairina E. The implementation of a chronic disease management program (Prolanis) in Indonesia: A literature review. In: *Journal of Basic and Clinical Physiology and Pharmacology*. De Gruyter; 2020. <https://doi.org/10.1515/jbcpp-2019-0350> PMID: 31860469
- [19] Yusransyah, Halimah E, Suwantina AA. Measurement of the quality of life of prolanis hypertension patients in sixteen primary healthcare centers in Pandeglang District, Banten Province, Indonesia, using the EQ-5D-5L instrument. *Patient Prefer Adherence*. 2020;14:1103–9. <https://doi.org/10.2147/PPA.S249085>
- [20] Cohen L, Curhan GC, Forman JP. Influence of age on the association between lifestyle factors and risk of hypertension. *Journal of the American Society of Hypertension*. 2012 Jul;6(4):284–90. <https://doi.org/10.1016/j.jash.2012.06.002> PMID: 22789880
- [21] Pustaka T, Age H, Rahmawati M, Kasih RP. Galenical is licensed under a Creative Commons Attribution-ShareAlike 4.0 International License. Vol. 2. 2023.
- [22] Research Innovation J, Lailatus Sa O, Martadani L, Taqiyuddin A, Economics University A Wahab Hasbullah Email FK. ANALYSIS OF EMPLOYEE PERFORMANCE DIFFERENCES AT PT SURYA INDAH FOOD MULTIRAS JOMBANG. 2021;2(2):515.
- [23] Kyriazis M. Aging as “Time-Related Dysfunction”: A Perspective. Vol. 7, *Frontiers in Medicine*. Frontiers Media S.A.; 2020. <https://doi.org/10.3389/fmed.2020.00371>
- [24] Mouhtadi BB, Kanaan RMN, Iskandarani M, Rahal MK, Halat DH. Prevalence, awareness, treatment, control and risk factors associated with hypertension in Lebanese adults: A cross-sectional study. *Glob Cardiol Sci Pract*. 2018;2018(1). <https://doi.org/10.21542/gcsp.2018.6>
- [25] Kusumawaty J, Hidayat N, Ginanjar E. Relationship between gender and hypertension intensity in the elderly in the Laknok Health Center working area, Ciamis Regency. *Mutiara Medika*. 2016;16(2):46-51.
- [26] Farhan Dito M, Alfarisi R, Kurniati M. THE RELATIONSHIP BETWEEN HERDERSTANDING HISTORY AND CHILD'S GENDER IN FAMILIES IN THE WORKING AREA OF THE BANDAR LAMPUNG RELIGIOUS AFFAIRS OFFICE [Internet]. Vol. 10, *Journal of Medical and Health Sciences*. 2023.
- [27] Li A le, Peng Q, Shao Y qin, Fang X, Zhang Y ying. The interaction on hypertension between family history and diabetes and other risk factors. *Sci Rep*. 2021 Dec 1;11(1). <https://doi.org/10.1038/s41598-021-83589-z> PMID: 33633182
- [28] Liu M, He Y, Jiang B, Wang J, Wu L, Wang Y, et al. Association between family history and hypertension among Chinese elderly. *Medicine (United States)*. 2015;94(48). <https://doi.org/10.1097/MD.0000000000002226> PMID: 26632912
- [29] Rodríguez-Moran M, Aradillas-García C, Simental-Mendía LE, Monreal-Escalante E, De La Cruz Mendoza E, Dávila Esqueda ME, et al. Family history of hypertension and cardiovascular risk factors in prepubertal children. *Am J Hypertens*. 2010 Mar;23(3):299–304. <https://doi.org/10.1038/ajh.2009.257> PMID: 20075847

- [30] Gluckman PD, Hanson MA. Developmental origins of disease paradigm: A mechanistic and evolutionary perspective. Vol. 56, *Pediatric Research*. Lippincott Williams and Wilkins; 2004. p. 311–7. <https://doi.org/10.1203/01.PDR.0000135998.08025.FB> PMID: 15240866
- [31] Manosroi W, Williams GH. Genetics of Human Primary Hypertension: Focus on Hormonal Mechanisms. *Endocr Rev*. 2019 Jun 1;40(3):825–56. <https://doi.org/10.1210/er.2018-00071>
- [32] Antonicelli R, Testa R, Bonfigli AR, Sirolla C, Pieri C, Marra M, et al. Relationship between lipoprotein(a) levels, oxidative stress, and blood pressure levels in patients with essential hypertension. *Clin Exp Med*. 2001 Sep 1;1(3):145–50. <https://doi.org/10.1007/s10238-001-8026-z>
- [33] Fauzi Mrwah S, Evelianti MS, Jemmy Wowor TF. FACTORS RELATED TO THE INCIDENCE OF ADULT HYPERTENSION DURING THE COVID-19 PANDEMIC IN PABUARAN CIBINONG BOGOR DISTRICT. Vol. 10, *Journal of Nursing*. 2022.
- [34] Pertiwi GAR, Aryawangsa AAN, Prabawa IPY, Manuaba IBAP, Bhargah A, Ratni NWS, et al. Factors associated with visit-to-visit variability of blood pressure in hypertensive patients at a Primary Health Care Service, Tabanan, Bali, Indonesia. *Fam Med Community Health*. 2018;6(4):191–9. <https://doi.org/10.15212/FMCH.2018.0124>
- [35] Farapti F, Nadhiroh SR, Sayogo S, Mardiana N. Urinary and dietary sodium to potassium ratio as a useful marker for estimating blood pressure among older women in Indonesian urban coastal areas. *Med J Nutrition Metab*. 2017 Aug 31;10(2):113–22. <https://doi.org/10.3233/MNM-17138>
- [36] Wahyu Wadarsih N, Ayu Dewi Sartika R. Factors Associated with Hypertension among Adults in West Java, Indonesia. *Indian J Public Health Res Dev*. 2020 Jan 1;11(1):1331. <https://doi.org/10.37506/v11/i1/2020/ijphrd/194028>
- [37] Shafrina AA, Sulastris D, Burhan IR. The Relationship between Salt Consumption Levels and the Incident of Hypertension in Southeast Asia. *Indonesian Journal of Health Sciences*. 2022 Aug 26;2(3):202–11. <https://doi.org/10.25077/jikesi.v2i3.452>
- [38] Jia Jiet L, Soma R. M. High salt diets in young university adults and the correlation with blood pressure, protein intake and fat free mass. *Bioscience Horizons: The International Journal of Student Research*. 2017 Jan 1;10. <https://doi.org/10.1093/biohorizons/hzx003>
- [39] Grillo, Salvi, Coruzzi, Salvi, Parati. Sodium Intake and Hypertension. *Nutrients*. 2019 Aug 21;11(9):1970. <https://doi.org/10.3390/nu11091970>
- [40] Caselli S, Serdoz A, Mango F, Lemme E, Vaquer Seguí A, Milan A, et al. High blood pressure response to exercise predicts the future development of hypertension in young athletes. *Eur Heart J*. 2019 Jan 1;40(1):62–8. <https://doi.org/10.1093/eurheartj/ehy810>
- [41] Schweiger V, Niederseer D, Schmied C, Attenhofer-Jost C, Caselli S. Athletes and Hypertension. *Curr Cardiol Rep*. 2021 Dec 16;23(12):176. <https://doi.org/10.1007/s11886-021-01608-x>
- [42] Purba EN, Santosa H, Siregar FA. The Relationship of Physical Activity and Obesity with the Incidence of Hypertension in Adults Aged 26-45 Years in Medan. *Open Access Maced J Med Sci*. 2019 Oct 14;7(20):3464–8. <https://doi.org/10.3889/oamjms.2019.447>
- [43] Misnaniarti, Nugraheni WP, Nantabah ZK, Restuningtyas FR, Hartono RK, Rachmawati T, et al. Smoking behavior and hypertension among health workers during the COVID-19 pandemic: a case study in Java and Bali-Indonesia. *Front Cardiovasc Med*. 2023 Aug 14;10. <https://doi.org/10.3389/fcvm.2023.114685>
- [44] Health J, Octavian Y, Setyanda G, Sulastris D, Lestari Y. Relationship between smoking and the incidence of hypertension in men aged 35-65 years in Padang City [Internet]. Vol. 4, *Andalas*. 2015.
- [45] Gao N, Liu T, Wang Y, Chen M, Yu L, Fu C, et al. Assessing the association between smoking and hypertension: Smoking status, type of tobacco products, and interaction with alcohol consumption. *Front Cardiovasc Med*. 2023 Feb 9;10. <https://doi.org/10.3389/fcvm.2023.1027988>