

Original Research Article

Effect of Organic Manure and Plant Growth Promoting Rhizobacteria Enriched Biocapsules on Growth Yield and Quality of Cape Gooseberry (*Physalis Peruviana* L.)

Abstract

The present experiment was conducted at Department of Horticulture, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj. Prayagraj during the session 2023 - 2024. The experiment was laid out in randomized block design with three replications, and the study consists of Ten treatment combinations including control by using different Effect of organic manure and plant growth promoting rhizobacteria enriched biocapsules on growth yield and quality of cape gooseberry. The best treatment was T₈ (Poultry manure @0.56 kg + 500 PPM Biocapsule) & T₇ (Green manure @1.12 kg + 500 PPM Biocapsule) which shows highest values in all the parameters viz., plant height (85.80 cm), number of leaf/plant (71.76), Leaf area (cm²) (64.62 cm²), Days to anthesis of 1st flower after planting (35.75), Flowering duration (138.36), flower/plant (118.86), Fruit setting after flowering (8.91), fruit set to maturity (50.10), 1st fruit harvesting (101.16), fruit/plant (84.00) fruit weight (10.77 g), fruit yield/plant (400.68 kg), TSS (17.68 %), Ascorbic acid (61.22 mg/100g), Acidity (0.49 %), total sugar (12.39 %). All the treatments were significantly superior in their growth, flowering, fruit yield and quality of cape gooseberry over control (T₀) and (T₉).

Key words: Cape gooseberry, organic, biocapsule, growth, yield, quality

Comment [WU1]: Replace suitable key words.

INTRODUCTION

Cape-gooseberry (*Physalis peruviana* L.) is a plant in the nightshade family (Solanaceae), native to warm temperate and subtropical regions throughout the world. *Physalis peruviana* was derived from the Greek word *physa* meaning bladder, for the calyx covering the fruit and *peruviana* meaning of Peru. It is commonly known as Goldenberry, Husk Cherry, Peruvian Ground Cherry, Poha Berry, Rasbhari, Makoi etc. The cape

gooseberry is an annual in temperate regions and a perennial in the tropics. In northern India, it is not cultivated above 1200 m, but in Southern India it thrives up to 1800 m above the mean sea level. It is an herbaceous, semi-shrub that is upright, perennial in subtropical zones and can grow until reaches 0.9 m. The fruit is 4–5 g in weight, remains protected by a calyx and covered by a brilliant yellow peel (**Mayorga et al., 2001**). The cape gooseberry is known as goldenberry in European countries, uchuva in Colombia, uvilla in Ecuador, aguaymanto in Peru and toptopo in Venezuela (**Puente et al., 2011**). Three types of cape gooseberry indigenous to Colombia, Kenya and South Africa are cultivated worldwide. Colombia is the world's largest producer of cape gooseberry followed by South Africa.

The organic manures such as FYM, vermicompost and press mud influence the physical chemical as well as biological properties of the soil which improve soil fertility, structure, porosity, aeration, drainage and water relation capacity. FYM supplies all major nutrients (N, P, K, Ca, Mg, S,) necessary for plant growth, as well as micronutrients (Fe, Mn, Cu and Zn). Hence, it acts as a mixed fertilizer. FYM improves soil physical, chemical and biological properties. Improvement in the soil structure due to FYM application leads to a better environment for root development. FYM also improves soil water holding capacity (**Tadesse et al., 2013**).

Poultry manure is an excellent organic fertilizer, as it contains high nitrogen, phosphorus, potassium and other essential nutrients. In contrast to chemical fertilizer, it adds organic matter to soil which improves soil structures, nutrient retention, aeration, soil moisture holding capacity and water infiltration (**Deksissa et al., 2008**).

Biocapsule, a bio-fertilizer technology developed by the IISR (Indian Institute of Spices Research). The reason biocapsules are significant is that they allow for the encapsulation of targeted microorganisms in a gelatin capsule, which is then transferred to the crops to improve soil nutrient solubilization, growth, and yield. Because a single gramme of capsules has the same amount of microorganisms as a kilogramme box of powdered biofertilizer or a litre bottle, they are incredibly effective. Additionally, unlike many liquid-based bio formulations, these microbial strains are not at risk of viability loss at room temperature since they are maintained in the dormant condition. It is made out of a carrier medium that is abundant in live microorganisms. Applied to seed, soil, or growing plants, it either boosts or organically releases the nutrients from the earth.

MATERIALS AND METHODS

This experiment was laid out during the August 2023 to March 2024 at Horticulture

Research Farm, Department of Horticulture, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj (U.P.). The horticulture research farm is situated at 25° 39' 42" N latitude, 81° 67' 56" E longitude and at an altitude of 98 m above mean sea level. The treatment consisted of T₀ - Control, T₁ - FYM @ 1.68 kg, T₂ - Vermicompost @ 0.56 kg, T₃ - Green Manure @ 1.12 kg, T₄ - Poultry Manure @ 0.56 kg, T₅ - FYM @ 1.68 Kg + 500 ppm Biocapsule, T₆ - Vermicompost @ 0.56 kg + 500 ppm Biocapsule, T₇ - Green Manure @ 1.12 kg + 500 ppm Biocapsule, T₈ - Poultry Manure @ 0.56 kg + 500 ppm Biocapsule. The experiment was laid out in a Randomized Block Design with 9 treatments and replicated thrice. Data recorded on different aspects of fruit crop, viz., growth, yield were subjected to statistical analysis by analysis of variance method. (Gomez and Gomez, 1976) and economic data analysis mathematical method.

Comment [WU2]: Cite latest reference

Add field photographs

Comment [WU3]: Add photos

RESULT AND DISCUSSION

Vegetative growth of Cape Gooseberry-

The data pertaining to Vegetative growth of cape gooseberry in different treatment combinations was recorded and are presented in table 1.

Plant height (cm)

At 120 DAT, significantly and maximum plant height (85.80 cm) was recorded in T₈ [Poultry Manure + 500 ppm Biocapsule] followed by the T₇ [Green Manure + 500 ppm Biocapsule] (84.97 cm), whereas minimum plant height of (75.33 cm) was recorded in T₀ [Control].

Biocapsules increases the nutrient fixation in the root and nanozeolite played an important role to make availability of nutrient in the soil. These results are in support with Bandana and Chandel (2017) and Padhan *et al.* (2019). It might be due to the more availability of nutrients by poultry manure throughout the growing season. Nitrogen increases photosynthetic activity and helps in maintaining higher auxin level which might have resulted in better plant height Verma *et al.*, (2017b).

Comment [WU4]: Check spelling

Number of Leaf per plant

At 120 DAT, significantly and maximum number of leaf per plant (71.76) was recorded in T₈ [Poultry Manure + 500 ppm Biocapsule] followed by the T₇ [Green Manure + 500 ppm Biocapsule] (69.20), whereas minimum number of leaf per plant of (62.47) was recorded in T₀ [Control].

The significant and maximum number of leaf/plant was with application of poultry manure might be due to with increased levels of NPK function in most of the physiological and metabolic processes resulting in increased growth and development, resulting in maximum leaf/plant. Similar result was also reported by **Verma *et al.*, (2017b)**. Further the significant and maximum leaf/plant was with application of biocapsule might be due to involvement of stimulation of cell division, photosynthetic process as well as formation of chlorophyll **Singh *et al.*, (2023)**.

Leaf area (cm²)

At 120 DAT, significantly and maximum leaf area (64.62 cm²) was recorded in T₈ [Poultry Manure + 500 ppm Biocapsule] followed by the T₇ [Green Manure + 500 ppm Biocapsule] (63.01 cm²), whereas minimum leaf area of (55.50 cm²) was recorded in T₀ [Control].

The significant and maximum leaf area cm² was with application of poultry manure might be due to with increased levels of NPK function in most of the physiological and metabolic processes resulting in increased growth and development, resulting in maximum leaf area cm². Similar result was also reported by **Verma *et al.*, (2017b)**. Further the significant and maximum leaf/plant was with application of biocapsule might be due to involvement of stimulation of cell division, photosynthetic process as well as formation of chlorophyll **Singh *et al.*, (2023)**.

Reproductive Growth of Cape Gooseberry

The data pertaining to reproductive growth of cape gooseberry after planting in different treatment combinations was recorded and are presented in table 2

Days to anthesis of 1st flower after planting

Significantly the minimum days to anthesis of 1st flower after planting of 35.75 days after planting was recorded at T₈ [Poultry Manure + 500 ppm Biocapsule] followed by

Comment [WU5]: Remove

T₇[Green manure + 500 ppm Biocapsule] of 37.60 days, whereas minimum days to anthesis of 1st flower after planting of 44.57 days was recorded in T₀ [Control].

The minimum in number of days to anthesis of cape gooseberry due to treatment might be due to fact that poultry manure regulate the growth by causing cell division and cell elongation in plant system. These results are in conformity with **Deepa *et al.*, (2014)**.

Flowering duration

Significantly the maximum flowering duration of 138.36 days after planting was recorded at T₈ [Poultry Manure + 500 ppm Biocapsule] followed by T₇[Green manure + 500 ppm Biocapsule] of 134.60 days, whereas minimum days to anthesis of 1st flower after planting of 116.33 days was recorded in T₀ [Control].

The results of present findings indicated that the sole application of organic manure did not have any significant effect on the flowering and fruiting but their combination of biocapsule showed better results. The earliness in flowering coupled with higher number of flowers after application of FYM, poultry manure and vermicompost might be due to enhanced level of nitrogen and phosphorus. They were proved to be better for synthesis of growth promoters and increased photosynthate production. These findings are in accordance with **Arancon *et al.* (2004)** in strawberry and **Yadav *et al.* (2011)** in papaya.

Total number of flowers plant⁻¹

Significantly the maximum total number of flowers plant⁻¹ 118.86 was recorded at T₈ [Poultry Manure + 500 ppm Biocapsule] followed by T₇[Green manure + 500 ppm Biocapsule] of 116.98, whereas minimum total number of flowers plant⁻¹ of 101.85 was recorded in T₀ [Control].

The increase in number of flowers might be due to organic fertilizers that produced significant increase in vegetative growth which in turn produced more photosynthates which were probably diverted towards the more flower production. The findings are in accordance with the results obtained by **Mohamed and El-Schrawy (2013)** in mango and **El-Miniawy *et al.* (2014)** in strawberry.

YIELD AND YIELD ATTRIBUTES

The data pertaining to yield and yield attributes in different treatment combinations was recorded and are presented in table 3

Total fruit number plant⁻¹

Significantly the maximum number of fruits per plant 84.00 was recorded at T₈ [Poultry Manure + 500 ppm Biocapsule] followed by T₇[Green manure + 500 ppm Biocapsule] of 82.33, whereas minimum number of fruits per plant 71.33 was recorded in T₀ [Control].

It may possibly be due to fact that INM application accelerated the development of fruits which are positively correlated with the number of fruits in the following spring. Increased number of fruits might have also resulted because of increase in number of flowers per plant. Similar observations were also reported by **Tripathi et al. (2010)** in strawberry.

Average fruit weight (g)

Significantly the maximum average fruit weight 10.77 was recorded at T₈ [Poultry Manure + 500 ppm Biocapsule] followed by T₇[Green manure + 500 ppm Biocapsule] of 9.10, whereas minimum average fruit weight 6.98 was recorded in T₀ [Control].

The higher fruit weight of kiwifruit in the combined application organic manures and inorganic fertilizers may be attributed to the relative increase in availability of nutrients and better solute uptake by plants (**Korwar et al., 2006**). These findings are in close conformity with **Ray et al. (2008)** in papaya and **Gautam et al. (2012)** in Mango.

Fruit yield plant⁻¹(g)

Significantly the maximum fruit yield per plant 918.97 g was recorded at T₈ [Poultry Manure + 500 ppm Biocapsule] followed by T₇[Green manure + 500 ppm Biocapsule] of 699.86 g, whereas minimum fruit yield per plant 497.84 g was recorded in T₀ [Control].

Further, the higher yield obtained with combined application of inorganic fertilizer and organic manure might be attributed to conducive physical and biological environment of soil which lead to higher and balanced nutrients uptake (**Deshpande and Senapathy,2010**).

QUALITY PARAMETERS:

The data pertaining to quality parameters in different treatment combinations was recorded and are presented in table 4

Total soluble solids (%)

Significantly the total soluble solids 17.68 was recorded at T₈ [Poultry Manure + 500 ppm Biocapsule] followed by T₇[Green manure + 500 ppm Biocapsule] of 16.98, whereas minimum total soluble solids 13.26 was recorded in T₀ [Control].

Comment [WU6]: No need to represent the combination repeatedly.

Generally, addition of organic manures for nutrients, moisture and growth promoting substances enhance the metabolic and hormonal activity of the plant. This promotes production of more photosynthates to be stored in fruits in the form of starch and carbohydrates. It is an established fact that the transformation of mature fruit into ripe form i.e. during the process of ripening in storage, the fruit undergoes physical, physiological and bio-chemical changes. The increase in total soluble solids (TSS) may be attributed to the conversion of reserved starch and other insoluble carbohydrates into soluble sugars. These results elucidate the findings of **Prasanna and Rajan (2001)** and **Yadav *et al.* (2011)**.

Comment [WU7]: Capitalize each word

Ascorbic acid (mg/100g)

Significantly the ascorbic acid 61.22 was recorded at T₈ [Poultry Manure + 500 ppm Biocapsule] followed by T₇[Green manure + 500 ppm Biocapsule] of 60.96, whereas minimum ascorbic acid 50.24 was recorded in T₀ [Control].

The ascorbic acid content was found higher with the application of organic supplements than the inorganic fertilizers. The negative relationship between nitrogen levels and vitamin-C content can be seen. This might be due to the exposure of plant with more of nitrogen, which increased protein production but reduce the carbohydrates. Similar findings have been reported by **Worthington, (2001)**; **Bahadur *et al.* (2006)** and **Citak and Sonmez, (2010)**. Ascorbic acid is vulnerable to loss as it is easily oxidizes in the presence of light, oxygen and heat.

Acidity (%)

Significantly the acidity 0.49 was recorded at T₈ [Poultry Manure + 500 ppm Biocapsule] followed by T₇[Green manure + 500 ppm Biocapsule] of 0.52, whereas minimum acidity 0.75 was recorded in T₀ [Control].

The decrease in acidity might be due to the degradation of organic acids. A gradual declining trend in the acidity content of fruit in all the treatments was observed with the advancement of storage period. Similar results were observed by **Rawat (2001)**. These results are also in agreement with the findings of **Sivakumar et al. (2010)** in Litchi.

Total sugar (%)

Significantly the total sugar 12.39 was recorded at T₈ [Poultry Manure + 500 ppm Biocapsule] followed by T₇ [Green manure + 500 ppm Biocapsule] of 0.52, whereas minimum total sugar 8.98 was recorded in T₀ [Control].

Generally, addition of organic manures for nutrients, moisture and growth promoting substances enhance the metabolic and hormonal activity of the plant. This promotes production of more photosynthates to be stored in fruits in the form of starch and carbohydrates. It is an established fact that the transformation of mature fruit into ripe form i.e. during the process of ripening in storage, the fruit undergoes physical, physiological and bio-chemical changes. The increase in total sugars (TS) may be attributed to the conversion of reserved starch and other insoluble carbohydrates into soluble sugars. These results elucidate the findings of **Prasanna and Rajan (2001)** and **Yadav et al. (2011)**.

ECONOMIC PARAMETER OF CAPE GOOSEBERRY

The calculations regarding the gross return, net return and benefit cost ratio of the cape gooseberry was recorded, tabulated and analyzed. Benefit cost ratio were found significantly higher in different treatments and is clear from the table 5

Cost of cultivation (Rs. /ha)

The maximum cost of cultivation (Rs. 105000) was observed in T₅ (Fym @ 1.68 + 500 ppm Biocapsule) followed by (Rs. 103000) was observed in T₆ (Vermicompost @ 0.56 Kg + 500 ppm Biocapsule) while the minimum cost of cultivation (Rs. 90000) was observed in T₀ (Control).

Gross Return (Rs. /ha)

The maximum gross return (Rs. 354156) was observed in T₈ (Poultry Manure + 500 ppm Biocapsule) followed by (Rs. 280476) was observed in T₆ (Vermicompost @ 0.56 Kg + 500 ppm Biocapsule) while the minimum gross return (Rs. 191860) was observed in T₀ (Control).

Net Return (Rs. /ha)

The maximum net return (Rs. 243156) was observed in T₈ (Poultry Manure + 500 ppm Biocapsule) followed by (Rs. 177476) was observed in T₆ (Vermicompost @ 0.56 Kg + 500 ppm Biocapsule) while the minimum cost of net return (Rs.101860) was observed in T₀ (Control).

Benefit Cost ratio (B:C)

The maximum benefit cost ratio (2.51) was observed in T₈ (Poultry Manure + 500 ppm Biocapsule) followed by (1.72) was observed in T₆ (Vermicompost @0.56 Kg + 500 ppm Biocapsule) while the minimum cost of cultivation (1.08) was observed in T₂ (Vermicompost @ 0.56 kg).

CONCLUSION

Based on the results of the present study:

It is concluded that, overall treatment T₈ (Poultry Manure + 500 ppm Biocapsule) performed best in terms vegetative as well as reproductive growth of cape gooseberry, yield, yield attributes and Economics of cape gooseberry was also obtained from this treatment.

Table 1 Effect of Organic Manure and Plant Growth Promoting Rhizobacteria Enriched Biocapsule vegetative growth of cape Gooseberry

Treatment Symbol	Treatment combinations	Plant height (cm)	Number of leaf/plant	Leaf area
		120 DAT	120 DAT	(cm ²)120 DAT
T ₀	CONTROL	75.33	62.47	55.50
T ₁	FYM @ 1.68 kg	78.47	64.09	57.50
T ₂	VERMICOMPOST @ 0.56 kg	76.56	63.33	56.88
T ₃	GREEN MANURE @ 1.12 kg	81.60	67.89	60.15
T ₄	POULTRY MANURE @ 0.56 kg	79.09	65.47	59.61
T ₅	FYM @ 1.68 kg + 500 PPM BIOCAPSULE	82.94	68.10	61.99
T ₆	VERMICOMPOST @0.56 kg + 500 PPM BIOCAPSULE	83.90	68.79	62.89
T ₇	GREEN MANURE @ 1.12 kg + 500 PPM BIOCAPSULE	84.97	69.20	63.01
T ₈	POULTRY MANURE @0.56 kg +500 PPM BIOCAPSULE	85.80	71.76	64.62
	F-test	S	S	S
	SEm(±)	2.33	1.92	1.83
	CD (p=0.05)	6.98	6.77	5.47

Table 2 Effect of Organic Manure & Plant Growth Promoting Rhizobacteria Enriched Biocapsule reproductive growth of cape

Treatment Symbol	Treatment combinations	Days to Anthesis of 1 st Flower after Planting	Flowering duration	Total number of flowers plant ⁻¹
T ₀	CONTROL	44.57	116.33	101.85
T ₁	FYM @ 1.68 kg	43.04	125.33	106.34
T ₂	VERMICOMPOST @ 0.56 kg	42.72	124.88	103.67
T ₃	GREEN MANURE @ 1.12 kg	41.46	126.11	110.89
T ₄	POULTRY MANURE @ 0.56 kg	38.58	125.98	107.98
T ₅	FYM @ 1.68 kg + 500 PPM BIOCAPSULE	40.35	133.73	111.29
T ₆	VERMICOMPOST @ 0.56 kg + 500 PPM BIOCAPSULE	37.60	132.96	114.79
T ₇	GREEN MANURE @ 1.12 kg + 500 PPM BIOCAPSULE	36.50	134.60	116.98
T ₈	POULTRY MANURE @ 0.56 kg + 500 PPM BIOCAPSULE	35.75	138.36	118.86
	F-test	S	S	S
	SEm(±)	1.57	4.06	3.30
	CD (p=0.05)	4.66	12.18	9.88

Gooseberry


Table 3 Effect of Organic Manure & Plant Growth Promoting Rhizobacteria Enriched Biocapsule yield of cape Gooseberry

Treatment Symbol	Treatment combinations	Fruit		
		Number of fruit plant ⁻¹	Average fruit weight (g)	yield plant ⁻¹ (g)
T ₀	CONTROL	71.33	6.98	497.84
T ₁	FYM @ 1.68 kg	75.00	7.97	597.70
T ₂	VERMICOMPOST @ 0.56 kg	73.33	7.35	538.93
T ₃	GREEN MANURE @ 1.12 kg	78.00	8.90	694.32
T ₄	POULTRY MANURE @ 0.56 kg	75.67	8.65	654.24
T ₅	FYM @ 1.68 kg + 500 PPM BIOCAPSULE	78.67	8.50	711.91
T ₆	VERMICOMPOST @ 0.56 kg + 500 PPM BIOCAPSULE	80.00	9.05	727.78
T ₇	GREEN MANURE @ 1.12 kg + 500 PPM BIOCAPSULE	82.33	9.10	699.86
T ₈	POULTRY MANURE @ 0.56 kg + 500 PPM BIOCAPSULE	84.00	10.77	918.97
	F-test	S	S	S
	SEm(±)	2.33	0.39	14.32
	CD (p=0.05)	6.98	1.18	42.93

Table 4: Effect of Organic Manure and Plant Growth Promoting Rhizobacteria Enriched Bio capsules quality of Cape Gooseberry

Treatment	Treatment Combination	Total soluble	Ascorbic acid	Acidity	Total sugar
-----------	-----------------------	---------------	---------------	---------	-------------

UNDER PEER REVIEW



Symbol		solids (%)	(mg/100g)	(%)	(%)
T ₀	CONTROL	13.26	50.24	0.75	8.98
T ₁	FYM @ 1.68 kg	13.98	57.79	0.68	9.91
T ₂	VERMICOMPOST @ 0.56 kg	13.50	57.12	0.72	9.66
T ₃	GREEN MANURE @ 1.12 kg	14.88	58.99	0.61	10.44
T ₄	POULTRY MANURE @ 0.56 kg	14.30	58.33	0.65	10.10
T ₅	FYM @ 1.68+ 500 PPM BIOCAPSULE	15.67	59.66	0.59	10.77
T ₆	VERMICOMPOST @0.56 kg + 500 PPM BIOCAPSULE	16.55	60.44	0.55	11.50
T ₇	GREEN MANURE @ 1.12 kg + 500 PPM BIOCAPSULE	16.98	60.96	0.52	11.98
T ₈	POULTRY MANURE @ 0.56 kg +500 PPM BIOCAPSULE	17.68	61.22	0.49	12.39
	F-test	S	S	S	S
	SEm(±)	0.67	1.77	0.02	0.02
	CD (p=0.05)	2.00	5.29	0.06	0.06

Table 5: Effect of Organic Manure and Plant Growth Promoting Rhizobacteria Enriched Bio capsules economics of Cape Gooseberry

Treatment Symbol	Treatment Combination	Cost of cultivation/ha	Gross Return/ha	Net Return/ha	B:C Ratio
T₀	CONTROL	90000	191860	101860	1.13
T₁	FYM @ 1.68 kg	101000	230346	129346	1.28
T₂	VERMICOMPOST @ 0.56 kg	100000	207696	107696	1.08
T₃	GREEN MANURE @ 1.12 kg	99000	267582	168582	1.70
T₄	POULTRY MANURE @ 0.56 kg	96000	252134	156134	1.63
T₅	FYM @ 1.68+ 500 PPM BIOCAPSULE	105000	274361	169361	1.61
T₆	VERMICOMPOST @0.56 kg + 500 PPM BIOCAPSULE	103000	280476	177476	1.72
T₇	GREEN MANURE @ 1.12 kg + 500 PPM BIOCAPSULE	102000	269715	167715	1.64
T₈	POULTRY MANURE @ 0.56 kg +500 PPM BIOCAPSULE	101000	354156	253156	2.51

REFERENCES

Comment [WU8]: Uniform formatting and style must be followed

- Arancon NQ, Edwards CA, Bierman P, Welch C and Metzger JD. 2004. Influences of vermicompost on field strawberries: 1. Effects on growth and yields. *Bioresource Technology* **93**:145–153.
- Bahadur A, Singh J, Singh KP, Upadhyay AK and Rai M. 2006. Effect of organic amendments and biofertilizers on growth, yield and quality attributes of Chinese cabbage (*Brassica pekinensis*). *Indian journal of agricultural science* **76**:596-598.
- Bandana, Chandel JS. Effect of growth regulators, PGPR and nitrogen on growth of apple nursery plants under protected conditions. *Journal of Hill Agriculture*. 2017;**8**(2):176-180.
- Citak S and Sonmez S. 2010. Effects of conventional and organic fertilization on spinach (*Spinacea oleracea* L.) growth, yield, vitamin C and nitrate concentration during two successive seasons. *Scientia horticulturae* **126**:415-420.
- Deepa, H. D., Anil, K. Y., & Pawan, K. (2014) Integrated Nutrient Management in Cape gooseberry (*Physalis peruviana* L.) for Peri Urban Horticulture. *Indian journal of applied research*, **12**(4):274-275.
- Deksissa, T., Short, I. and Allen, J. 2008. Effect of soil amendment with compost on growth and water use efficiency of Amaranth. In: Proceedings of the UCOWR/NIWR annual conference: International water resources: challenges for the 21st century and water resources education, July 22 – 24, 2008, Durham, NC.
- Deshpande HH and Senapathy P. 2010. Effect of green manuring and organic manures on yield, quality and economics of rice (*Oryza sativa* L.) under lowland condition. *Karnataka Journal of Agricultural Science* **23**:235-238.
- El-miniawy, S. M., Ragab, M. E., Youssef, S. M. and Metwally, A. A., 2014, Influence of foliar spraying of seaweed extract on growth, yield and quality of strawberry plants. *J. Appl. Sci. Res.* **10**(2): 88-94
- Gautam US, Singh R, Tiwari N, Gurjar PS and Kumar A. 2012. Effect of integrated nutrient management in mango cv. Sunderja. *Indian Journal of Horticulture* **69**:151-155.
- Gomez, K. A. and Gomez, A. A. (1976). Statistical procedures for agriculture Research, 2nd Edition, John Wiley and Son, New York, 680p.
- Korwar GR, Pratibha G, Ravi V and Palanikumar D. 2006. Influence of organics and inorganics on growth, yield of aonla (*Emblica officinalis*) and soil quality in semi-arid tropics. *Indian Journal of Agricultural Science* **76**:457-461.

- Mayorga, H., Knapp, H., Winterhalter, P., and Duque, C. Glycosidically bound flavour compounds of cape gooseberry (*Physalis peruviana* L.). *Journal of Agriculture and Food Chemistry*, **49**: 1904–1908, 2001.
- Mohamed, A. Y. and El-sehrawy, O. A., 2013, Effect of seaweed extract on fruiting of Hindy Bisinnara mango trees. *J. Sci.*,**9**(6): 537-544.
- Padhan A, Mishra S, Bahadur V. Effect of growing media on growth, development and establishment of low chilling variety of apple HRMN-99 under Prayagrajagro climatic conditions. *Journal of Pharmacognosy and Phytochemistry*. 2019;**8**(3):1227-1230.
- Prasanna KP and Rajan S. 2001. Effect of organic farming on storage life of brinjal fruits. *South Indian Horticulture* **49**:255-256.
- Puente, L. A., Pinto, S. A., Castro, E. S. and Cortés, M. *Physalis peruviana* L., The multiple properties of a highly functional fruit: a review. *Food Research International*, **44**: 1733-1740, 2011
- Rawat VR, Tomar YK, Rawat JM. 2010. Influence of foliar application of micronutrients on the fruit quality of guava cv. Lucknow-49. *Journal of Hill Agriculture*.
- Ray PK, Singh AK and Kumar Arun. 2008. Performance of Pusa Delicious papaya under organic farming. *Indian Journal of Horticulture***65**:21-24.
- Shivakumar BS, Dharmatti PR and Channal HT. 2012. Effect of organic cultivation of papaya on yield, economics and soil nutrient status. *Karnataka Journal of Agricultural Sciences***25**:488-492.
- Singh, R. K., Mishra, S., & Bahadur, V. (2023) Effect of Nano-chitosan, Nano-micronutrients and Bio capsules on Vegetative growth, flowering and fruiting attributes of Strawberry (*Fragaria* × *ananassa*) cv. Winter dawn. *Biological Forum – An International Journal***15**(5),289-295.
- Tadesse, T., Dechassa, N., Bayu, W. and Gebeyehu, S. 2013. Effects of farm yard manure and inorganic fertilizer application on soil physic-chemical properties and nutrient balance in rain-fed lowland rice ecosystem. *Am. J. Plant Sci.* **4**:309-316.
- Tripathi, V. K., Kumar, N., Shukla, H. S. and Mishra, A. N. Influence of Azotobacter, Azospirillum and PSB on growth, yield and quality of strawberry cv. Chandler, Paper presented in National Symposium on Conservation Hort., during March, 21-23, 2010 at Dehradun, pp 198-199, 2010.

- Verma, N., Dwivedi, D. H., Kishor, S., & Singh, N. (2017b). Impact of integrated nutrient management on growth and fruit physical attributes in Cape gooseberry, *Physalis peruviana*. *Bioscience biotechnology research communications*, **10**(4), 672-675.
- Worthington V. 2001. Nutritional quality of organic versus conventional fruits, vegetables, and grains. *The Journal of Alternative and Complementary Medicine***7**:161-173.
- Yadav PK, Yadav AL, Yadav AS, Yadav HC and Singh YP. 2011. Effect of integrated nutrient nourishment on yield attributes and economics of papaya (*Carica papaya* L.) fruit cv. Pusa dwarf. *Plant Archives***11**:307-309.

UNDER PEER REVIEW