

# FUNCTIONAL AND PHYSICOCHEMICAL PROPERTIES OF FLOUR AND COOKIES FROM WHEAT SOYBEAN AND ORANGE FLESHED POTATOES BLEND

## ABSTRACT

The incorporation of nutritionally-rich underutilized local crops into popularly consumed snacks such as cookies could be employed to address the above challenge. Hence, the feasibility of producing cookies from wheat flour, soybean and orange fleshed sweet potatoes was investigated. A completely randomized design (CRD) was used in the design of experiment. A total of 5 samples of the composite flour and one control (100 % wheat flour) in varying proportions of 100:0:0, 75:20:5, 70:20:10, 65:20:15, 60:20:20 were obtained and used for the production of the cookies. The cookies were subjected to functional, proximate, antinutrient and physical analyses using standard laboratory procedures. The functional properties of the flour increased in addition of soybean and OFSP. The proximate results of the cookies showed that protein, ash, fat, crude fibre, moisture and Carbohydrate ranged from (10.54 - 11.39), (1.61 - 2.95), (1.60 - 3.30), (1.25 - 3.38), (10.77 - 14.36) and (65.99 - 74.22) respectively. The antinutrients analysed showed a range of (4.81 - 11.87), (11.71 - 19.03), (6.87 - 10.73 TIU/g) and (7.35 - 12.67 mg/100g) for oxalate, tannin, trypsin inhibitor and phytate respectively. The minerals (iron and zinc) were bioavailable in the cookies and the effect of phytate could not hinder their bioavailability. The physical properties (weight, diameter and thickness) of the cookies substituted with constant 20% soybean in combination with variations of OFSP increased significantly ( $p < 0.05$ ) except the spread ratio. The study suggests that incorporation of soybean and OFSP flour into wheat flour will improve the nutritional quality of cookies mostly consumed by the populace.

*Keywords: functional, proximate, mineral, antinutritional, molar ratio, soybean, orange fleshed sweet potatoes, wheat*

## 1. INTRODUCTION

“Snacks including cookies are widely consumed all over the world” [1]. “They represent an important part of human diet especially as a source of energy and may be consumed in between meals. The consumption of cereal snacks foods, such as biscuits, cookies, wafers and short bread has become very popular in Nigeria especially among children of which cookies are one of the most consumed cereal foods apart from bread, because they are readily

available in local shops as ready to eat, cheap, convenient and appetizing food products" [2]. [3] observed that "carbohydrates and added sugars tend to be over-consumed at snacking occasions. They opined that replacement of current snack choices with nutrient-dense foods could lower the risks of nutrient deficiencies and help lower excess nutrient consumption of which can be obtained from a composite flour". "Composite flour is a mixture of varying proportion of two or more flour which may or may not contain wheat flour and used for production of bread, pastries, cake and other confectionery products that are conventionally produced from wheat flour with the intention of increasing the essential nutrients in human diet and increase the economic relevance of indigenous crops" [4].

According to [5], "composite flour are potential gateways to stimulate and scale up the consumption of underutilized crops such as orange sweet fleshed potatoes or neglected crops. Cookies with high sensory ratings have been produced from blends of malted barley/wheat, wheat/jackfruit seed and other products. Wheat (*Triticum aestivum* L.) has been used since time immemorial for the manufacture of several bakery products because of the presence of unique gluten protein which is responsible for providing the viscoelastic characteristics to dough". [6] discovered that "in wheat products such as bread, gluten network formation is desirable for gas retention and better volume of product, while in products such as biscuits and cookies, extensibility is required, so gluten formation is undesirable". "Due to the limited amount of wheat produced in Nigeria and cost of importation, government has collaborated with research institutes to encourage the use of composite flour in the production of bread and related food products such as cookies" [7]. "Soybean (*Glycine max*) is an edible legume highly famed for being nutritious" [8]. "As they contain a high amount of protein, essential vitamins, minerals, complex carbohydrates, and dietary fibre" [9]. "Soybeans also contain phytochemicals that contribute to the health of the human body" [10]. "Orange Fleshed Sweet Potatoes (OFSP) serve as a cheap and sustainable source of Pro-Vitamin A, especially for vulnerable populations. OFSP is a special type of biofortified sweet potato that contains 76kcal energy, 1.37g protein, 2.5g fibre, 0.72mg iron, 0.2mg zinc, 588 mg RAE of vitamin A and 12.8g of vitamin C". [11]. "It appears that there is a growing demand and interest for OFSP because it is a good source of beta-carotene, the major precursor of vitamin A. Depending on variety, the beta-carotene content in OFSP may vary between 5091 µg/100 g and 16,456 µg/100 g" [12]. "OFSP has been suggested to have great potentials to be used in food-based programs to address vitamin A deficiency" [13]. [7] observed that "in Nigeria, the reliance on wheat flour by baking industries has over the years restricted the use of other cereals, tuber and oilseed crops available for domestic use. Cookies are grossly nutritionally inadequate as they consist mainly of un-supplemented cereals deficient in Pro-Vitamin A and other essential amino acids such as lysine, threonine and tryptophan which are necessary for proper growth and development". OFSP and soybean which are valuable sources of vitamin A carotenoids, antioxidants, minerals, proteins, fats, dietary fibre etc., could be exploited in the production of cookies. The nutritional content of cookies is expected to improve as a result of addition of Pro-vitamin A carotenoids from OFSP flour and minerals, proteins, fat from soybean flour. Formulating complementary food with wheat, soybean and OFSP flour could not only improve the nutrient composition of the complementary food but also reduces post-harvest losses of this locally grown food items. The thrust of this study therefore is to improve the nutritional content of cookies through fortification of composite flours obtained from underutilized crops such as OFSP and soybeans.

## **2. MATERIAL AND METHODS**

### **2.1 Materials and Sample Collection**

Dangote wheat flour, Dangote granulated sugar, Peak powdered milk, topper baking fat, Dangote salt and Longman baking powder were procured from Wuse market, Abuja FCT, Nigeria. Soybean seeds and orange fleshed sweet potatoes tubers were obtained from Gosa market, airport road, Abuja FCT, Nigeria. The chemicals used were of analytical grade.

## 2.2 Preparation of Soybean Flour

The method described by [14] was employed. The soybeans were sorted and washed with clean water. The seeds were then soaked overnight in cold water for 12 h at 30 °C and de-hulled. It was then boiled at 100 °C for 30 min. The seeds were drained through a nylon sieve for 1 h. The seeds were oven dried at 60 °C for 30 min, milled and sieved using 0.5 mm mesh. The flour was stored in low density dark coloured polythene bags stored in plastic containers with airtight lids at room temperature (30±2°C) until analysis.

## 2.3 Preparation of OFSP Flour

The orange flesh sweet potato flour was produced using the method described by [15]. The sweet potato tubers were peeled and cut into thin pieces manually. The potato slices were then soaked for 30 min to prevent browning reactions. Drying of sweet potato slices was done in an oven at 60 °C till constant weight. The dried sweet potato chips were milled into flour using the laboratory grinder and passed through 0.5 mm mesh sieve, packed in airtight containers and stored in the refrigerator till further use.

## 2.4 Research Design

The research design is Completely Randomized design (CRD). Three different flours (wheat, soybean and OFSP) were mixed at five different proportions to give 100g composite.

## 2.5 Flour Blend Formulation

The composite flour was prepared by replacing wheat flour (WF) with soybean and OFSP flours at the ratios of 100:0:0, 75:20:5, 70:20:10, 65:20:15 and 60:20:20 respectively as shown in Table 1. Sample ratio of 100:0:0 with 100% WF served as a reference sample. The winnowed materials were oven dried at 60 °C for 30 min and milled with the aid of a Kenwood Chef blender to form fine particle size prior to its use. The wheat, soybean and orange fleshed sweet potatoes flours were weighed separately with the aid of a digital electronic weighing balance at different proportions. The formulation was done so as to enhance the nutrient (particularly protein and Pro-vitamin A) content of the cookies.

**Table 1: Formulation of wheat, soybean and orange fleshed sweet potatoes flour blends for cookies production**

W:S:OFSP	Wheat Flour	Soybean	OFSP Flour
100:0:0	100	20	00
75:20:5	75	20	5
70:20:10	70	20	10
65:20:15	65	20	15
60:20:20	60	20	20

## 2.6 Preparation of Cookies from Composite Flours

The cookies were prepared using the method described by [16]. The essential ingredients for the production of cookies in their various proportions were weighed in. Sweeteners (sucralose and shortening) were creamed for 10 mins until light and fluffy. Egg and flour were added to the mixture followed by milk, salt and baking powder using a Kenwood hand mixer. The mixture was thoroughly kneaded into consistent dough for about 30 min and allowed to rest. The

dough was rolled on flat wooden surface sprinkled with flour to form a uniform thickness and cut into predetermined size and shape using a biscuit cutter. The dough was arranged in greased trays and baked in a preheated oven operating at 178 °C for 25 min. The biscuits were allowed to cool down to the room temperature for 5 – 10 min before packaged in polythene bags and stored for evaluation.

## **2.7 Determination of Functional Properties**

The parameters of functional properties determined were OAC, BD, GT, SI and WAC as described by [17].

## **2.8 Proximate Analysis**

The crude protein content, crude fibre content, moisture content, fat content and ash content of the composite flour were determined in triplicate using established analytical procedures of [18]. The carbohydrate content was estimated by difference from 100% after accounting for moisture, protein, fibre, ash, and fat.

## **2.9 Antinutritional Composition**

The Tannin, oxalate, trypsin inhibitor and phytate was determined as described by [19].

## **2.10 Determination of Phytate Mineral Molar Ratio**

The moles of phytate and minerals were obtained by dividing the weight of phytate with minerals of each mineral molecular weight as described by [19]. The phytate mineral molar ratio was obtained by dividing the moles of phytate with the moles of each mineral.

## **2.11 Determination of Physical Properties**

The weight, diameter, thickness and spread ration of cookies were determined according to the method described by [20].

## **2.12 Statistical Analysis**

The experiments were conducted in triplicates and data generated were analysed using analysis of Variance (ANOVA). Means were separated by Duncan multiple range test. Significant difference was accepted at 5 % level of probability ( $p < 0.05$ ) using Statistical package for social sciences (SPSS) version 23.

# **3. RESULTS AND DISCUSSION**

## **3.1 Functional Properties of Wheat, Soybean and OFSP Composite Flour**

“Functional properties of flour are important for the selection of crops for use in value-added product development. The swelling index is the degree of starch from the flour that absorbs water. It is worthy to note that swelling capacity is evidence of non-covalent bonding between molecules within starch granules and also a factor of the ratio of  $\alpha$ -amylose and amylopectin ratios. However, the swelling capacity of flours depends on size of particles, types of variety and types of processing methods or unit operations” [21]. The overall values were lower (2.213 - 3.617) than wheat-soy-sweet potato composite flour reported by [22] who observed that there was decrease in swelling capacity as the level of incorporation of sweet potatoes increased. This observation was also reported by [21] submitting that the swelling index of composite flours are highly affected by the level of potatoes flour, because potatoes is a rich source of starch. Swelling power of flour granules is an indication of the extent of associative forces within the granule. It depends on sizes of particles, types of variety and processing methods [23]. Lower swelling index observed in samples with 60:20:20 % of W:S:OFSP could be attributed to the increased substitution and fat content of soybean which interferes with starch

granules by forming films around the granules and high protein which forms a starch protein complex thus reducing the swelling capacity [24]. The 5% OFSP showed a higher swelling index due to high content of wheat which have high amylopectin content as observed by [25].

Water absorption capacity (WAC) reflects the amount of water that the flour can absorb and retain. The overall values were similar to the values obtained by [26] for chickpea flour (1.92 g/g) and teff and amaranth flours (0.95 g/g). However, it was observed that the WAC increased as the proportion of OFSP increased from 5 – 20%. These values were similar to the values (1.57 – 2.01) reported by [23] for varieties of OFSP breeding lines. This shows that OFSP had the ability to enhance the absorption of water during of flour processing into dough. The high values of WAC can be attributed to the amount and nature of hydrophilic constituent and nature of the protein. Agreeing to the result of this study, [23] suggests that “the flours from the OFSP would be useful in foods such as bakery products which require hydration to improve handling characteristics”. “For an improved food texture of baked products (such as bread, cookies), higher values for water absorption are desired. Higher water absorption values were attributed to the higher content of starch and fibre” [27]. [27] also stated that “higher protein content tends to increased water absorption therefore this shows that 60:20:20 exhibited the highest water absorption capacity”. [23] reported that “water absorption property indicates the ability of a product to associate with water under conditions when water is limiting such as making of dough and paste”. “The water and oil absorption capacity depend on the type of protein, amino acid composition and protein polarity and hydrophobicity” [28].

The oil absorption capacity (OAC) of the control (100% wheat) had a lowest value. The increase in values obtained for fortified samples were higher than values (1.8 – 1.9 g/g) obtained by [26] for gluten free flours such as amaranth, tiger nut, and chickpea flours and values (1.10 - 1.40 g/ml) obtained by [23] for varieties of OFSP. In their opinion, the oil absorption capacity of flours is important for the development of new food products as well as their storage stability (particularly for flavour binding and in the development of oxidative rancidity). “Both the protein content and type contribute to the oil-retaining properties of food materials. Oil absorption capacity is attributed mainly to the physical entrapment of oils. It is an indication of the rate at which protein binds to fat in food formulations” [29]. “They noted that high OAC flours are suitable for retaining the flavour and enhance the mouthfeel when used in foods. Moreover, variation in the amylose/amylopectin ratio and content of individual flours can contribute to differences in the water as well as oil absorption capacity of flour” [26].

The bulk densities increased ( $P = .05$ ) significantly in the fortified samples while a reduction in value was observed as the incorporation of OFSP increased from 5% to 20% respectively. The bulk density observed in this study is within the range 0.68 – 0.82 g/cm<sup>3</sup> as reported by [30]. [31] reported “increase in bulk density which agrees with this work”. “The increase in bulk density could be due to the increased addition of composite flour” [21]. “Nutritionally, low bulk density promotes digestibility of foods especially in children with immature digestive systems while high bulk density decreases the caloric and nutrient intake of children resulting in growth faltering” [31]. “Low bulk density then implies that the product can easily be packaged for economic use. Moreover, low bulk density has nutritional and economic significance as more of the products can be eaten thus leading to high energy and nutritional density. However, bulk density is a measurement of the porosity of a product which influences the design of the packaging material. Bulk density is affected by the moisture content and the particle sizes of the starch. The differences in the particle size may be the cause of variations in bulk density of the OFSP. The high bulk density of the sweet potato breeding lines is an indication that they can be used as thickeners” [23].

“The variation in the thermal properties among the different flours are influenced by some factors such as: size of the starch granule, molecular structure of the amylopectin (branch, length and weight), starch, protein and dietary fibre content, as well as the presence of other compounds” [32]. “The gelatinization temperature of the control (100:0:0) was lower than

values obtained from the fortified samples. However, there was a significant ( $p < 0.05$ ) temperature increase in the fortified samples. During starch gelatinization, the helix structure and crystallinity of the starch is lost and the granule is disrupted. Albeit, the increase in temperature may be due to higher fat and fibre content of soybean and sweet potatoes. The swelling of the starch granule is disturbed by the presence of non-starch compounds such as fat, which leads to higher gelatinization temperatures" [33]. [34] reported that "gelatinization temperature of the flour samples blends generally increased with increasing addition of OFSP and cashew nut flour. Increasing fibre content appears to delay gelation and subsequently its temperature. Thus, higher heat energy is required to attain significant gelation. They discovered that waxy and regular maize gelatinize at 62 - 72°C, whereas high-amylose starches begin to swell below 100°C, temperatures greater than 130°C are required to fully disperse these starches. This is because more amylose molecules are involved in the crystalline regions of the high amylose starch than in waxy and regular starches". However, [35] submitted that "decrease in gelation capacity is due to the addition of composite flours thus increasing the concentration of protein and enhancing the interaction among the binding forces which in turn increases the gelling ability of the flour". "Gelation properties are related to water absorption capacities and takes place at high water protein concentration because of the great intermolecular contact during heating" [35]. The formation of gel provides a structural matrix for holding water and other soluble materials like sugar and flavour [36].

**Table 2: Functional properties of wheat, soybean and OFSP composite flour**

W:S:OFSP (%)	SI	WAC (g/ml)	OAC (g/g)	BD (g/cm <sup>3</sup> )	GT (°C)
100:0:0	1.20 <sup>d</sup> ±0.02	1.31 <sup>e</sup> ±0.04	1.61 <sup>d</sup> ±0.03	0.730 <sup>e</sup> ±0.002	68.0 <sup>c</sup> ±0.0
75:20:5	1.38 <sup>a</sup> ±0.03	1.87 <sup>d</sup> ±0.05	2.32 <sup>c</sup> ±0.08	0.825 <sup>a</sup> ±0.003	73.0 <sup>b</sup> ±1.4
70:20:10	1.35 <sup>ab</sup> ±0.00	1.99 <sup>c</sup> ±0.02	2.51 <sup>b</sup> ±0.02	0.793 <sup>b</sup> ±0.003	74.0 <sup>ab</sup> ±1.4
65:20:15	1.31 <sup>bc</sup> ±0.01	2.10 <sup>b</sup> ±0.00	2.59 <sup>b</sup> ±0.03	0.783 <sup>c</sup> ±0.001	75.5 <sup>ab</sup> ±0.7
60:20:20	1.27 <sup>c</sup> ±0.01	2.31 <sup>a</sup> ±0.01	2.75 <sup>a</sup> ±0.01	0.764 <sup>d</sup> ±0.004	76.5 <sup>a</sup> ±0.7

*Values are means ± standard deviations of triplicate determinations. Data in the same column bearing different superscripts differ significantly  $P=0.05$*

### 3.2 Proximate Composition of Wheat Soybean Orange Fleshed Sweet Potatoes Composite Flour

The MC in samples was observed to increase in the fortified samples as OFSP increased with a constant 20% rate of soybean. There was no significant ( $P = .05$ ) difference between samples with the control (100% wheat) and fortified flours. The values obtained were similar to 10.20 - 12.22% obtained for wheat, soybean moringa leaf composite flours [37] and also in agreement with the research work of [38]. The moisture content values obtained were within the recommended moisture content (10%) for storage stability [39] thus, the sweet potato flour will have a longer shelf life if properly stored under a good condition. "However, the moisture content increased with decrease in the carbohydrate value of the samples enhancing nutritional products, thus implying that the flour can be stored if properly packaged" [40].

The ash contents were observed to increase as shown in the results. There was a significant ( $P = .05$ ) difference between the control (100% wheat) and the fortified samples, however the ash content increased as the rate of incorporation increased, a similar trend also observed by [41]. [42] reported lower ash contents values than those of this work. The increase in the ash content was as a result of the OFSP flour. According to [43] the increase in ash content implies that the samples are good sources of minerals. [44] confirmed "the high crude ash contents

(4.60 - 7.20%) of OFSP (KJP specie) could be a rich source of mineral salts". "Ash is an inorganic compound present in a food which helps in the breaking down of some organic compounds such as protein, fat and carbohydrates" [39].

The fat content increased ( $P = .05$ ) significantly in the fortified samples but decreased as the rate of incorporation increased. [45] reported a decrease in fat content when meat balls were produced with increased addition of African yam bean (AYB). The increase in fat was reported in complementary flours due to higher fortification with chickpea [46], a similar legume like soybean. The increased fat content could be due to the composition of AYB and due to its high fat content [47], a similar observation in soybean. The low-fat content obtained in this study agrees with the findings of [48] that sweet potato is low in fat.

There was a significant ( $P = .05$ ) increase in the fibre content in the fortified samples. A similar observation was reported by [37] with values (2.05% - 3.00%) for wheat soybean moringa leaf composite flour. They submitted that the crude fibre content of the flour was enhanced with increase in moringa leaf flour and 20% soybean flour constant. This could be so as soybean, moringa leaf and OFSP are excellent sources of fibre. However, [40] reported "a decrease in fibre with increased addition of soybean and carrot, this study observed an increase due to the addition of OFSP showing the important benefit that fibre can help increase the utilization and absorption of some other micronutrients" [40]. The fibre content was higher in samples with 20% OFSP than other samples.

The protein content of the fortified samples increased significantly ( $P = .05$ ) but decreased as the level of incorporation of OFSP increased. [49] reported an increase in protein when pap and *agidi* jollof were produced by the increased addition of AYB flour in line with this work. The protein values (10.24% - 28.81%) obtained by [37] was similar to the values in this work. They reported that the protein content of the flour increased with a decrease in wheat, an increase in moringa leaf and 20% soybean flour constant. According to [50], increase in protein was observed due to the addition of cowpea which also agrees with this work. This increase in the protein content is due to the increased addition of soybean which is an excellent source of dietary protein. The increase in protein can result in amino acid complementation from soybean to those in the cereal protein [49;40]. Protein is required especially during weaning in order to prevent protein energy malnutrition [51].

The carbohydrate content significantly ( $P = .05$ ) decreased was in line with [40] who opined that carbohydrate content decreased due to increased substitution of OFSP which agrees with this work. The carbohydrate content was similar to values (67.67 - 44.94%) obtained for wheat soybean moringa leaf composite flour [37]. According to [14] carbohydrate provides heat and energy for all forms of body activity.

**Table 3: Proximate composition of wheat soybean and OFSP composite flour**

W:S: OFSP (%)	Moisture	Ash	Fat	Crude fibre	Protein	Carbohydrate
100:0:0	10.54 <sup>d</sup> ±0.08	1.61 <sup>e</sup> ±0.07	1.60 <sup>d</sup> ±0.00	1.25 <sup>d</sup> ±0.07	10.77 <sup>e</sup> ±0.10	74.22 <sup>a</sup> ±0.19
75:20:5	10.72 <sup>c</sup> ±0.02	2.42 <sup>d</sup> ±0.01	3.30 <sup>a</sup> ±0.14	2.67 <sup>c</sup> ±0.04	14.36 <sup>a</sup> ±0.04	65.99 <sup>e</sup> ±0.16
70:20:10	10.85 <sup>c</sup> ±0.07	2.60 <sup>c</sup> ±0.02	3.10 <sup>ab</sup> ±0.14	2.90 <sup>b</sup> ±0.02	14.11 <sup>b</sup> ±0.04	66.33 <sup>d</sup> ±0.09
65:20:15	11.06 <sup>b</sup> ±0.03	2.70 <sup>b</sup> ±0.00	2.90 <sup>b</sup> ±0.00	3.23 <sup>a</sup> ±0.09	13.42 <sup>c</sup> ±0.09	66.78 <sup>c</sup> ±0.01
60:20:20	11.39 <sup>a</sup> ±0.03	2.95 <sup>a</sup> ±0.01	2.65 <sup>c</sup> ±0.07	3.38 <sup>a</sup> ±0.04	12.69 <sup>d</sup> ±0.05	67.46 <sup>b</sup> ±0.10

*Values are means ± standard deviations of triplicate determinations. Data in the same column bearing different superscripts differ significantly  $P = .05$*

### 3.3 Antinutritional Contents of Wheat, Soybean and Orange Fleshed Sweet Potatoes Composite Flour

There were significant ( $P = .05$ ) difference in the varieties of composite flours examined in Table 4. 20% OFSP had the highest oxalate value. The variations observed might be attributed to the increased rate of incorporation of OFSP. The oxalate values obtained in this study were lower compared to the values reported by [52] for unpeeled OFSP. Oxalate has been reported by [44] “to have a harmful effect on human nutrition and health because it has the ability to reduce calcium absorption in the form of calcium oxalate in the blood and thus aid the formation of a kidney stone”.

The highest tannin was recorded for 20% OFSP and the least value was observed in control (100:0:0). The results obtained were lower than values reported by [53]. [54] further reported similar values (0.09 - 0.26 mg/100g) of wheat-AYB-tigernut biscuits. However, the tannin content reduced significantly which could be because of the soaking processing method used. A similar finding was reported by [55] “for cookies made from tigernut flour. Based on their findings, [54] submitted that tannin inhibits the activities of some enzymes such as trypsin, amylase, and lipase to form insoluble complexes with protein and divalent ions like  $Fe^{2+}$  and Zn thereby reducing their absorption in the body”.

The phytate content of the composite flour decreased with the level of incorporation of OFSP where the 75:20:5 had a highest value and the 60:20:20 had a lower value. High values of anti-nutrient (phytate, oxalate and tannin) in food are undesirable because they form complexes with minerals and proteins resulting in its unavailable to the body system [53] thereby leading to carcinogenesis, shock and renal damage. “Anti-nutritional factors are substances that destructively affect the nutritional composition of a food thereby reducing both the mineral and protein bioavailability, digestion and its utilizations in the body” [53].

“The trypsin inhibitor showed significant ( $P = .05$ ) increase in the fortified flour (6.87 – 10.73 TIU/g) where the 20% OFSP had the lower value of 8.56 TIU/g and 5% had a highest value of 10.73 TIU/g. It was observed that all the molar ratios were below the critical limits (Phy: Ca > 1.56, Phy: Fe > 14 and Phy: Zn > 10)” as reported by [56]. “This implies that the bioavailability of Ca, Fe & Zn are not inhibited by the concentration of phytate in the samples examined. Hence, all the minerals (Ca, Fe and Zn) in the fortified samples will be adequately absorbed by the body when consumed. The molar ratio between phytate and divalent cations (Ca, Fe and Zn) indicates the impact of phytate on the bioavailability (ability of the body to absorb and digest minerals in a food after consumption) of dietary minerals and the absorption of these cations were not adversely affected by the amount of phytate in the varieties of sweet potato examined” [53].

**Table 4: Antinutrients concentration of wheat, soybean and OFSP composite flour**

W:S:OFSP	Oxalate (mg/100g)	Tannin (mg/100g)	Trypsin inhibitor (TIU/g)	Phytate (mg/100g)
100:0:0	4.81 <sup>e</sup> ±0.04	11.71 <sup>e</sup> ±0.04	6.87 <sup>d</sup> ±0.11	7.35 <sup>e</sup> ±0.00
75:20:5	9.96 <sup>d</sup> ±0.10	11.92 <sup>d</sup> ±0.00	10.73 <sup>a</sup> ±0.12	12.67 <sup>a</sup> ±0.12
70:20:10	10.73 <sup>c</sup> ±0.03	17.31 <sup>c</sup> ±0.04	10.49 <sup>a</sup> ±0.12	12.08 <sup>b</sup> ±0.06
65:20:15	11.21 <sup>b</sup> ±0.05	17.91 <sup>b</sup> ±0.02	9.36 <sup>b</sup> ±0.05	11.76 <sup>c</sup> ±0.05
60:20:20	11.87 <sup>a</sup> ±0.04	19.03 <sup>a</sup> ±0.17	8.56 <sup>c</sup> ±0.04	10.71 <sup>d</sup> ±0.09

*Values are means ± standard deviations of triplicate determinations. Data in the same column bearing different superscripts differ significantly  $P=.05$*

### 3.4 Antinutritional Content of Wheat Soybean Orange Fleshed Sweet Potatoes Cookies

The oxalate content of the fortified cookies had the highest values in sample with 20 % OFSP and the least in sample with 5 % OFSP. Samples containing OFSP had a significant ( $P = .05$ ) increase as the percentage OFSP increased. The tannin content was observed to decrease for all samples where the control sample was observed to have a least tannin value of 5.89 mg/100g. This trend was similar to the values obtained by [54] for biscuit produced from composite flour of wheat, AYB and tigernut. They opined that the reduction could be because of the soaking processing method used. Samples containing OFSP had the highest trypsin inhibitor value in sample with 5 % OFSP and the least in sample with 20 % OFSP.

The phytate content of the samples increased for all samples where the control sample recorded a least phytate value. The values obtained in this study were lower compared to the phytate values reported by [52] “for peeled and unpeeled OFSP (77.75 and 95.15 mg/100g, respectively)”. “Phytate produces phytic acid which is a major phosphorous storage component that chelate metallic ion such as Zinc, Calcium and Iron, thereby reducing their bioavailability” [57]. Findings from [58] buttressed that “cookies in present study were safe to consumers due the processing methods and preparing cookies in the oven decreased the phytate content. Their findings stated that phytic acid decreased during bakery product making due to the action of phytase as well as cooking temperature”.

Bioavailability of iron and zinc values show that iron and zinc of the cookies were bioavailable to consumers without the binding effect of phytate. This is due to the molar ratio of phytate: iron values were below the critical value (14) and phytate: zinc values were also below the critical value (10). The measured values of phytate: iron decreased progressively when OFSP amounts were increased; so, bioavailability of iron is more when OFSP was more in the blended cookies. The measured values of phytate: zinc decreased when the amount of OFSP increased in the developed cookies. This may increase the zinc bioavailability in more OFSP added cookies due to high values of phytate: zinc ratios. The iron and zinc contents in the present study cookies can be taken without hindering effect of the phytate. Generally, the phytate contents of the composite flour cookies and the molar ratios of phytate: minerals imply that, phytate in the cookies of the present study could not impair the bioavailability of iron and zinc to consumers.

**Table 5: Antinutrients concentration of the wheat, soybean and orange fleshed sweet potatoes cookies**

W:S:OFSP (%)	Oxalate (mg/100g)	Tannin (mg/100g)	Trypsin inhibitor (TIU/g)	Phytate (mg/100g)
100:0:0	0.52 <sup>d</sup> ±0.04	5.89 <sup>e</sup> ±0.03	0.99 <sup>d</sup> ±0.04	0.92 <sup>e</sup> ±0.00
75:20:5	1.22 <sup>c</sup> ±0.04	8.95 <sup>a</sup> ±0.00	2.28 <sup>a</sup> ±0.07	1.87 <sup>a</sup> ±0.02
70:20:10	1.21 <sup>c</sup> ±0.01	8.50 <sup>b</sup> ±0.04	2.04 <sup>b</sup> ±0.04	1.68 <sup>b</sup> ±0.01
65:20:15	1.39 <sup>b</sup> ±0.02	7.65 <sup>c</sup> ±0.02	1.74 <sup>c</sup> ±0.02	1.50 <sup>c</sup> ±0.09
60:20:20	1.64 <sup>a</sup> ±0.02	7.08 <sup>d</sup> ±0.06	1.66 <sup>c</sup> ±0.02	1.22 <sup>d</sup> ±0.02

*Values are means ± standard deviations of triplicate determinations. Data in the same column bearing different superscripts differ significantly  $P = .05$*

**Table 6: Molar ratio of Phytate/Mineral (bioavailability) of wheat, soybean and OFSP cookies**

Samples (W:S:OFSP)						
Molar Ratio						Critical Limits
	100:0:0	75:20:5	70:20:10	65:20:15	60:20:20	
Phy-Ca	0.007	0.0013	0.0011	0.0010	0.0008	>1.56
Phy-Mg	0.0005	0.0010	0.0009	0.0008	0.0006	
Phy-Ph	0.0006	0.001	0.0009	0.0009	0.0007	
Phy-Na	0.0017	0.0025	0.0023	0.0022	0.0019	
Phy-K	0.0003	0.0004	0.0004	0.0004	0.0003	
Phy-Fe	0.032	0.05	0.04	0.03	0.03	>14
Phy=Zn	0.04	0.066	0.055	0.047	0.03	>10

*Values are means  $\pm$  standard deviations of triplicate determinations. Data in the same column bearing different superscripts differ significantly  $P=0.05$*

### 3.5 Physical Properties of Wheat, Soybean and Orange Fleshed Sweet Potatoes Cookies

“The mean values of physical characteristics of cookies prepared from composite flour blends of wheat soybean and OFSP are presented in Table 7. The result showed that the physical characteristics of the prepared cookies varied with the variation in the proportion of individual flours in the different blends. Similar observations have been reported by other authors” [59]. The highest weight was observed in samples with 5 % OFSP while 20% OFSP had the least value. The result obtained in this study where lower than 16.62 - 20.14g for rice, unripe banana and sprouted soybean cookies by [60]. This observation agrees with the reports by [20]. The recorded lower weight of cookies with 5% OFSP flour addition could be due to higher fat content of the soybean flour relative to other flours.

The diameter obtained for all samples were similar to the value (3.91 – 4.20) obtained by [60] for rice, unripe banana and sprouted soybean cookies. The diameter obtained from the study also followed a similar trend as the weight (50.2 - 57.3 mm) for composite wheat flour and malted barley cookies reported by [61]. This could be attributed to the amount of fat added to the flour blends during production. Similarly, an increasing trend for the diameter (38.90 – 40.20 mm) of cookies made from wheat brewers spent grain flour blends was reported by [62].

“The thickness obtained for all samples were similar to the results (0.73cm - 0.91cm) obtained for unripe banana and sprouted soybean cookies by [60]. Variations in cookies diameter and thickness are reflected in the spread ratio. Increase in thickness could be due to the high adsorption of moisture of the dough with increase in malted barley bran level, owing to the presence of water binding components. Moreover, it could also be attributed to the inconsistent rolling thickness of the dough which was exhibited as a result of the addition of high fat content” [61]. A similar finding for thickness was reported by [63] for cookies produced from blends of wheat flour and oat bran. However, [64] reported that thickness of cookies

developed from OFSP supplemented with wheat flour was reduced from 6.89 to 6.50 mm with addition of more OFSP during the cookie development. They considered that the heat applied to cookies during baking decreased the thickness of cookies with more OFSP than wheat.

The spread ratio obtained in this study is in contrast with the report obtained by [34] who opined that the average spread ratio of wheat orange fleshed sweet potatoes cashew nut cookies showed significant decreased as the proportion of orange flesh sweet potato and cashew nuts flour was increased in the formulation. In their discovery, they suggested that the spread ratio of cookies is strongly correlated to the water absorption capacities of flour. Further research by [60] revealed that “spread ratio of cookies has long been used as an important characteristic for determining the quality of flour for cookies production. Cookies with higher values of spread ratio are considered to be more desirable than those with lower values. Results obtained were similar to values (4.67 – 6.38) reported by [64] for wheat-Bambara protein isolate-ripe banana mash cookies”. However, [61] reported significant ( $P = .05$ ) increase in the spread ratio of the cookies with increasing level of malted barley bran (MBB) substitution from 0 to 50%, an observation [20] agreed with. According to [60] discovery, “doughs with lower viscosity cause cookies to spread at faster rate and vice versa”. They documented that the spread ratio of cookies increased with increase in the content of non-wheat protein. Increase in spread ratio could also be attributed to increase in the hydrophilic sites in the dough mixture leading to increase in water absorption and swelling index” [66]. “The control of cookies spread ratio is a serious problem encountered during production; cookies that spread so much cannot be filled into the package and those that spread slightly causes slack fill or excess height for package, thus creating havoc on the packaging line” [61].

**Table 7: Physical properties of wheat, soybean and orange fleshed sweet potatoes cookies**

W:S:OFSP (%)	Weight (g)	Diameter (cm)	Thickness (cm)	Spread ratio
100:0:0	7.85 <sup>e</sup> ±0.01	4.51 <sup>e</sup> ±0.02	0.90 <sup>a</sup> ±0.00	5.01 <sup>e</sup> ±0.02
75:20:5	8.11 <sup>d</sup> ±0.01	4.65 <sup>d</sup> ±0.04	0.86 <sup>b</sup> ±0.01	5.37 <sup>d</sup> ±0.00
70:20:10	8.52 <sup>c</sup> ±0.02	4.77 <sup>c</sup> ±0.03	0.81 <sup>c</sup> ±0.01	5.85 <sup>c</sup> ±0.01
65:20:15	9.17 <sup>b</sup> ±0.01	4.96 <sup>b</sup> ±0.08	0.77 <sup>d</sup> ±0.01	6.44 <sup>b</sup> ±0.01
60:20:20	9.45 <sup>a</sup> ±0.01	5.36 <sup>a</sup> ±0.02	0.77 <sup>d</sup> ±0.01	6.95 <sup>a</sup> ±0.09

*Values are means ± standard deviations of triplicate determinations. Data in the same column bearing different superscripts differ significantly  $P=0.05$*

#### 4. CONCLUSION

Functional properties such as bulk density and water absorption increased upon incorporation of OFSP and soybean flours while a decrease was observed in the swelling index. Proximate **composition** of flour had high protein, ash, fat and fibre contents due to OFSP and soybean substitution however, a decrease in carbohydrate content was observed. The antinutritional properties of the flour and cookies increased upon incorporation of soybean and OFSP however, the overall values obtained were below the critical limits as shown in the molar ratio. The physical properties of the fortified cookies were affected by the increased amount of OFSP.

## REFERENCES

1. Okpala L, Okoli E, Udensi E. Physico-chemical and sensory properties of cookies made from blends of germinated pigeon pea, fermented sorghum, and cocoyam flours. *Food Sci Nutr*. 2013(1):8-14.
2. Dauda A, Abiodun O, Arise A, Oyeyinka S. Nutritional and consumers acceptance of biscuit made from wheat flour fortified with partially defatted groundnut paste. *LWT-Food Science and Technology*. 2018(90):265-269. DOI:10.1016/J.LWT.2017.12.039. Corpus ID: 90344813. <https://api.semanticscholar.org/CorpusID:90344813>.
3. Hess J, Slavin J. Snacking for a cause: nutritional insufficiencies and excesses of U.S. children, a critical review of food consumption patterns and macronutrient and micronutrient intake of U.S. children. *Nutrients*. 2014 Oct 30;6(11):4750-9. doi: 10.3390/nu6114750. PMID: 25360509; PMCID: PMC4245561.
4. Ubbor SC, Arukwe DC, Ejechi, ME, Ekeh JI. Physiochemical And Sensory Evaluation of Cookies Produced from Composite Flours of Wheat, Bambara Nut and Orange Fleshed Sweet Potato. *J Agric Fd Sci*. 2022:(20)1:60 – 77.
5. Jukić M, Nakov G, Komlenić DK, Vasileva N, Šumanovac F, Lukinac J. Quality Assessment of Cookies Made from Composite Flours Containing Malted Barley Flour and Wheat Flour. *Plants (Basel)*. 2022;12;11(6):761. doi: 10.3390/plants11060761. PMID: 35336642; PMCID: PMC8948820.
6. Kumar N, Khatkar BS, Kaushik R. Effect of reducing agents on wheat gluten and quality characterisation of flour and cookies. *Journal of Food Science and Technology*. 2014. DO - 10.1007/s13197-013-1224-3.
7. Azeez L.A., Adedokun S. O, Elutilo O. O. and Alabi A. O. (2021). Quality attributes of cookies produced from the blends of sorghum, unripe plantain and watermelon seed flours. *International Journal of Research - Granthaalayah*, 2021: 9(2):309-319. <https://doi.org/10.29121/granthaalayah.v9.i2.2021.3565>
8. Adelakun OE, Duodu KG, Buys E, Olanipekun BF. Potential Use of Soybean Flour (Glycine max) in Food Fortification, Soybean - Bio-Active Compounds, Hany A. El-Shemy, IntechOpen, 2013. DOI:10.5772/52599 (February 20th 2013).
9. Sibiya H, Bhagwat P, Amobonye A, Pillai S. Effects of flaxseed and soybean supplementation on the nutritional and antioxidant properties of mahewu—A South African beverage. *South African Journal of Botany*, 2022(150):275–284. [CrossRef].
10. Yu X, Meenu M, Xu B, Yu H. Impact of processing technologies on isoflavones, phenolic acids, and antioxidant capacities of soymilk prepared from 15 soybean varieties. *Food Chemistry*. 2021:345. 128612. [CrossRef].
11. Helen Keller international (HKI). Orange fleshed sweet potato situation analysis and needs assessment international potato centre, 2012.
12. Laurie S, Van Jaarsveld P, Faber M, Philpott M, and Labuschage M. Trans-β-carotene, selected mineral content and potential nutritional contribution of 12 sweet potato varieties. *Journal of Food Composition and Analysis*, 2012(27):151-159.

13. Kurabachew H. The role of orange fleshed sweet potato (*Ipomea batatas*) for combating vitamin A deficiency in Ethiopia: A review. *International Journal of Food Science and Nutrition Engineering*. 2015;5(3), 141 – 146.
14. Shiriki D, Igyor MA, Gernah DI. Nutritional Evaluation of Complementary Food Formulations from Maize, Soybean and Peanut Fortified with Moringa oleifera Leaf Powder. *Food and Nutrition Sciences*. 2015: 6(1), 494–500
15. Kudadam J, Solomon K, Chikpah K, Hensel O, Pawelzik E, Sturm B. Effect of orange - fleshed sweet potato flour particle size and degree of wheat flour substitution on physical, nutritional, textural and sensory properties of cookies. *European Food Research and Technology*. 2021: 247(4), 889–905. <https://doi.org/10.1007/s00217-020-03672-z>.
16. Bashir A, Ashraf AS, Khan MA, Azad AA. Development and Compositional Analysis of Protein Enriched Soybean-Pea-Wheat Flour Blended Cookies. *Asian Journal of Clinical Nutrition*. 2015;7(3): 76-83.
17. Onwuka GI. Food analysis and instrumentation. Theory and practices. Revised edition. Naphtali Prints Lagos, Nigeria. 2018: 10-20.
18. AOAC. Official Methods of Analysis. Association of Official Analytical Chemists. 18th Edition, AOAC, Arlington. 2015: 806-814.
19. Norhaizan M, NorFaizadatul AW. Determination of phytate, iron, zinc, calcium contents and their molar ratios in commonly used raw and prepared food in Malaysia. *Malays. J. Nutr*. 2009;15 (2), 213–222.
20. Dabel N, Igbabul BD, Julius A, Iorliam B. Nutritional composition, physical and sensory properties of cookies from wheat, acha and mung bean composite flours. *International Journal of Nutrition and Food Science* 2016: 5(6):401-406, 2016.
21. Chandra S, Singh S, Kumari D. Evaluation of functional properties of composite flours and sensorial attributes of composite flour biscuits. *J Food Sci Technol*. 2015;52(6): 3681–3688.52(6):3681–3688. DOI 10.1007/s13197-014-1427-2. PMCID: PMC4444897. PMID: 26028751
22. Okafor J, Obiegbuna N, James E, Agu HO. Functional and Pasting Properties of Composite Flour from Wheat, Sweet Potato and Soybean. *International Journal of Environmental and Agriculture Research (IJOEAR)*, 2021:(7):2454-1850, Issue-12, December- 2021].
23. Kanu NA, Afuape SO, Ezeocha VC, Nwafor JO. Proximate Composition and Functional Properties of Improved Orange-Fleshed Sweetpotato Breeding Lines Developed In Umudike, Abia State. *Nigerian Agricultural Journal*. 2018: (49):125-133 No. 1, April 2018. ISSN: 0300-368X. Available online at: <http://www.ajol.info/index.php/naj>.
24. Menon L, Majumdar SD, Ravi U. Development and analysis of composite flour bread. *Journal of Food Science and Technology*, 2015;52 (7): 4156 – 4165.
25. Klunklin W, Savage G. Physicochemical, antioxidant properties and in vitro digestibility of wheat–purple rice flour mixtures. *International Journal Food Science and Technology*, 2018:(53):1962–1971. [CrossRef]

26. Culetu A, Susman IE, Duta DE, Belc N. Nutritional and Functional Properties of Gluten-Free Flours. *Applied Sciences*. 2021; 11(14):6283. <https://doi.org/10.3390/app11146283>
27. Patil SP, and Arya SS. Nutritional, functional, phytochemical and structural characterization of gluten-free flours. *Journal Food Measurement and Characterization*, 2017, 11, 1284–1294. [CrossRef]
28. Chandra S. Assessment of functional properties of different flours. *Afr. J. Agric. Res.* 2013;8(38):4849–4852. [Google Scholar]
29. Onimawo IA and Akubor PI. *Food Chemistry (Integrated Approach with Biochemical background)*. 2012. 2nd ed. Joytal printing press, Agbowo, Ibadan, Nigeria.
30. Etudaiye HA, Emmanuel O, Aniedu C, Majekodunmi OR. Utilization of sweet potato starches and flours as composites with wheat flours in the preparation of confectioneries. *Afri J Biotechno* 2014;14(1):17-22. DOI:10.5897/AJB12.2651
31. Olaitan NI, Eke MO, Uja EM. Quality Evaluation of Complementary Food Formulated from Moringa Oleifera Leaf Powder and Pearl Millet (*Pennisetum glaucum*) Flour. *The International Journal of Engineering and Science*. 2014;3 (11): 59 – 63.
32. Schirmer M, Jekle M, Becker T. Starch gelatinization and its complexity for analysis. *Starch* 2015;67:30–41. [CrossRef]
33. Jhan, F., Gani, A., Shah, A., Ashwar, B. A., Bhat, N. A. And Ganaie, T. A. (2021). Gluten-free minor cereals of Himalayan origin: Characterization, nutraceutical potential and utilization as possible anti-diabetic food for growing diabetic population of the world. *Food Hydrocolloids*, 2021, 113, 106402. [CrossRef]
34. Donaldben N.S, Amanze L.C and Ewelike I.C (2019). Quality Characteristics of Cookies from Wheat and Orange Flesh Sweet Potatoes Composite Flour Supplemented with Cashew Nut Powder.
35. Kisambira, A., Muyonga, J. H., Byaruhanga, Y. B., Tukamuhabwa, P., Tumwegamire, S. And Grüneberg, W. J. (2015). Composition and functional properties of Yam bean (*Pachyrhizus spp.*) seed flour. *Food and Nutrition Sciences*, 6: 736 – 746.
36. Shad, A. M., Nawaz, H., Noor, M. and Ahmad, H. B. (2013). Functional properties of maize flour and its blends with wheat flour: optimization of preparation conditions by response surface methodology. *Pakistan Journal of Botany*, 45 (6): 2027 – 2035.
37. Verem TB, Dooshima IB, Ojotu EM, Owolabi OO, and Onigbajumo A. Proximate, Chemical and Functional Properties of Wheat, Soy and Moringa Leaf Composite Flours. *Agricultural Sciences*. 2021;12:18-38. <https://doi.org/10.4236/as.2021.121003>
38. Nour, A. A. M. and Ibrahim, M. A. E. M. (2016) Effect of Supplementation with Moringa Leaves Powder (MLP) and Fermentation on Chemical Composition, Total minerals Contents and Sensory Characteristics of Sorghum Flour. *International Journal of Science Resource*, 5, 672-677. <https://doi.org/10.21275/v5i3.NOV161822>
39. Iwe MO, Michael N, Madu NE, Obasi NE, Onwuka GI. Physicochemical and pasting properties high quality cassava flour (HQCF) and wheat flour blends. *Agrotechnology*. 2017;6:167.

40. Barber LI, Obinna-Echem PC, Ogburia EM. Proximate composition micronutrient and sensory properties of complementary food formulated from fermented maize, soybeans and carrot flours. *Sky Journal of Food Science*. 2017;6 (3): 33 – 39.
41. Oyarekua MA. Effect of co-fermentation on nutritive quality and pasting properties of maize/cowpea/sweet potato as complementary food. *African Journal of Food Agriculture, Nutrition and Development*. 2013;13 (1): 7171 – 7191.
42. Damian L. Formulation and characterization of sweetpotato-based complementary food. Master of Philosophy (M.Phil) Thesis, Department of Food Science and Technology, Kwame Nkrumah University of Science and Technology, Kumasi College of Science, Ghana. 2016: 123 pp.
43. Okoye, J. I. and Ojobor, C. C. (2016). Proximate composition, energy content and sensory properties of complementary foods produced from blends of sorghum and African yam bean flour. *International Journal of Scientific and Technology Research*, 5 (7): 274 – 277.
44. Oloniyo RO, Omoba OS, Awolu OO. Biochemical and antioxidant properties of cream and orange-fleshed sweet potato. *Heliyon*, 7 (2021) e06533. <https://doi.org/10.1016/j.heliyon.2021.e06533>. Published by Elsevier Ltd
45. Banigo BE, Kiin-Kabari DB. Effect of African Yam Bean (AYB) (*Sphenostylis stenocarpa*) on the quality characteristic of extended meat ball. *Journal of Food and Nutrition Research*, 2016: 4 (2): 121 – 125.
46. Desalegn BB, Abegaz K, Kinfe E. Effect of Blending Ratio and Processing Technique on Physicochemical Composition, Functional Properties and Sensory Acceptability of Quality Protein Maize (QPM) Based Complementary Food. *International Journal of Food Science and Nutrition Engineering*. 2015;5 (3): 121-129.
47. Okoye, J. I. and Obi, C. D. (2017). Chemical Composition and Sensory Properties of Wheat-African Yam Bean Composite Flour Cookies. *Discourse Journal of Agriculture and Food Sciences*, 5(2): 21-27.
48. Alam M, Rana Z, Islam S. Comparison of the proximate composition, total carotenoids and total polyphenol content of nine orange-fleshed sweet potato varieties grown in Bangladesh. *Foods*. 2016;5 (3), 64.
49. Ukegbu PO, Uwaegbule AC, Nnadi CM. Organoleptic and nutritional evaluation of African yam bean (*Sphenophyllis stenocarpa*) flour enriched complementary food. *Scholars Journal of Agriculture and veterinary Sciences*, 2015: 2 (2A): 97 – 101.
50. Ikujenlola AV, Adurotoye EA. Evaluation of quality characteristics of high nutrient dense complementary food from mixtures of malted quality protein maize (*Zea mays* L.) and steamed cowpea (*Vigna unguiculata*). *Journal of Food Process Technology*, 2014: 5 (1): 1 – 5.
51. Achidi AU, Tiencheu B, Tenyang N, Womeni HM, Moyeh MN, Ebini LT, Tatsinkou F. Quality evaluation of nine instant weaning foods formulated from cereal, legume, tuber, vegetable and crayfish. *International Journal of Food Science and Nutrition Engineering*, 2016: 6(2), 21–31. [Google Scholar]
52. Dako E, Retta N, Desse G. Comparison of three sweet potato (*Ipomoea Batatas* (L.) Lam) varieties on nutritional and anti-nutritional factors. *Global Journal Science of Frontier Research, (GJSFR): D Agriculture and Veterinary*. 2016;16 (4), 7–19.

53. Tiruneh Y, Urga K, Tassew G, Bekere A. Biochemical compositions and functional properties of orange-fleshed sweet potato variety in Hawassa, Ethiopia. *American Journal of Food Science and Nutritional Research*. 2018;5 (1), 17.
54. Dada MA, Bello FA, Omobulejo FO Olukunle FE. Nutritional quality and physicochemical properties of biscuit from composite flour of wheat, African yam bean and tigernut. *Heliyon*, 9 (2023) e22477. <https://doi.org/10.1016/j.heliyon.2023.e22477>.
55. Nzeagwu OC, Onwudiwe ET. Nutrient composition and sensory properties of tigernut (*Cyperus esculentus*) biscuit. *Nigerian Journal of Nutritional Science*, 2016;37 (1):157–164.
56. Haile A, Getahun D. Evaluation of nutritional and anti-nutrition factors of orange-fleshed sweet potato and haricot bean blended mashed food for pre-school children: the case of Dale Woreda, Southern Ethiopia. *Food Science and Technology*. 2018;6 (1), 10–19.
57. Connorton JM, Balk J, Rodríguez-Celma J. Iron homeostasis in plants—a brief overview. *Metallomics*. 2017;9 (7), 813–823.
58. Laelago, T., Haile, A. and Fekadu, T. Production and quality evaluation of cookies enriched with  $\beta$ -carotene by blending orange-fleshed sweet potato and wheat flours for alleviation of nutritional insecurity. *International Journal of Food Science and Nutrition Engineering*, 2015;5:209-217.
59. Man S, Paucean A, Muste S. Preparation and quality of gluten-free biscuit. *Bull UASVM Food Science and Technology*. 2014;71(1):39-41, 2014.
60. Inyang UE, Daniel, EA, Bello FA. Production and quality evaluation of functional biscuits from whole wheat flour supplemented with acha (fonio) and kidney bean flours. *Asian Journal of Agriculture and Food Sciences*. 2018;6(6):193-201.
61. Ikuomola DS, Otutu OL, Oluniran DD. Quality assessment of cookies produced from wheat flour and malted barley (*Hordeum vulgare*) bran blends, *Cogent Food and Agriculture*, 2017;3:1. DOI: 10.1080/23311932.2017.1293471.
62. Gernah, D. I., Ariahu, C. C. and Umeh, U. (2012). Physical and Microbiological Evaluation of Food Formulations from Malted and Fermented Maize (*Zea mays* L.) Fortified with Defatted Sesame (*Sesamun indicum* L.) Flour. *Advance Journal of Food Science and Technology* 4(3): 148 – 154.
63. Abdul WK, Javid A, Tariq M, Muhammad A, Mohammad P, Said H. Effect of oat bran on the quality of enriched high fibre biscuits. *World Journal of Dairy and Food Sciences*. 2015;10;68–73. Google Scholar.
64. Shazia S, Muhammad M, Ahmad H, Kausar S, Parveen SM, Abdusalam. Effect of sweet potato flour on quality of cookies. *Journal of Agriculture Research*, 2012;50(4): 1 - 3.
65. Arise AK, Akeem SA, Olagunju OF, Opaleke OD, Adeyemi DT. Development and Quality Evaluation of Wheat Cookies Enriched with Bambara Groundnut Protein Isolate alone or in Combination with Ripe Banana Mash. *Applied Food Research*, Volume 1, Issue 1, 2021, 100003, ISSN 2772-5022, <https://doi.org/10.1016/j.afres.2021.100003>. (<https://www.sciencedirect.com/science/article/pii/S2772502221000032>)

66. Igbabul BD, Iorliam BM, Umana EN. Physicochemical and sensory properties of cookies produced from composite of wheat, cocoyam and African yam bean. *Journal of Food Research*. 2015;4(2):150-158, 2015.

UNDER PEER REVIEW