

## Original Research Article

### **Spillage of Akosombo and Kpong dams in Ghana: perspectives on public health impacts on affected populations and proposed mitigation strategies**

#### **ABSTRACT**

**Background:** The negative impact of floods on humans and the environment cannot be overemphasized. Annually, different parts of Ghana get flooded resulting in the loss of lives and significant damage to property.

**Objective:** This study assessed the public health impacts of the controlled spillage of the Akosombo and Kpong dams in Ghana on people living in downstream and upstream communities.

**Methods:** This was an observational study conducted in the affected communities from September 15 and October 30, 2023. Government reports, published media reports, NGO publications, and field visits were also used to assess the extent of damage, challenges faced by local communities, and mitigation measures initiated.

**Results and Discussion:** The spillage led to the loss of critical infrastructure such as schools, houses, toilet facilities, water supply systems and electricity. Overcrowding in temporary shelters provided for displaced persons could serve as a conduit for the transmission of communicable disease. The suspension of healthcare services in areas inundated by flood waters worsened the plight of residents including vulnerable members of the society such as pregnant women, children, the aged, persons living with disability and the very poor. The submersion of farmlands, crops and drowning of livestock and poultry had serious implication for food security and livelihoods in the affected areas. The current study comprehensively captured the scale of devastation of the floods on people and their livelihoods in 7 out of the 16 regions within 21 administrative districts across Ghana, and proposed workable strategies to mitigate future happenings.

**Conclusion:** The public health impacts resulting from the controlled spillage of the Akosombo and Kpong Dams on affected individuals was enormous. Possible surge in vector-borne disease transmission such as

malaria, dengue, zika, and a probable increase in communicable and non-communicable diseases. Although short term measures were adopted to mitigate the impact of the flood on affected individuals through the provision of clean water, beddings, emergency food relief, long-term resilience strategies including early warning systems, climate sensitive interventions, effective collaboration among key stakeholders and disaster preparedness at the local level will help avert or reduce the severity of future floods. The study underscored the urgent need for comprehensive measures to mitigate the impacts of future disasters and enhance community resilience.

**Keywords:** Floods, Volta River Authority, Akosombo dam spillage, public health impacts, vector-borne diseases

## 1. INTRODUCTION

The research carried out by the International Flood Initiative in 2003 highlighted the significant impact of floods, which represent the predominant category of water-related natural disasters. These floods not only result in considerable harm to human lives and property but also have far-reaching consequences for cultural and ecological resources [1]. Causes of floods can be categorized into two main groups, as noted by Nott in 2006 [2] namely, natural factors, often driven by climate forces, and human-induced factors, which include urban development and vegetation clearance. While natural factors play a significant role in causing floods globally, changes in flood patterns are primarily attributed to human activities, such as deforestation [3].


Floods are not classified as natural disasters until they result in damage to human lives or property. However, their adverse effects can be mitigated through the implementation of advanced warning systems, as emphasized by Sinclair and Pegram in 2003 [4]. Additionally, many economically disadvantaged individuals live near riverbanks, often due to the lack of alternative housing options. Unfortunately, these residents not only face a higher risk of flooding due to their locations but also due to their limited financial resources [5].

The impact of floods on human beings is of utmost significance, as it extends its reach across various aspects of human existence. This encompasses the physical environment, human health, and agricultural

produce [6]. Depending on its magnitude and speed, flooding has the potential to harm a wide range of structures, such as bridges, vehicles, buildings, sewerage systems, roads, and canals. Additionally, it can lead to water contamination [7].

In Ghana, the Akosombo and Kpong hydro-electric power plants located on the Volta River Basin are owned and operated by the Volta River Authority (VRA) [8]. The power generating capacity of the Akosombo and Kpong dams are 1,020MW and 160MW respectively [9]. One of the most notable outcomes of the Akosombo hydroelectric power plant is its capacity to supply electricity to neighboring countries, strengthening regional ties and fostering economic cooperation. Ghana extends its electrical generosity to nations such as Togo, Benin, and Cote d'Ivoire, offering a valuable resource for domestic and commercial purposes [10]. This interconnectedness through electricity exchange serves as a testament to the VRA's pivotal role not only in Ghana but also in bolstering the energy security and economic growth of its neighboring countries.

This study assessed the negative impacts of the 2023 controlled spillage of the Akosombo and Kpong dams, resulting from increased water inflow, with severe impacts on downstream communities, and the urgent need for comprehensive measures to mitigate the impacts of future disasters and enhance community resilience. The Akosombo hydroelectric dam and the opening of the the spillways are shown in Fig. 1.a and Fig. 1.b respectively.



**Fig. 1.a** Akosombo dam constructed in 1965



**Fig. 1.b** Akosombo dam opens spillways.

## **2. METHODS**


### **2.1 Study design**

This was an observational study conducted in the affected areas from September 15 to October 30, 2023. Government reports, published media reports, NGO publications, and field visits were also used to assess the extent of damage, challenges faced by local communities, and mitigation measures initiated.

### **2.2 Study area**

Ghana, the study area, is a country in West Africa located between latitudes 4.5°N and 11.5 °N and longitude 3.5 °W and 1.5 °E (see Fig 2). Ghana is bordered to the east by the Republic of Togo, to the west by the Republic of Cote d'Ivoire, to the north by the Republic of Burkina Faso, and to the south by the Gulf of Guinea. Tropical monsoon typically defines the climatic condition of Ghana and about 65% of the land is

under agricultural use [11, 12]. The West African Monsoon (WAM) controls the rainfall in Ghana, where the atmospheric pressure system influenced by the temperature gradient and energy between the Sahara and the Gulf of Guinea [13]. The Inter-Tropical Discontinuity (ITD) modulates this atmospheric pressure system and the oscillatory motion of the ITD between the south and the north of Ghana results in the bi-modal and uni-modal rainfall distribution across the country [14, 15]. Uni-modal rainfall is experienced in northern Ghana from April through to mid October and peaks in August or September. However, southern Ghana is characterized by bi-modal rainfall distribution. First rainfall occurs between March to July and peaks in June whereas the second occurs between September and mid-November and peaks in October. Several factors which influence rainfall pattern in the country include sea level temperature, local convective activity and atmospheric dynamic stability [16]. Ghana experiences the harmattan as a result of the north-east desert wind between December and February. The harmattan which results in low humidity, hotter days and cooler nights is prevalent in the northern part of Ghana compared to the southern part.



**Fig 2: Geography of Ghana Credit: [13]**

Ghana is categorized into five distinct climatic zones due to the significant changes in climatic variables such as rainfall and temperature over time. These distinct climatic zones include; the Sudan Savannah, Guinea Savannah, Transition, Forest and Coastal zones [17]. The new zonation is a result of the expansion in the size of Sudan and Guinea Savannah southwards, shrinking of the Transition zone as the Guinea Savannah took over a greater portion of it in the southeastern part of Ghana. Additionally, the shrinking of the Forest zone in size together with the shift in the northwest, while the coastal belt has expanded significantly in size to cover the entire coast of Ghana. These changes are strong indications of climate change and its possible impact on weather patterns.

### 2.3 Source of data

We conducted field visits and extensive review of existing literature, including academic papers, electronic media reports, government reports, and non governmental organization (NGO) publications. This provided a baseline understanding of the issue and the existing mitigation efforts. The field visits were conducted in areas directly impacted by the spillage of the dams to ascertain the conditions and challenges faced by local communities, assess the extent of damage, and identify ongoing mitigation measures.

The literature search strategy consisted of five key elements: Akosombo dam spillage, floods in Ghana, climate change, effects of climate change, and man-made disasters. Search terms for 1) Akosombo dam spillage was combined with search terms for 2) floods in Ghana (including regions and districts). Search terms for 3) climate change was combined with search terms for 4) man-made disasters and 5) effects of climate change. These combinations were necessary to compile a comprehensive search strategy to address the aim of the study. Academic literature databases (Pubmed, Web of Science, Google Scholar), Websites of Media outlets in Ghana (Myjoyonline.com, Citinewsroom.com, Ghana News Agency, Ghana Broadcasting Corporation); Websites of government agencies, online search engines such as Google for “grey” literature.

### 2.4 Data management

Finalized search strings and references were used to search electronic databases. The search results were imported into Mendeley Desktop (version 1.19.4, Mendeley Ltd., London, United Kingdom) for storage and the removal of duplicates (Christopher Yaw Dumevi, Christopher Mfum Owusu-Asenso, Bright Darko Amoah). The inclusion and exclusion criteria guided the search, selection and review of the relevant literature such as abstracts, commentaries, news reports and full articles. Three author-reviewers independently screened the literature (Christopher Yaw Dumevi, James-Paul Kretchy, Nicholas Tete Kwaku Dzifa Dayie) and a fourth independent reviewer (Patrick F. Ayeh-Kumi) resolved any conflicts. Irrelevant literature was excluded paving the way for the compilation of a final list of literature included in this study. There was no grading of included literature.

**Participant involvement:** This was an observational study with no human subject participation directly or indirectly.

**Data analysis:** Due to the heterogeneity in the study design, study settings, and impact of the flood on affected individuals, a thorough narrative synthesis was done to address the objective of the study. The findings of the study were graphically represented and tabulated highlighting the region, district and communities that were negatively impacted.


### 3. RESULTS

The Volta River Authority on September 15, 2023 began a controlled spillage of the Akosombo and Kpong dams as a result of a consistent inflow of water into the Akosombo dam. The water level prior to the spillage was 272.50 ft (83.058m) to protect the dam from imminent collapse. This necessitated the opening of the spillways of the Akosombo and Kpong dams between September 15 and October 30, 2023 which led to severe flooding in downstream communities along the Volta River Basin. Water and electricity supply to affected communities were cut, personal belongings, and domestic animals were washed away by the raging water levels leaving many affected individuals without food and potable water for days .

The shutting down of the Ghana Grid Company (GRIDCo) sub-station (electricity power distributor) at Fievie near Sogakope in the South Tongu district of the Volta region severely hampered healthcare delivery in Sogakope and Adidome hospitals. Some districts severely impacted by the floods include, North Tongu, Central Tongu, South Tongu, Anlo, and Keta all in the Volta Region; Asuogyaman and Ada districts in the Eastern and Greater Accra Regions respectively. Some communities in the Keta and Anloga were also flooded due to the overflow of lagoons in the Keta basin as a result of the Akosombo dam spillage. Although VRA has spilled water from the Akosombo dam in the past, the current spillage was most devastating compared to the impact of previous spillages in 1966, 1967, 1968, 1969, 1970, 1971, 1972, 1974, 1991, and 2010 [18].

An estimated 150,000 people have been affected by the flooding directly or indirectly with many homes, personal belongings and livelihoods lost due to the floods [19]. Latest records revealed flooding in 21 administrative districts within seven (7) regions which lie along the upper and lower Volta basin due to increasing flow of water from the Bagre Dam in Burkina Faso into the Akosombo and Kpong dams

resulting in spillage of excess water [19]. The widespread devastation is shown in Fig. 3. Although the impact of the spillage was felt largely along the lower volta basin, other areas upstream were severely impacted (**Table 1**).



**Fig. 3.** Areas affected by the Akosombo and Kpong dam spillage.

**Table 1. Devastating effect of the spillage of the Bagre dam (in Burkina Faso), and Akosombo and Kpong dams (in Ghana) on Ghanaian communities.**

Region	District /Municipal	Some affected Communities
Bono East	Pru East	Yeji, Futudeke, and environs
Eastern	Asuogyaman	Abume, Kudikope, Kokonte Kpedzi, Ahenbrom, Dzidzokope, Mama Kope and environs
	Lower Manya Krobo	Akuse and environs
Greater Accra	Shai Osodoku	Asutsuare, Blonya, Ashimi, Fawkpi, Avakpo, Volivo Landor, Alabonu, Kewum Atrobinya, Kesegakope, Torgome, and environs
	Ada East	Ada, Agavedo, Tuanikope Island, Anazome, parts of Ada Foah, Big Ada and communities at the estuary
Northern	Kpandai	Lonto, Sika Kura, no 1&2, Kpadjai, Gyeakope, Tetekura, Vuvukope, and environs
Oti	Biakoye	Tapa-Abotoase, Kwamekrom, Obosumanu, Fahiakoho, Bumbula, and environs
	Krachi East	Dambai, Kove Kope, Cannab, Mepe Kope, Teflekordzi, Kudorkope, Kekpodzi, Wulubu, Adakponu, Korvekope, Asikanfo-Ambatem, Wulubu,
	Krachi West	Kekte-Krachi, and environs
	Krachi Nchumuru	Bagamsi, Buafori, D/A Primary and Junior High School at Papatia Grubi, Anyinamae and environs

	Nkwanta South	Kabiti, Odumasi and environs ,
	Nkwanta North	Kabonwule, Damanko
Savannah	Central Gonja	Buipe and environs
Volta	Kpando	Kpando Tokor and environs
	Anloga	Parts of Anloga township, Anloga market& lorry park, Atsito, and environs
	Keta	Adzido, Keta central, Galosota, Fiakor, Kedzi, Azizadzi, Keta Business College (Ketabusco), Anyako, and environs
	Ketu South	Havedzi, Blekusu, Adina, Horvi, Amutinu
	South Dayi	Tongor-Dzemeni and environs
	Central Tongu	Adidome, Afetorgborkope, Mafi Dugame, Kebedogo, Akpokofe, Devime, Bekpo Avadimewoe-kope, Akpokope, Kebegodo, Siamekope, Atsemkope, Dokpoe and environs
	North Tongu	About 90% of Mepe township flooded,  Battor Aveyime,
	South Tongu	Fievie, Sogakope, St. Comboni Catholic Hospital, Tefle, Vume, Sokpoe, Alikekope, Agorme, Agbave, Villa Cisneros, Sogakope Beach Resort and Spa, Holy Trinity Spa and Health Farm, Ahiatrogakope, Havorkope, Adadzikope, Agbokope, Sukladzi, Ashiagborkope, Tsinuto, and environs

#### 4. Discussion

The current study explored the devastation caused by the controlled spillage of the Akosombo and Kpong dams in Ghana that led to the displacement of about 150,000 people. This is in line with the report by Adegoke et al., [20] which stated that, more than 9,500 people in Sikasso, Kolikoro, Segou, Mopti, Timbuktu, Gao, Kidal and Menaka in Mali were affected by flooding. The Akosombo Dam, which historically stood as a linchpin of Ghana's electricity generation infrastructure, has become a focal point of despair and devastation due to controlled spillage of excess water. Critical social infrastructure such as schools, hospitals, water treatment facilities, electricity sub-stations, roads, bridges among others were severely affected by the floods. Taylor and Davies [21] reported similar negative impacts of frequent flooding on roads, railway lines, electricity supply, and communication infrastructure in Cape Town, South Africa. In Vietnam, Duy et al., [22] reported that, transport systems are frequently inundated with flood waters leaving commuters either stranded or risk being swept away by the torrential current of the flood waters. Okuyama et al., [23] opined that, the destruction of social and physical infrastructure such as transport networks is the “direct damages” of flooding. The resultant effect is reduction in economic activities, leading to change production and or consumption behaviour as well as investment decisions.

In the South Tongu district of the Volta Region of Ghana, the Agordomi water treatment and distribution plant was shutdown leaving the inhabitants of Sogakope without potable water for weeks. Similar study conducted by Augusto & Santos, [24] on the operation of the Weija Dam in Accra reported the challenges of managing excess water during periods of heavy rainfall, leading to spillage and downstream flooding leading to contamination of water sources for domestic use. The negative environmental impact and loss of economic activities in downstream communities calls for a rethink in managing excess water solely by spillage according to Owusu-Ansah et al., [25]. The devastation caused by the flood waters in some affected areas are shown in Fig. 4



**Fig. 4.** devastation caused by the flood waters in some affected areas

**Key**

**A.** Flooded homes. **Photo credit:** myjoyonline.com

**B.** Flooded Dambai market, Oti Region. **Photo credit:** myjoyonline.com

**C.** St. Comboni Secondary Technical School Teachers' bungalow, South Tongu district  
26/10/2023 at 14:12 GMT **Photo credit:** Jerry Clinton Donkor.

**D.** Flooded homes. **Photo credit:** myjoyonline.com

**E.** Flooded toilet facility at Mepe, North Tongu District, Volta Region. **Photo credit:** Jerry  
Amenyo-Quarm 25/10/2023 at 8:07GMT

**F.** St. Comboni Secondary Technical School, Girls' lavatory, South Tongu district 26/10/2023 at  
14:09 GMT. **Photo credit:** Jerry Clinton Donkor.

The spread of waterborne diseases through flooding is a serious public health threat [26, 27]. The floodwaters, often a turbulent mixture of rainwater, sewage, and pollutants, serves as a breeding ground for a host of waterborne pathogens [25]. Moreover, the displacement of thousands of people from their homes created overcrowding conditions in temporary shelters, raising concerns about the potential spread of communicable diseases. Cholera and typhoid fever, which are transmitted through contaminated water as a result of insanitary and compromised environmental sanitation [28]. This finding agrees with Mirza, [29] and Dewan [30] who opined that, the pollution of water for domestic use, risk of the spread of waterborne diseases such as diarrhoea, typhoid, cholera, dysentery, and epidemics are commonly associated with flooding in Bangladesh and Nepal with overwhelming impact on the poor, women, children, and the elderly.

The forced displacement of individuals from their homes and communities, has profound psychological burden on many. The loss of homes had a deep emotional and psychological rift that could manifest as anxiety, depression, post-traumatic stress, and a pervasive sense of grief. The survivors grapple with a sense of emptiness, a feeling of being adrift, as they come to terms with the altered landscape of their lives. A study conducted by Fitzgerald et al., [31] revealed that, comparatively, resource limited-countries experience severe impacts of floods such physical trauma, anxiety, panic attacks than developed countries and that one-third of deaths associated with flooding in developing countries are linked to traumatic experience and heart attack.

These mental health challenges, often underestimated in the immediate aftermath of a disaster, persisted long after the floodwaters receded, requiring sustained attention and support to help individuals and communities rebuild not only their physical homes but also the sanctuary of their mental and emotional well-being.

The floodwaters submerged fields and pastures, destroying crops and livestock in affected communities across Ghana. The loss of these vital resources meant not only economic hardship but also a severe reduction in the availability of locally produced foods. Communities that relied heavily on subsistence farming were especially vulnerable particularly staples such as cassava, maize, banana, plantains, and groundnut. In the Anloga, Keta, and Ketu South districts of the Volta region, large tracts of vegetable farms;

carrot, cabbage, sweet pepper, and onions have all been submerged when the lagoon overflowed its banks as a result of increased volume of water in River Volta and its tributaries. With local food production severely hampered, affected populations are at risk of facing acute food shortages. Many individuals and families struggled to secure an adequate and nutritious diet, which can potentially lead to malnutrition. The destruction of crops, farmlands, and livestock due to flooding in Nepal has negative impact on the food security situation of the country. Also, starvation due to lack of food resulted in the death of day laborers in Bangladesh [30]. Malnutrition, therefore, can have long-lasting consequences on the physical and cognitive development, immune function, and overall well-being of children, pregnant women and the aged [20].

Vector-borne diseases (VBDs) such as malaria, dengue, Zika, and Lyme disease pose significant public health threats, particularly in resource-limited regions [33]. The surge in VBDs, exacerbated by flooding, is a pressing concern in areas like South Asia, where vulnerable populations are mostly affected [29]. Flooding, intensified by climate change, urbanization, and anthropogenic factors such as hydroelectric dam spillage, profoundly impacts the dynamics and epidemiology of VBDs [32, 34]. The formation of stagnant water bodies resulting from dam spillage provides ideal breeding habitats for vectors, particularly mosquitoes. These conditions can lead to a surge in mosquito vector populations, heightening the risk of diseases like malaria and dengue as the floodwaters recede [35, 36]. Additionally, the expansion of floodplains can facilitate the geographic spread of vector species, introducing VBDs to new areas [37]. The recent dam spillage in Ghana, affecting diverse ecological zones, highlights the potential for serious public health implications through the introduction of novel vector species in previously uncolonized areas.

Flooding not only boosts vector abundance but also increases vector-human contact [38]. Higher humidity levels and numerous water bodies enhance mosquito and tick activity, leading to more frequent disease transmission. This is particularly concerning for malaria and arboviral disease transmission, as communities remain at greater risk during the post-flood period [39]. Moreover, the displacement and crowding in emergency shelters during floods create conditions favorable for vector-human interactions, thus enhancing disease transmission [40, 41]. Damaged sanitation infrastructure and contaminated water sources further exacerbate the risk of waterborne diseases, which may co-occur with VBDs during floods. The potential for

outbreaks increases, necessitating stringent vector control measures and health education programs [42]. These findings emphasize the critical need for integrated vector management and proactive public health strategies to mitigate the impact of flooding on VBD transmission.

#### Lapses identified

- a. Ineffective community engagement and sensitization by Volta River Authority. Most affected communities were not aware of the spillage and had short time to evacuate or move to higher grounds. This led to the loss of valuable personal belongings and properties.
- b. Late response by the National Disaster Management Organization (NADMO) to affected communities. Although the Ghana Navy helped to evacuate about 8,000 people in some parts of the Volta Region, there was no such response in other affected regions. The late arrival of NADMO officials with some relief items days after victims were displaced exposed the vulnerability of many affected persons. Many island communities were completely cut off from the inland for weeks without any essential supplies.
- c. Ineffective collaboration among state institutions. That was necessary as to broaden consultation and sensitization of communities along the Volta Basins on the impending spillage of the Akosombo and Kpong dams. These key stakeholders include, the Volta River Authority, Ghana Meteorological Agency, National Commission on Civic Education, local administrative councils and other relevant stakeholders
- d. Lack of disaster preparedness by local government authorities..

#### Limitations of the study

The results of the study were based on available data as described in the methods and not from the affected individuals.

## 5. CONCLUSION

The devastating flood and the controlled spillage of the Akosombo and Kpong dams in September 2023 had profound and far-reaching impacts on the affected populations and regions. The floods affected critical infrastructure, with overarching effect on education, food security, healthcare, energy, business among others. This study recommends the strengthening of existing collaboration between VRA and key stakeholders to avert the negative impact of any future controlled spillage. We advocate a comprehensive mental health assessment and immediate psychological intervention for all survivors in the affected communities. This would be one giant step to getting the lives of affected individuals back to normalcy.

## REFERENCES

1. European Union (EU) Floods directive (2007) DIRECTIVES, date accessed 16th August 2012 from <http://eurlex.europa.eu/LexUriServ/LexUriServ.do?uri=OJ:L:2007:288:0027:0034:EN>: **Google Scholar CrossRef**
2. Nott J. *Extreme Events: A Physical Reconstruction and Risk Assessment*. Cambridge. 2006 University Press. New York. Available from: <https://www.preventionweb.net/publication/extreme-events-physical-reconstruction-and-risk-assessment> **Google Scholar**
3. Odunnuga S, Oyebande L, & Omojola AS. Socio-economic Indicators and Public Perception on Urban Flooding in Lagos Nigeria", Special Publication of the *Nigeria Association of Hydrological Science*. 202 Available from: <http://www.unaab.edu.ng/Nov.2013>. **Google Scholar**
4. Sinclair S, & Pegram G. A flood nowcasting system for the eThekweni Metro, Volume 1: Urgent Nowcasting using Radar-An Integrated Pilot Study. Water Research Commission (WCR). University of Natal. Silowa Printers South Africa. 2003 **Google Scholar**
5. Olajuyigbe AE, Rotowa OO, & Durojaye E. An assessment of flood hazard in Nigeria: The case of mile 12, Lagos. *Mediterranean Journal of Social Sciences*, 3(2), pp.367-375. Available from: <https://doi.org/10.5901/mjss.2012.v3n2.367> **Google Scholar**
6. Adeaga O. Adoption of Global Navigation Satellite System (GNSS) Technology in Flood Disaster Management and Response; A case study of Lagos Mega City: A paper Delivered at International Symposium on GNSS. 2008. **Google Scholar**
7. Theron M. *Climate Change and Increasing Floods in Africa: Implication for Africa's Development*. 2007 **Google Scholar**
8. Agbenyo PK. Effects of the Akosombo and Kpong power schemes on six selected Mafi-

communities in the Volta region. Diss. University of Cape Coast. 2009 **Google Scholar**

9. Mishio M. Improved management of the water and energy resources in the Volta Basin. MS thesis. 2021. NTNU. Available from: <https://ntnuopen.ntnu.no/ntnu-xmlui/handle/11250/2824233> **Google Scholar**
10. Gadzanku S. Evaluating electricity generation expansion planning in Ghana. Diss. Massachusetts Institute of Technology. MIT Institute for Data, Systems and Society. 2019 Available from: <https://globalchange.mit.edu/publication/17409> **Google Scholar**
11. Wemegah CS, Yamba EI, Aryee JN, Sam F, Amekudzi LK. Assessment of urban heat island warming in the greater Accra region. *Scientific African*. 2020;8:e00426. Available from: <https://doi.org/10.1016/j.sciaf.2020.e00426>. **Google Scholar**
12. Owusu K, & Waylen PR. The changing rainy season climatology of mid-Ghana. *Theoretical and Applied Climatology*. 112(3-4):419–430. 2013 Available from: <https://doi.org/10.1007/s00704-012-0736-5> **PubMed** | **Google Scholar**
13. Amekudzi LK, Yamba EI, Preko K, Asare EO, Aryee J, Baidu M, Codjoe SNA. Variabilities in rainfall onset, cessation and length of rainy season for the various agro-ecological zones of Ghana. *Climate*. 2015;3(2):416–434. Available from 3(2):416-434. Available from: <https://doi.org/10.3390/cli3020416> **Google Scholar**
14. Aryee J, Amekudzi L, Quansah E, Klutse N, Atiah W, Yorke C. Development of high spatial resolution rainfall data for Ghana. *International Journal of Climatology*. 2018;38(3):1201–121. Available from: <https://doi.org/10.1002/joc.5238> **Google Scholar**
15. Ansah SO, Ahiataku MA, Yorke CK, Otu-Larbi F, Yahaya B, Lamptey PNL, Tanu, M. Meteorological analysis of floods in Ghana. *Advances in Meteorology*, vol. 2020, Article ID 4230627, 14 pages. Available from: <https://doi.org/10.1155/2020/4230627> **Google Scholar**
16. Baidu M, Amekudzi LK, Aryee JN, Annor T. Assessment of long-term spatio-temporal rainfall variability over Ghana using wavelet analysis. *Climate*. 5(2):30 2017. Available from: <https://doi.org/10.3390/cli5020030> **Google Scholar**
17. Yamba EI, Aryee JNA, Quansah E, Davies P, Wemegah CS, Osei MA, et al. Revisiting the agro-climatic zones of Ghana: A re-classification in conformity with climate change and variability. *PLOS Clim* 2(1): e0000023. 2023 Available from: <https://doi.org/10.1371/journal.pclm.0000023> **Google Scholar**
18. Daily Graphic Online <https://www.graphic.com.gh/news/general-news/akosombo-dam-spillage-navy-rescues-8-000-flood-victims.html> Accessed on 30th October, 2023 at 17:20 GMT.
19. Volta River Authority, 2023: [https://www.vra.com/about\\_us/profile.php](https://www.vra.com/about_us/profile.php) Accessed on October 30, 2023 at 14:44 GMT

20. Adegoke J, Sylla M, & Bossa AY, Taylor C, & Klein C, Ogunjobi K, & Adoukpè, J, (2019). On the 2017 Rainy Season Intensity and Subsequent Flood Events Over West Africa. DOI: 10.33183/rccs.2019.p10. **Google Scholar**
21. Taylor A, & Davies H., 2019. An overview of climate change and urban development in cape town. Climate change and urban development: lessons from Cape Town. Cape Town: UCT Press.) **Google Scholar**
22. Duy PN, Chapman L, & Tight M. (2019). Resilient transport systems to reduce urban vulnerability to floods in emerging-coastal cities: A case study of Ho Chi Minh city Vietnam. *Travel Behaviour and Society*, 15, 28–43. <https://doi.org/10.1016/j.tbs.2018.11.001>. **Google Scholar**
23. Okuyama Y, Hewings GJ, & Sonis M. (2004). Measuring economic impacts of disasters: interregional input-output analysis using sequential interindustry model. In *Modeling spatial and economic impacts of disasters* (pp. 77-101). Berlin, Heidelberg: Springer Berlin Heidelberg. **Google Scholar**
24. Augusto c, & santos g. (2008). Runoff estimates into the weija reservoir and its implications for water. *Journal of urban and environmental engineering*, 2(2), 33–40. <https://doi.org/10.4090/juee.2008.v2n2.033040> **Google Scholar**
25. Owusu-Ansah, J. K., Dery, J. M., & Amoako, C. (2018). Flood vulnerability and coping mechanisms around the Weija Dam near Accra, Ghana. *GeoJournal*, 3. <https://doi.org/10.1007/s10708-018-9939-3> **Google Scholar**
26. Chan EYY, Tong KHY, Dubois C, McDonnell K, Kim JH, Hung KKC, & Kwok KO. Narrative Review of Primary Preventive Interventions against Water-Borne Diseases: Scientific Evidence of Health-EDRM in Contexts with Inadequate Safe Drinking Water. *International journal of environmental research and public health*, 18(23), 2021 12268. Available from: <https://doi.org/10.3390/ijerph182312268> **PubMed** | **Google Scholar**
27. Osunla CA, & Okoh AI. *Vibrio Pathogens: A Public Health Concern in Rural Water Resources in Sub-Saharan Africa*. *International journal of environmental research and public health*, 14(10), 2017 1188. Available from: <https://doi.org/10.3390/ijerph14101188> **PubMed** | **Google Scholar**
28. Fazal-ur-Rehman M. Polluted water borne diseases: Symptoms, causes, treatment and prevention. *J Med Chem Sci* 2.1: 2019 21-6. **Google Scholar**
29. Mirza MMQ. Climate change, flooding in South Asia and implications. *Reg Environ Change* 11 (Suppl 1), 95–107 (2011). <https://doi.org/10.1007/s10113-010-0184-7> **Google Scholar**
30. Dewan TH, Societal impacts and vulnerability to floods in Bangladesh and Nepal, *Weather and Climate Extremes*, Volume 7, 2015, Pages 36-42, <https://doi.org/10.1016/j.wace.2014.11.001> **Google Scholar**
- Fitzgerald WDG, Clark M, Hou X, Health impact of floods. *Prehosp. Disaster Med.*, 25 (2010), pp. 265-272 **Google Scholar**
31. Marques AH, Bjørke-Monsen AL, Teixeira AL, & Silverman MN. Maternal stress, nutrition and physical activity: Impact on immune function, CNS development and psychopathology. *Brain research*, 1617, 2015 28–46. Available from: <https://doi.org/10.1016/j.brainres.2014.10.051> **PubMed** | **Google Scholar**
32. World Health Organization. Vector-borne diseases. <https://www.who.int/news-room/fact->

sheets/detail/vector-borne-diseases. 2020 Available from: <https://www.who.int/news-room/factsheets/detail/vector-borne-diseases> Accessed 3 March, 2023.

33. Parham PE, Waldo J, Christophides GK, Hemming D, Agosto F, Evans KJ, et al. Climate, environmental and socio-economic change: weighing up the balance in vector-borne disease transmission. *Philosophical transactions of the Royal Society of London. Series B, Biological sciences*, 370(1665), 2015 20130551. Available from: <https://doi.org/10.1098/rstb.2013.0551> **PubMed** | **Google Scholar**
34. Shaman J, Yang W, & Kandula S. Inundation, Mosquito Dynamics, and the Spread of West Nile Virus. *PLOS Neglected Tropical Diseases*, 13(5), e0007320. 2019 **Google Scholar**
35. Yasuoka J, & Levins R. Impact of deforestation and agricultural development on anopheline ecology and malaria epidemiology. *The American Journal of Tropical Medicine and Hygiene*, 76(3), 2007 pp 450-460. Available from: <https://pubmed.ncbi.nlm.nih.gov/17360867/> **PubMed** | **Google Scholar**
36. Patz JA, Graczyk TK, Geller N, & Vittor AY. Effects of environmental change on emerging parasitic diseases. *International journal for parasitology*, 30(12-13), 2000 1395–1405. Available from: [https://doi.org/10.1016/s0020-7519\(00\)00141-7](https://doi.org/10.1016/s0020-7519(00)00141-7) **PubMed** | **Google Scholar**
37. Coalson JE, Anderson EJ, Santos EM, Madera GV, Romine JK, Dominguez B, Richard DM, Little AC, Hayden MH, & Ernst KC. The Complex Epidemiological Relationship between Flooding Events and Human Outbreaks of Mosquito-Borne Diseases: A Scoping Review. *Environmental health perspectives*, 129(9), 2021 96002. Available from: <https://doi.org/10.1289/EHP8887> **PubMed** | **Google Scholar**
38. Gething, P. W., Smith, D. L., Patil, A. P., Tatem, A. J., Snow, R. W., & Hay, S. I. (2010) Climate change and the global malaria recession. *Nature*, 465(7296), 342–345. Available from: <https://doi.org/10.1038/nature09098> **PubMed** | **Google Scholar**
39. Nigusie A, Gizaw Z, Gebrehiwot M, & Destaw B. Vector-Borne Diseases and Associated Factors in the Rural Communities of Northwest Ethiopia: A Community-Based Cross-Sectional Study. *Environmental health insights*, 15, 2021 11786302211043049. Available from: <https://doi.org/10.1177/11786302211043049> **PubMed** | **Google Scholar**
40. Akanda AS, Johnson K, Ginsberg HS, Couret J, Prioritizing Water Security in the Management of Vector-Borne Diseases: Lessons From Oaxaca, Mexico. *Geohealth*. 2020 Mar 1;4(3):e2019GH000201. Available from: <https://doi.org/10.1029/2019GH000201>. **PubMed** | **Google Scholar**
41. World Health Organization. Dengue: Guidelines for diagnosis, treatment, prevention, and control. World Health Organization. 2014 Available from: <https://www.who.int/publications-detail-redirect/9789241547871> Accessed on March 29, 2024
42. Akanda AS, Johnson K, Ginsberg HS, Couret J, Prioritizing Water Security in the

Management of Vector-Borne Diseases: Lessons From Oaxaca, Mexico. *Geohealth*. 2020 Mar 1;4(3):e2019GH000201. Available from: [https://doi: 10.1029/2019GH000201](https://doi.org/10.1029/2019GH000201). **PubMed| Google Scholar**

Ehimen FA, Akpan IS, Osagiede EF, Ofili AN, Okukpon PO, Airefetalor IA. Assessment of Standard Precautions' Practices among Health Care Workers in a Rural Area of South-South Nigeria. *Curr. J. Appl. Sci. Technol.* [Internet]. 2020 Jun. 16 [cited 2024 May 24];39(15):8-21. Available from: <https://journalcjast.com/index.php/CJAST/article/view/3413>

Cherif MKC-, Sakyi DA-, Maiga B, Diop S, Dolo A, Nébié I, -Blomberg MT. Patients' Perception of Health Care Services in Malaria Endemic Area in Mali: Villages with Health Care Services Versus villages Without Health Care Services. *Asian J. Med. Health.* [Internet]. 2021 Jan. 2 [cited 2024 May 24];18(11):91-9. Available from: <https://journalajmah.com/index.php/AJMAH/article/view/501>

Mulderij-Jansen V, Gerstenbluth I, Duits A, Tami A, Bailey A. Evaluating and strengthening the health system of Curaçao to improve its performance for future outbreaks of vector-borne diseases. *Parasites & Vectors*. 2021 Dec;14:1-7.