

## Original Research Article

### Response of different levels of PGR on gerbera (*Gerbera jamesonii*) cv. Pink elegance under protected cultivation

#### Abstract

An experiment was conducted to evaluate the influence of plant growth regulators on the growth, flower quality and yield of gerbera (*Gerbera jamesonii* L.) cv. Pink Elegance under a naturally ventilated polyhouse in the Department of Horticulture at SHUATS, Allahabad, U.P., during 2023-24. The experiment followed a Randomized Block Design with three replications and ten treatments. Gerbera plants were sprayed with GA<sub>3</sub>, Cycocel and NAA at different concentrations of 50, 100 and 150 ppm along with control. The results from the study concluded that the treatment GA<sub>3</sub> @ 150 ppm was found in superior among other treatments in terms of plant height (33.30 cm), number of leaves per plant (30.96), plant spread (49.68), leaf area (152.34m<sup>2</sup>), leaf area index (3.77), number of flowers per plant (12.00), flower weight (40.12g), flower diameter (12.31cm), root length (26.80cm), stalk length (47.88cm), shelf life (9.95), flower yield per m<sup>2</sup> (112.89), flower yield per 1000 polybags (12000) and minimum days to first flowering (69.33). Thus, the 150 ppm GA<sub>3</sub> treatment was identified as the most effective for promoting both excellent vegetative and reproductive growth. Thus, GA<sub>3</sub> @ 150 ppm can be recommended for plant growth and flowering of Gerbera proving the multifaceted role of PGR in gerbera cultivation, helping growers optimize plant growth, flowering & yield while maintaining high-quality flower production.

**Keywords:** cCycocel, GA<sub>3</sub>, gGerbera, growth, NAA, yield

#### Introduction

Gerbera (*Gerbera jamesonii* L.), a prominent cut flower, ranks fifth among the top ten cut flowers in the global market and is extensively cultivated under protected conditions in regions like Bangalore, Pune, and Uttarakhand. Native to tropical Asia and Africa, this species belongs to the Asteraceae family and known by various names such as Transvaal Daisy, Barberton Daisy, or African Daisy and can be grown under a wide range of climatic conditions (Swarup, 1997). Discovered by Gronovious and named in honor of German naturalist Traugott Gerber, gerbera is valued for its wide range of colors, including yellow, red, orange, cream, white, and various shades in between (Prasad, 2007). These features, along with its use in floral arrangements and decorations, make it a popular choice in India.

Gerbera is a dwarf herbaceous perennial, characterized by solitary flower heads on long slender stalks, which thrive in polyhouses and shade houses (Sahu *et al.*, 2016). These environments enhance plant growth, producing larger, greener leaves with high dry matter content. The commercial production of gerbera is in high demand both nationally and internationally, especially during winter when few other flowers bloom (Darras, 2021).

Gerbera flowers are in high demand nationwide due to their enduring nature, elegant appearance, resilience, and export potential. Optimizing the application of PGRs can lead to increased productivity and improved quality of gerbera flowers, thus enhancing the economic viability of gerbera production. Understanding the specific effects of different PGRs on gerbera growth, flowering, and fruiting can enable growers to tailor their application methods to maximize desired outcomes while minimizing costs and environmental impacts (Fawzy *et al.*, 2012). Moreover, research in this area can contribute to the development of sustainable

**Commented [mm1]:** On abstract: please add introduction, prolems, aims, methods, results

cultivation practices by reducing the reliance on conventional chemical inputs and promoting environmentally friendly alternatives and gerbera plants can inform breeding efforts aimed at developing new cultivars with enhanced responsiveness to PGRs or intrinsic resistance to environmental stresses.

Plant growth regulators (PGRs) have revolutionized the floriculture industry by significantly enhancing the growth, yield, and quality of ornamental crops like gerbera. PGRs, including gibberellic acid (GA<sub>3</sub>), cycocel and naphthalene acetic acid (NAA), are chemical substances used in small amounts to regulate plant development and stimulate desired growth responses (Wu *et al.*, 2024). They help overcome growth limitations, stimulate flowering, and extend the vase life of flowers (Thakur *et al.*, 2023). Given the significance of PGRs, the present study aimed to determine the optimal concentrations of GA<sub>3</sub> and NAA for improving the growth, yield, and quality of gerbera under naturally ventilated polyhouses. This research is conducted to study the effect of different levels of PGR on growth, flowering & yield of Gerbera.

### Materials and methods

The experiment was conducted during the winter season of 2023-24 at the Department of Horticulture, Sam Higginbottom University of Agriculture, Technology and Sciences, Allahabad, UP. The experimental site is located at a latitude of 20°15' North and a longitude of 60° East, at an altitude of 98 meters above mean sea level (MSL). The soil at the experimental site was sandy loam in texture with a pH of 6.87, an electrical conductivity (E.C) of 0.15 dS/m, and a carbon content of 43%. The available nitrogen, phosphorus, and potassium contents were 216.62 kg/ha, 37.34 kg/ha, and 215.86 kg/ha, respectively. The experiment was laid out in a Randomized Block Design with three replications and ten treatments. Seedlings of cv. Pink Elegance, each with 4-5 leaves, were planted on raised beds on November 13, 2023 in polybags. Three plant growth regulators, gibberellic acid (GA<sub>3</sub>), cycocel and naphthalene acetic acid (NAA), were used at three concentrations 50, 100 and 150ppm along with the control. The details of the treatment combinations are as follows: T<sub>0</sub>- Control, T<sub>1</sub>- GA<sub>3</sub> 50 ppm, T<sub>2</sub>- GA<sub>3</sub> 100 ppm, T<sub>3</sub>- GA<sub>3</sub> 150 ppm, T<sub>4</sub>- Cycocel 100 ppm, T<sub>5</sub>- Cycocel 100 ppm, T<sub>6</sub>- Cycocel 150 ppm, T<sub>7</sub>- NAA 50 ppm, T<sub>8</sub>- NAA 100 ppm and T<sub>9</sub>- NAA 150 ppm.

**Commented [mm2]:** On methods: add references on all parameter

### Results and Discussion

#### Influence of plant growth regulators on Growth parameters

The data on growth parameters presented in Table 1 show that the maximum plant height (35.68 cm), number of leaves (30.96), plant spread (51.21 cm), leaf area (178.64m<sup>2</sup>) and leaf area index (3.77) were observed in treatment T<sub>3</sub> (GA<sub>3</sub> @ 150 ppm). In contrast, the minimum plant height (21.12 cm), number of leaves (30.96), plant spread (38.02 cm), leaf area (129.65 m<sup>2</sup>) and leaf area index (1.71) were recorded in treatment T<sub>0</sub> (control). The significantly greater plant height, number of leaves, plant spread, leaf area and leaf area index in T<sub>3</sub> can be attributed to the effect of gibberellic acid, which promotes cell division and cell elongation, leading to enhanced vegetative growth. Similar findings regarding maximum plant height were reported by Nair *et al.* (2002) and Dalal *et al.* (2009) in gerbera. The maximum plant spread at 150 ppm was observed by Patra *et al.* (2015) in gerbera. Sharma *et al.* (2004) suggested that GA<sub>3</sub> at 100 ppm is highly effective in promoting vegetative growth in gladiolus and Dogra *et al.* (2012) reported an increase in leaf area index due to GA<sub>3</sub> treatments. Similarly, Porwal *et al.* (2002) observed the impact of plant growth regulators on Damask Rose, noting that GA<sub>3</sub> at 250 ppm resulted in the maximum plant height, number of shoots per plant, and plant spread. Additionally, GA<sub>3</sub> promotes the increase in the photosynthetic area by inducing

cell elongation and hyponasty of leaves, which affects their orientation and ultimately contributes to an increase in leaf area (Dutta, 2015).

#### **Influence of plant growth regulators on flowering parameters**

The data on qualitative parameters presented in Table 2 indicate that the minimum number of days to first flower bud emergence or earliness, which occurred in 69.33 days, was observed in T<sub>3</sub> (GA<sub>3</sub> @ 150 ppm), while T<sub>0</sub> (control) took the maximum number of days (93.11 days) to produce a visible flower bud. Maximum number of flowers per plant (12.00), flower weight (40.12), flower diameter (12.31), root length (26.80), stalk length (47.88), and shelf life (9.95) was observed with T<sub>3</sub> - GA<sub>3</sub> @ 150 ppm. Conversely, the minimum number of flowers per plant (4.67), flower weight (26.72), flower diameter (7.24), root length (14.86), stalk length (35.55), and shelf life (5.63) were found in treatment T<sub>0</sub> (control).

Gibberellic acid is a plant growth regulator known to play a crucial role in various physiological processes, including cell division, elongation, and differentiation. When applied exogenously, GA<sub>3</sub> can stimulate these processes, leading to significant changes in plant morphology and development (Ridha *et al.*, 2016). In the case of gerbera plants, the application of GA<sub>3</sub> at 150 ppm stimulated the cell expansion and enlargement of floral structures which have accelerated the initiation and emergence of flower buds, resulting in earlier flowering, promoted flower bud formation, enhanced floral organ development, increased flower weight and diameter (Saini and Arora, 2016). Moreover, GA<sub>3</sub> (150 ppm) treated plants displayed enhanced root and stalk growth, with longer root systems and taller stalks, which can support larger and more abundant flowers. It also improved the post-harvest characteristics such as shelf life, as GA<sub>3</sub> at 150 ppm treated flowers exhibited delayed senescence and prolonged freshness compared to untreated flowers (Zosser *et al.*, 2017). Overall, the application of GA<sub>3</sub> at 150 ppm effectively optimized qualitative parameters in gerbera plants, resulting in enhanced flower quality and overall plant performance.

Patel *et al.* (2013) observed similar results in gerbera, with GA<sub>3</sub> at 150 ppm resulting in the fewest days to first flower initiation (108.33 days), maximum flower diameter (8.76 cm), flower weight (5.93 g), and shelf life of flowers (8.00 days). GA<sub>3</sub> also increased the number of suckers, and foliar application of 150 ppm GA<sub>3</sub> was found to be the best for enhancing plant growth, yielding the maximum number of cut blooms with longer stalks and larger flower sizes, as reported by Sharifuzzaman *et al.* (2011) in chrysanthemum and Jamal *et al.* (2014) in gerbera and Sainath *et al.* (2018) in chrysanthemum observed similar earliness in the full opening of flowers due to GA<sub>3</sub> treatments. Chauhan *et al.* (2014) found that GA<sub>3</sub> at 150 ppm led to the earliest appearance of the first flower bud (50.98 days) and the largest flower diameter (11.37 cm) in gerbera. Furthermore, Salem *et al.* (2016) reported that in gerbera, the highest yield and quality parameters (number of flowers per plant, stalk length, flower bud diameter, and stalk diameter), as well as the shortest time to flower bud emergence and first flowering, were observed with GA<sub>3</sub> at 150 ppm.

#### **Influence of plant growth regulators on Yield parameters**

The data on yield parameters presented in Table 3 indicate that the maximum flower yield per m<sup>2</sup> (112.89) and maximum flower yield per 1000 polybags (12000) was observed with T<sub>3</sub> - GA<sub>3</sub> @ 150 ppm. However, the minimum flower yield per m<sup>2</sup> (65.15) and flower yield per 1000 polybags (4670) were recorded in treatment T<sub>0</sub> (control). The increase in flower yield could be attributed to the fact that plants treated with gibberellic acid produced a greater number of leaves and more number of flowers. This increased leaf production likely led to the production and accumulation of more photosynthates, which were then redirected to the floral

organs, resulting in a higher number of flowers and flower weight. These findings are consistent with the results reported by Dalal *et al.* (2009) in gerbera, Tariq *et al.* (2011) in gladiolus, Sainath *et al.* (2018) in chrysanthemum and Patel *et al.* (2010) in chrysanthemum, Kumar *et al.* (2014) and Patel *et al.* (2015) in African marigold. .

## Conclusion

From the present investigation, it is concluded that among the different treatments, the treatment T<sub>3</sub> - GA<sub>3</sub> @ 150 ppm was found in superior among other treatments in terms of Plant height (33.30 cm), Number of leaves per plant (30.96), plant spread (49.68), leaf area (152.34m<sup>2</sup>), leaf area index (3.77), number of flowers per plant (12.00), flower weight (40.12g), flower diameter (12.31cm), root length (26.80cm), stalk length (47.88cm), shelf life (9.95), flower yield per 1000 polybags (12000) and minimum days to first flowering (69.33). Thus, GA<sub>3</sub> @ 150 ppm can be recommended for plant growth and flowering of Gerbera proving the multifaceted role of PGR in gerbera cultivation, helping growers optimize plant growth, flowering & yield while maintaining high-quality flower production

## References

- Chauhan, R.V., Kav, K.P., Babariya, V.J., Pansuria, P.B. and Savaliya, A.B. 2014. Effect of gibberellic acid on flowering and cut flower yield in Gerbera under protected condition. *Asian Journal of Horticulture*, 9(2):404-407.
- Dalal, S.R., Somavanshi, A.V. and Karale, G.D. (2009) Effect of gibberellic acid on growth, flowering, yield and quality of Gerbera under polyhouse condition; *International journal of agriculture sciences*, 5(2): 355-356.
- Darras, A. (2021). Overview of the dynamic role of specialty cut flowers in the international cut flower market. *Horticulturae*, 7(3), 51.
- Dogra, S., Pandey, R. K., and Bhat, D. J., 2012. Influence of gibberellic acid and plant geometry on growth, flowering production in gerbera under Jammu agroclimate, *International Journal of Pharma and Bio Sciences*, 3(4): 1083-1090.
- Dutta, S.C. (2015). *Plant Physiology*. Wiley Eastern Limited. New Age International Limited, pp. 466-558.
- Emongor, V.E. (2004) Effect of gibberellic acid on post-harvest quality and vase life of Gerbera cut flowers (*Gerbera jamesonii*), *Journal of agronomy*, 3(3): 191-195.
- Fawzy, Z. F, El-Shal, Z. S, Yunsheng L., Zhu O., Sawan O. M. 2012. Response of gerbera plants to foliar spraying of some bio- stimulants under sandy soil condition. *Applied Scientific Research*. 8 (2): 770-776.
- Girisha, R., Shirol, A. M., Reddy, B. S., Kulkarni, B. S., Patil, S. and Murthy, G. H. K. 2012. Growth, quality and yield characteristics of daisy (*Aster amellus*) cultivar Dwarf Pink as influenced by different plant growth regulator; *Karnataka J. Agric. Sci.*, 25(1), 163-165.
- Jamal Uddin AFM, Mehraj H, Taufique T, Ona AF and Parvin S. 2014. Foliar Application of Gibberellic Acid on Growth and Flowering of Gerbera Cultivars, *Journal of Bioscience and Agriculture Research*, 02(01): 52-58.
- Jamil, M. K., Rahman, M. M., Hosssain, M. M., Hosssain, M. T. and Karim, A. J. M. S. (2015) Effect of plant growth regulators on flowering and bulb production of Hippeastrum (*Hippeastrum hybridum*). *Bangladesh J. Agril.Res.* 40(4): 591-600.

Commented [mm3]: Add references using limited 5 years ago

- Kumar, M., Singh, A. K. and Kumar A. 2014. Effect of plant growth regulators on flowering and yield attributes of African marigold (*Tagetes erecta* L.) cv. Pusanarangigainda, *Plant Archives*, 14(1): 363-365.
- Kumar, R., Ahmed, N., Singh, D. B., Sharma, O. C., Lal, S., and Salmani, M. M. 2013. Enhancing blooming period and propagation coefficient of Tulip (*Tulipa gesneriana* L.) using growth regulators. *African Journal of Biotechnology*, 12(2):168-174.
- Kumar, M., Singh, A. K., Kumar, A. 2014. Effect of plant growth regulators on flowering and yield attributes of African marigold (*Tagetes erecta* L.) cv. Pusa Narangi Gainda. *Plant Archives*, 14(1):363-365.
- Nair, S. A., Singh, V., & Sharma, T. V. R. S. 2002. Effect of plant growth regulators on yield and quality of gerbera under Bay Island conditions. *Indian Journal of Horticulture*, 59(1), 100-105.
- Padaganur, V. G., Monakshi, A. N., and Patel, V. S. 2005. Effect of plant growth regulators on growth and yield of Tuberose cv. Single, *Karnataka J.Agric.Sci.*, 18(2):469-473.
- Palei, S., Das, A. K. and Dash, D. K. 2016. Effect of plant growth regulators on flowering and yield attributes of African marigold (*Tagetes erecta* L.), *International education and research journal*, 2(6):2454-9916.
- Patel, B. B., Desai, J. R., Patel, G. D., and Patel, H. F. 2013. Influence of foliar application of nitrogen and plant growth regulators on growth, flowering production of gerbera, *Bioinfolet. Journal*, 10 (2A): 415 – 417.
- Patil, D., Chopde, N., Lokhande, S., and Bhande, M. H. 2016. Studies on response of African marigold to plant growth regulators for seed production. *Plant Archives*, 16 (1): 423425.
- Patra, S. K., Beura, S., and Shasani, T. 2015. Efficacy of GA<sub>3</sub> on growth and flowering regulation of in vitro raised hybrid Gerbera under shade net. *Agric. Sci. Digest.*, 35 (3):173-177.
- Porwal, R., C.L. Nagda and J.P.S. Pundir. 2002. Influence of plant growth regulator on vegetative growth and flower earliness of Damask rose. *South Indian Hort.*, 50 (1-3): 119 - 123.
- Prasad, K. V. 2007. Bulbous ornamental crops: Status of global trade. *Floriculture Today*. 11(8), 22-36.
- Ridha A. A. S., S. Saravanan and Prasad V. M. 2016. Effect of Gibberellic Acid Spraying on yield and Flowers of Gerbera (*Gerbera jamesonii*) Cv. Dennis. *International J. of Scientific Res.* 5(3), 234-236.
- Sahu, M. K., Tirkey, T., Sharma, G., Tiwari, A., & Kushram, T. (2016). Effect of foliar application of micronutrients on growth and flower production of gerbera under protected condition. *Journal of Pure and Applied Microbiology*, 10(4), 3099-3103.
- Sainath, D. S., Uppar, V. S., Patil, V. K., Deshpande and Husije, R., 2018. Effect of different growth regulators on seed yield and quality attributes in annual chrysanthemum (*Chrysanthemum coronarium* L). *Journal of Agricultural Sciences*, 27(2): 131-134.
- Saini S and Arora S. 2016. Effect of gibberellic acid on growth, flowering, yield and quality of gerbera under polyhouse conditions, *Indian Nutrient Digest*, 11: 233-236.

- Sajid, M., Amin, N., Ahmad, H., and Khan, K. 2016. Effect of gibberellic acid on enhancing flowering time in *Chrysanthemum morifolium*; *Pak.J. Bot.*, 48(2): 477-483.
- Salem, R. A. A., Saravanan, S., and Prasad V. M. 2016. Effect of gibberellic acid spraying on yield and flowers of Gerbera (*Gerbera jamesonii*) cv. Dennis. *International journal of scientific research*, 5 (3):2277-2280.
- Sharifuzzaman, S. M., Ara, K. A., Rahman, M. H., Kabir, K. and Talukdar, M. B., 2011. Effect of GA<sub>3</sub>, CCC and MH on vegetative growth, flower yield and quality of *Chrysanthemum*. *Int. J. Expt. Agric.* 2(1):17-20.
- Sharma, J.R., R.B. Gupta and R.D. Panwar. 2004. Growth, flowering and corm production of gladiolus cv. Friendship as influenced by foliar application of nutrients and growth regulators. *Journal of Ornamental Horticulture*. 7 (3-4): 154 – 158.
- Sujatha, A. N., Singh, V. and Sharma, T. V. R. S. 2002 Effect of plant growth regulators on yield and quality of Gerbera under Bay island condition. *Indian J. Hort.*, 59(1):100-105.
- Suvalaxmi Patel, A. K. Das and D. K. Dash. 2015. Effect of Plant Growth Regulators on Growth and Yield Attributes of African Marigold (*Tagetes erecta* L.) *International Education & Res. J.* 2(6):44-45.
- Swarup, V. 1997. Garden Flowers, *National Book Trust*, pp.184-185.
- Tariq S., Hassan, I., Akhtar, N., Abbasi, Jilani. G. 2013. Effect of gibberellic acid on the vase life and oxidative activities in senescing cut gladiolus flowers. *Article in Plant Growth Regulation*. 72:89-95.
- Thakur, R., Thakur, R., Chandermohan, N. C., & Kanwar, B. (2023). Effect of Gibberellic acid and benzyl adenine on ornamental plants, *The Pharma Innovation Journal*, 12(2): 101-105.
- Uddin, A. F. M. J., Mehraj, H., Taufique, T., Ona, A. F. and Parvin, S. 2014. Foliar Application of gibberellic acid on growth and flowering of Gerbera cultivars. *Journal of Bioscience and Agriculture Research*, 2(1): 52-58.
- Uddin, M. S., Hossain, A. V. M. S., Normania, O., Boyce, A. N. and Moneruzzaman, K. M. 2009. Effect of Naphthalene acetic acid and Gibberellic acid in prolonging bract longevity and delaying discoloration of *Bougainvillea spectabilis*, *J. of Biotechnology*, 8 (3): 3434-350.
- Wu, X., Gong, D., Zhao, K., Chen, D., Dong, Y., Gao, Y. & Hao, G. F. (2024). Research and development trends in plant growth regulators. *Advanced Agrochem*, 3(1), 99-106.
- Zosser, C. N., Sangma, Devi, S. and Urfi, F. 2017. Effect of plant growth regulators on growth, yield and flower quality of Gerbera (*Gerbera jamesonii* L.) cv. Pink Elegance under naturally ventilated polyhouse (NVPH), *International Journal of Current Microbiology and Applied Sciences*. 6(10): 468-476.

**Table 1. Effect of Plant growth regulators on growth parameters of gerbera**

Tr.	Plant height (cm)	Number of leaves/plant	Plant spread (cm)	Leaf Area (m <sup>2</sup> )	Leaf Area Index (cm <sup>2</sup> )
T <sub>0</sub>	21.12	18.16	38.02	129.65	1.71

T <sub>1</sub>	31.66	24.86	47.02	146.40	2.78
T <sub>2</sub>	33.30	28.79	49.68	171.19	3.19
T <sub>3</sub>	35.68	30.96	51.21	178.64	3.77
T <sub>4</sub>	27.43	19.25	39.72	124.06	2.04
T <sub>5</sub>	29.77	21.67	41.35	152.34	2.46
T <sub>6</sub>	29.33	23.56	44.85	164.18	2.53
T <sub>7</sub>	24.04	22.24	42.02	164.48	2.36
T <sub>8</sub>	25.49	25.91	43.35	153.13	2.94
T <sub>9</sub>	27.04	26.73	45.35	146.58	3.12
<b>F-Test</b>	<b>S</b>	<b>S</b>	<b>S</b>	<b>S</b>	<b>S</b>
<b>S.Ed (+)</b>	<b>0.94</b>	<b>1.06</b>	<b>0.87</b>	<b>2.69</b>	<b>0.06</b>
<b>CD<sub>(0.05)</sub></b>	<b>2.00</b>	<b>2.24</b>	<b>1.84</b>	<b>5.70</b>	<b>0.13</b>

**Table 2. Effect of Plant growth regulators on floral parameters of gerbera**

Tr.	Days to first flowering	Flower Weight (g)	Flower Diameter (cm)	Stalk Length (cm)	Shelf life (Days)	Root length (cm)
T <sub>0</sub>	93.11	26.72	7.24	35.55	5.63	14.86
T <sub>1</sub>	72.48	36.99	10.65	44.35	7.67	23.66
T <sub>2</sub>	70.27	38.64	11.72	45.93	8.69	24.85
T <sub>3</sub>	69.33	40.12	12.31	47.88	9.95	26.80
T <sub>4</sub>	88.01	29.39	7.83	36.22	6.63	17.42
T <sub>5</sub>	84.88	31.03	8.09	38.47	6.61	18.19
T <sub>6</sub>	80.29	32.24	8.75	38.69	6.34	18.36
T <sub>7</sub>	77.73	34.72	9.98	39.33	7.52	19.44
T <sub>8</sub>	75.26	34.81	10.26	40.62	8.31	22.21
T <sub>9</sub>	78.33	35.36	10.31	42.31	8.66	22.30
<b>F-Test</b>	<b>S</b>	<b>S</b>	<b>S</b>	<b>S</b>	<b>S</b>	<b>S</b>
<b>S.Ed (+)</b>	<b>1.78</b>	<b>0.64</b>	<b>0.21</b>	<b>1.13</b>	<b>0.44</b>	<b>0.52</b>
<b>CD<sub>(0.05)</sub></b>	<b>3.77</b>	<b>1.35</b>	<b>0.45</b>	<b>2.40</b>	<b>0.94</b>	<b>1.11</b>

**Table 3. Effect of Plant growth regulators on yield parameters of gerbera**

Treatments	Number Of Flowers Per Plant	Flower Yield per m <sup>2</sup>	Flower Yield per 1000 Polybags
T <sub>10</sub>	4.67	65.15	4670
T <sub>1</sub>	8.23	99.39	8230
T <sub>2</sub>	10.00	107.58	10000
T <sub>3</sub>	12.00	112.89	12000
T <sub>4</sub>	6.86	77.58	6860
T <sub>5</sub>	7.00	76.11	7000
T <sub>6</sub>	7.57	90.23	7570
T <sub>7</sub>	7.61	82.05	7610
T <sub>8</sub>	8.12	86.54	8120

<b>T<sub>9</sub></b>	9.20	92.15	9200
<b>F-Test</b>	<b>S</b>	<b>S</b>	<b>S</b>
<b>S.Ed (+)</b>	<b>0.70</b>	<b>1.55</b>	<b>180.414</b>
<b>CD (5%)</b>	<b>1.48</b>	<b>3.28</b>	<b>381.972</b>

UNDER PEER REVIEW