

Growth and yield of summer sesamum (*Sesamum indicum* L.) as influenced by varying organic nutrient management practices

ABSTRACT

A field experiment was conducted during *Zaid* season of 2023 at Agricultural Research Station, Binjhagiri, Chhatabar, Faculty of Agricultural Sciences, Siksha 'O' Anusandhan (Deemed to be University), Bhubaneswar, Odisha. The experiment was laid out in Randomized Block Design with ten treatments each replicated thrice. Soil of the experiment field was sandy loam in texture, atslightly acidic(6.40), low in organic carbon (0.45%) and low in available nitrogen (179 kg ha⁻¹), medium in available phosphorous (20.0) and potassium (142). A keen observation of data revealed that application of various organic sources like FYM, jeevamrutha, beejamrutha, panchagyavya, vermicompost and rock phosphate significantly influenced the growth parameters, yield parameters, yield and economic attributes like net return and return per rupee invested.

Keywords: FYM, vermicompost, panchagavya, jeevamrutha, rock phosphate, sesame

1.INTRODUCTION

Sesame is one of the most important oilseed crop in India, grown next to groundnut, rapeseed and mustard. It is cultivated in an area of 19.53 lakh hectares in India with an annual production of 7.84 lakh tonnes and productivity of 463 kg/ha. It is mostly cultivated in Maharashtra, Uttar Pradesh, Rajasthan, Odisha, Andhra Pradesh, Madhya Pradesh, Tamil Nadu and Karnataka. These six states account about 64 percent of an area and 78 percent of the production of sesame in the country [1]. India ranks first in area, production and export of sesame in the world. Sesame ranks third in terms of total oilseed area and fourth in terms of total oilseed production in India. Lower productivity of sesame is due to use of sub-optimal rate of fertilizer, poor management and cultivation of sesame in marginal and sub-marginal lands where deficiency of macronutrients such as nitrogen, phosphorus, potassium and micronutrients is predominant. Increased usages of chemical fertilizers without adequate organic recycling has not only aggravated multi-nutrient deficiencies in soil-plant system, but also deteriorated soil health and created environmental pollution. Therefore, it is right time to evaluate the feasibility and efficiency of organic agriculture not only for improving building up soil fertility, but also for increasing the efficiency of chemical fertilizers. Now-a-days there is a huge demand for organic sesame in the global market. India has greater scope to produce sesame as it is traditionally grown without much chemical fertilizer and plant protection [2].

Organic agriculture is a production system, which avoids or largely excludes the use of chemical fertilizers, pesticides and growth regulators. To the extent feasible, organic farming relies on crop rotations, crop residues, animal manures, green manures, off farm organic wastes and aspects of biological pest management to maintain soil productivity and to manage insects and pests. The

Commented [M1]: at the Agricultural

Commented [M2]: and each treatment replicated thrice

Commented [M3]: slightly acidic (6.4),

Commented [M4]: (0.45 %), low in available nitrogen (179 kg ha⁻¹),

Commented [M5]: (20.0 ...?)

Commented [M6]: (142...?)

Commented [M7]: The results showed that the application

Commented [M8]: Significantly affected

Commented [M9]: Attributes of summer sesamum

Commented [M10]: Return per rupee.

Commented [M11]: Summer sesame, organic nutrients, rock phosphate, seed yield

Commented [M12]: crops

Commented [M13]: that grown

Commented [M14]: [Reference number]

Commented [M15]: summer sesame

Commented [M16]: summer sesamum

Commented [M17]: soil and plant system

Commented [M18]: [Reference number]

concept of soil as a living system that promotes the activities of beneficial organisms is central to this definition. For successful organic farming in any crops depends upon the availability of nutrients at right time and in right amount, particularly in short duration crops [1].

Commented [M19]: [Reference number]

Organic manures like FYM add much needed organic and mineral matter to the soil. The organic matter added is an indispensable component of soil and plays an important role in maintenance and improvement of soil fertility and productivity. Farm yard manure is known to play an important role in improving the fertility and productivity of soils through its positive effects on soil physical, chemical and biological properties and balanced plant nutrition. Vermicompost is considered as a high nutrient biofertilizer with diverse microbial communities, it plays a major role in improving growth and yield of different field crops, vegetables, flower and fruit crops [2]. Panchagavya is a mixed culture of naturally occurring beneficial microbes mostly lactic and bacteria (Lactobacillus), yeast (Saccharomyces), actinomyces (Streptomyces), photosynthetic bacteria (Rhodospirillum rubrum) and certain fungi (Aspergillus) which promotes the growth and yield of different crops. Jeevamruth is one of those organic fertilizer which have large number of nutrients like nitrogen, phosphorus, calcium and other micronutrients. This will ensure higher yield by enhancing the availability of nutrients through faster decomposition of bulky organic manures by boosting the microbial activity in the soil [3]. Use of beejamrutha, a mix of cow dung, cow urine, water, lime and a handful of soil has been given importance in sustainable agriculture since age old days. It is also one such organic product helpful for the plant growth. The beneficial microorganisms present in beejamrutha are known to protect the crop from harmful soil-borne and seed-borne pathogens. In this context, it is worth noting that nutrient management through organics play a major role in maintaining soil health due to build up of soil organic matter, beneficial microbes and enzymes, besides improving soil physical and chemical [4].

Commented [M20]: [Reference number]

Commented [M21]: [Reference number]

Commented [M22]: [Reference number]

Thus, the research aims to evaluate the effect of different organic manuring schedule on growth and yield of sesame, to determine the nutrient uptake under different sets of treatment, and to work out the economics for different organic manuring practices.

Commented [M23]: The objective of present study

2. MATERIALS AND METHODS

Commented [M24]: After dot, a space should be considered, like this 2. MATERIALS AND METHODS

2.1 Experimental Area and Treatments

Commented [M25]: Must be removed

The experiment was carried out at Agricultural Research Station, Binjhagiri, Chhatarbar (20°26 North and 85° 67 East, 45 meters above mean sea level) in the Faculty of Agricultural Sciences, Siksha 'O' Anusandhan (Deemed to be University), Bhubaneswar, Odisha. The experiment was conducted on sandy loam soil was slightly acidic (6.40), low in organic carbon (0.45%) and low in available nitrogen (179 kg ha⁻¹), medium in available phosphorous (20.0) and potassium (142). The experiment was laid out in a Randomized Block Design with ten treatments and each treatment was replicated three times. The gross plot size was 5m x 3m and net plot size was 4m x 2.6m. The treatment combinations are T₁- Control, T₂ - 100 % RDN

Commented [M26]: What is unit?

Commented [M27]: What is unit?

Commented [M28]: What is unit?

through 50% of FYM and 50% of vermicompost + remaining RDP through rock phosphate, T₃- 50 % RDN through 50% of FYM and 50 % of vermicompost + remaining RDP through rock phosphate, T₄- Seed treatment with beejamrutha 10 % solution @ 20 ml kg⁻¹ seed + application of panchagavya @ 3 % as one soil application and two foliar application (branching and flowering), T₅- Seed treatment with beejamrutha 10 % solution @ 20 ml kg⁻¹ seed + application of jeevamrutha @ 10 % as one soil application and two foliar application (branching and flowering), T₆- Seed treatment with beejamrutha 10 % solution @ 20 ml kg⁻¹ seed + application of panchagavya @ 3 % and jeevamrutha @ 10 % as soil application, T₇- Seed treatment with beejamrutha 10 % solution @ 20 ml kg⁻¹ seed + application of panchagavya @ 3 % and jeevamrutha @ 10 % as one soil application and two spray of panchagavya @ 3 % and jeevamrutha @ 10 % as foliar application (branching and flowering), T₈-T₃+T₄, T₉ -T₃+T₅ , T₁₀- T₃+T₄+T₅.

3.RESULTS AND DISCUSSION

The data depicted in the Tab-1 showed significant variation in case of plant height, no. of branches and LAI of sesame at 45, 60 DAS due to application of various organic sources. The tallest plants and maximum no of branches were found in combined use of organic sources of nutrient i.e. 100% RDN through 50% of FYM and 50% of vermicompost + remaining RDP through rock phosphate (T₂) and 50% of RDN through FYM and vermicompost + remaining RDP through rock-phosphate + seed treatment with beejamrutha + application of panchagavya @3% as one soil application and two foliar application + application of jeevamrutha @10% as one soil application and two foliar application (T₁₀) over rest of the treatments. The findings are in close conformity with the results of Chaubey *et al.* (2003), Naugraiya and Singh (2004) with respect to plant height and number of branch plant⁻¹ in sesame. The increase in plant height also influenced by vermicompost and FYM due to the availability of plant growth regulation and humic acid, which is produced by increasing the activity of microbes in vermicompost (Sharma *et al.*, 2017) []. Nitrogen is associated with increase in protoplasm, cell division and cell enlargement resulting in taller plants (Tisdale *et al.*, 1985). Presence of naturally occurring, beneficial, effective microorganisms in Panchagavya predominantly, lactic acid bacteria, yeast, actinomycetes, photosynthetic bacteria and certain fungi besides beneficial and proven the beneficial effect especially in improving soil quality, growth and yield of crops (Xuand Xu, 2000 and Papen *et al.* (2002).

Maximum leaf area index was obtained with application of 100 % RDN through 50% of FYM and 50 % of vermicompost + remaining RDP through rock phosphate (T₂) (1.25, 1.39, 1.24 at 30, 45, 60 DAS and at harvest) which was found statistically at par with treatment i.e. 50 % RDN through 50% of FYM and 50 %of vermicompost + remaining RDP through rock phosphate + seed treatment with beejamrutha + application of panchagavya @ 3 % as one soil application and two foliar application+ application of jeevamrutha @ 10 % as one soil application and two foliar application (T₁₀) and significantly found superior over rest of the treatments.

Commented [M29]: After dot, a space should be considered, like this 3. RESULTS AND DISCUSSION

Commented [M30]: Plant height, number of braches plant⁻¹ and leaf area index (LAI) of summer sesame at 45 and 60 DAS showed significant variation due to application of various organic sources (Table 1).

Commented [M31]: Plant height

Commented [M32]: Number of branches plant⁻¹

Commented [M33]: Over other treatments

Commented [M34]: Chaubey *et al.* (2003) [number of reference], Naugraiya and Singh (2004) [number of reference].

Commented [M35]: Should be removed

Commented [M36]: (Sharma *et al.* 2017), [Number of reference]

Commented [M37]: (Tisdale *et al.* 1985), [Number of reference].

Commented [M38]: (Xu and Xu, 2000 and Papen *et al.* 2002), [Number of reference].

Commented [M39]: Maximum leaf area index (1.25, 1.39 and 1.24 at 30, 45, 60 DAS and at harvest) was obtained with application of 100 % RDN through 50% of FYM and 50 % of vermicompost + remaining RDP through rock phosphate (T₂).

The data relating to number of capsules plant⁻¹ as influenced by different organic sources are presented in Table 2. It is clear from table that application of 100 % RDN through 50% of FYM and 50 % of vermicompost + remaining RDP through rock phosphate (T₂) registered significantly higher in the number of capsules per plant (22.87) which was found statistically at par with application of 50 % RDN through 50% of FYM and 50 % of vermicompost + remaining RDP through rock phosphate + seed treatment with beejamrutha + application of panchagavya @ 3 % as one soil application and two foliar application + jeevamrutha @ 10 % as one soil application and two foliar application (T₁₀) (22.59) and was superior over the other treatments (T₈, T₇, T₆ and T₅). However, control treatment (T₁) indicated lowest no. of capsules plant⁻¹ (18.35) among all the treatments. The effect of organic sources on the number of seeds capsule⁻¹ depicted in Table-2 and appraisal of the data revealed that the highest seeds capsule⁻¹ (40.77) was observed with the treatment T₂ (100 % RDN through 50% of FYM and 50 % of vermicompost + remaining RDP through rock phosphate) and the minimum value was observed with control T₁ (29.97), respectively during the experimentation. Among other treatments, application of 50 % RDN through FYM and vermicompost + remaining RDP through rock phosphate + seed treatment with beejamrutha + application of jeevamrutha @ 10 % as one soil application and two foliar application (T₉) was found statistically similar (38.96) to treatment T₃ (35.98).

The seed yield of sesame was significantly influenced by different organic nutrient management practices and the data on seed yield is presented in Table-2. Among all the treatments, significantly highest yield of seed ha⁻¹ (808 kg ha⁻¹) was recorded by application of 100 % RDN through 50% of FYM and 50 % of vermicompost + remaining RDP through rock phosphate (T₂) which was statistically at par with the application of 50% of RDN through FYM and vermicompost + remaining RDP through rock phosphate + seed treatment with beejamrutha + application of panchagavya @ 3 % as one soil application and two foliar application + jeevamrutha @ 10 % as one soil application and two foliar application (T₁₀) (803 kg ha⁻¹) followed by 50 % RDN through FYM and vermicompost + remaining RDP through rock phosphate + seed treatment with beejamrutha + application of jeevamrutha @ 10 % as one soil application and two foliar application (T₆) (711 kg ha⁻¹) and 50 % RDN through FYM and vermicompost + remaining RDP through rock phosphate (T₃) (680 kg ha⁻¹). These treatments are superior over T₈, T₇, T₆, T₅ and T₄. The lowest seed yield was observed in control treatment T₁ (415 kg ha⁻¹). The combined application of FYM, vermicompost, rock phosphate and jeevamrutha results in better growth and yield attributes resulting in increased seed yield over FYM at 100 per cent N equivalent and without application of jeevamrutha thus, FYM and jeevamrutha can effectively and efficiently be used to get higher seed yield. Better seed yield due to balanced and timely supply of nutrients from diversified sources of nutrients (FYM, vermicompost, microbial consortia) that resulted in prolonged availability of nutrients to crop (Dharati *et al.*, 2017).

Among the different treatments, higher net returns (36,701 Rs. ha⁻¹ & 34,575 Rs. ha⁻¹) and returns per rupee investment were obtained maximum (1.14 and 1.01) with application of 100 % RDN through 50% of FYM and 50 % of vermicompost + remaining RDP through rock phosphate (T₂) and 50 % RDN through FYM and vermicompost + remaining RDP through rock

Commented [M40]: (22.59) (T₁₀).

Commented [M41]: Should be removed

Commented [M42]: Number

Commented [M43]: Shown in Table 2.

Commented [M44]: Recorded

Commented [M45]: value (29.97) was found in control treatment (T₁) respectively.

Commented [M46]: Should be removed

Commented [M47]: With T₃

Commented [M48]: The seed yield of summer sesame was significantly influenced by different organic nutrient management practices (Table 2).

Commented [M49]: Seed yield

Commented [M50]: Should be removed

Commented [M51]: Should be removed, Treatment number is enough

Commented [M52]: Should be removed

Commented [M53]: Should be removed

Commented [M54]: Should be removed

Commented [M55]: (Dharati *et al.* 2017), [reference number].

Commented [M56]: Should be mentioned the unit of this data.

phosphate + seed treatment with beejamrutha + application of panchagavya @ 3 % + jeevamrutha @ 10 % (T₁₀), respectively. The lowest net returns (15,225 Rs ha⁻¹) and returns per rupee investment (0.60) were obtained with seed treatment with beejamrutha 10 % solution + application of panchagavya @ 3 % (T₄) which might be due to the higher cost of cultivation and lesser yields. Similar views were expressed by Quddus *et al.* (2012), Jamkhogin (2015) and Sipai *et al.* (2016).

Commented [M57]: What is unit

Commented [M58]:

Commented [M59]: With application of

Commented [M60]: Similar results were found by Quddus *et al.* (2012) [reference number], Jamkhogin (2015) [reference number] and Sipai *et al.* (2016) [reference number].

UNDER PEER REVIEW

Table-1: Effect of Organic Nutrient Management Practices on Growth and Physiological Parameters of Sesamum

Commented [M61]: Table 1. Effect of Organic Nutrient Management Practices on growth behaviour of summer sesamum

Treatments	Plant Height(cm)		No of Branches ⁻¹		LAI	
	45 DAS	60 DAS	45 DAS	60 DAS	45 DAS	60DAS
T1: Control	24.06	56.14	2.28	2.48	1.05	1.18
T2: 100% RDN (50% FYM& 50 % VC) + remaining RP	41.32	72.9	2.87	3.64	1.25	1.39
T3: 50% RDN (50% FYM& 50% VC) + remaining RP	36.74	67.60	2.71	3.41	1.19	1.32
T4: Seed treatment with BM+ PG application (one soil & twofoliar spray)	26.62	60.9	2.34	3.05	1.09	1.2
T5: Seed treatment with BM+ JM application (one soil & twofoliar spray)	29.56	59.55	2.43	3.09	1.13	1.21
T6: Seed treatment with BM+ soil application of PG & JM	30.25	61.03	2.51	3.11	1.15	1.23
T7: Seed treatment with BM+ soil application & foliar sprayof PG & JM	33.18	63.87	2.57	3.26	1.17	1.26
T8: T3 + T4	34.11	64.85	2.65	3.32	1.18	1.29
T9: T3 + T5	37.65	68.77	2.75	3.48	1.21	1.34
T10: T3 + T4 + T5	40.30	71.3	2.81	3.53	1.23	1.36
S.Em (±)	0.91	0.97	0.06	0.11	0.02	0.02
CD (P =0.05)	2.55	2.73	0.15	0.30	0.05	0.06

Commented [M62]: Plant height (cm)

Commented [M63]: Title of these two columns should be merged

Commented [M64]: Number of Branches⁻¹

Commented [M65]: Title of these two columns should be merged

Commented [M66]: Leaf Area Index (LAI)

Commented [M67]: Title of these two columns

Commented [M68]: All of these should be Align center

*RDN- Recommended dose of nitrogen, BM- Beejamrutha, VC- Vermicompost, PG-Panchagavya, JM- Jeevamrutha, RP- Rock phosphate

Table-2: Effect of Organic Nutrient Management Practices on Yield, Yield Attributes and Economics of Sesamum

Treatments	No. of capsules plant ⁻¹	No. of seeds capsules ⁻¹	Seed yield (kg ha ⁻¹)	Net Return (Rs ha ⁻¹)	Return per rupee investment
	18.35	29.97	415	13263	0.60
T1: Control	22.87	40.77	808	36701	1.14
T2: 100% RDN (50% FYM& 50 % VC) + remaining RP	21.81	35.98	680	31031	1.12
T3: 50% RDN (50% FYM& 50% VC) + remaining RP	19.21	31.28	477	15325	0.60
T4: Seed treatment with BM+ PG application (one soil & twofoliar spray)	20.60	32.43	549	21113	0.83
T5: Seed treatment with BM+ JM application (one soil & twofoliar spray)	20.78	33.84	563	22767	0.90
T6: Seed treatment with BM+ soil application of PG & JM	21.31	35.65	643	27509	1.00
T7: Seed treatment with BM+ soil application & foliar sprayof PG & JM	21.22	36.80	646	24929	0.81
T8: T3 + T4	22.06	38.96	711	30163	0.97
T9: T3 + T5	22.59	40.21	803	34575	1.01
T10: T3 + T4 + T5	0.93	1.44	26.58	-	-
S.Em (±)	18.35	29.97	415	13263	0.60
CD (P =0.05)	22.87	40.77	808	36701	1.14

*RDN- Recommended dose of nitrogen, BM- Beejamrutha, VC- Vermicompost, PG-Panchagavya, JM- Jeevamrutha, RP- Rock phosphate

Commented [M69]: Table 2. Effect of Organic Nutrient Management Practices on yield attributes and yield of summer sesamum

Commented [M70]: Number of capsules plant⁻¹

Commented [M71]: Number of of seeds capsules⁻¹

Commented [M72]: All of these should be Align center

4. CONCLUSION

Based on the findings of the present investigation, the following conclusions could be drawn: Application of organic source of nutrients i.e. 100 % RDN through 50% of FYM and 50% of vermicompost + remaining RDP through rock phosphate (T₂) resulted in maximum growth parameters, yield attributes and yield of *sesamum*, but it was found to be statistically on par with combined application of 50 % of RDN through FYM & vermicompost + remaining RDP through rock phosphate + seed treatment with beejamrutha + application of panchagavya @ 3 % as one soil application & two foliar application + application of jeevamrutha @ 10 % as one soil application & two foliar application (T₁₀). In the organic nutrient treatments, application of 100 % RDN through 50% of FYM and 50 % of vermicompost + remaining RDP through rock phosphate resulted in maximum uptake of nutrients by seeds and stover and this was found to be the best combination for increasing the productivity of *sesamum* and also resulted in increased residual fertility levels of the soil compared to different sources organic nutrient combinations. Highest gross returns, net returns and return per rupee investment were obtained with application of 100 % RDN through 50% of FYM and 50 % of vermicompost+ remaining RDP through rock phosphate.

REFERENCE

1. Chaubey AK, Kaushik MK, Singh SB. Response of sesame (*Sesamum indicum* L.) to nitrogen and sulphur in light textured entisol. *New Agriculturist*. 2003; **14**(1/2): 61-64.
2. Dharati P, Patel R A and Sonaka Ghosh (2017). Effect of irrigation, vermicompost and sulphur on growth and yield of Summer sesamum (*Sesamum indicum* L.). *International Journal Current Microbiology and Applied Science*, **6**(11); 1647-1652.
3. Jamkhogin Lhungdim Devi, K N Influence of FYM, vermicompost and biofertilizers on economics and yields of urdbean (*Vigna mungo*) under acidic soil of NEH region. *Environment and Ecology* **33**(1) 73-77, 2015.
4. Naugraiya MN, Jhapatsngh P. Role of nitrogen and sulphur on performance of sesamum indicum under plantation of Dalbergia sissou in marginal land of Chhattisgarh. *Indian Journal of Agroforestry*. 2004; **6**(1): 89-91.
5. Papen, H A, Gabler, EZ and Rennenbeg, H, 2002, Chemolitho autotrophic nitrifiers in the phyllosphere of a spruce ecosystem receiving high nitrogen input. *Current Microbiology*, **44**: 55-60.

Commented [M73]: Summer sesamum

Commented [M74]: at

Commented [M75]: summer sesamum

Commented [M76]: source of

Commented [M77]: The number of references (9 references) is very few and should be increased the numbers of references to meet the criteria of an article.

Commented [M78]: Dharati P, Patel R A, Sonaka G.. Effect of irrigation, vermicompost and sulphur on growth and yield of Summer sesamum (*Sesamum indicum* L.). *International Journal Current Microbiology and Applied Science*. 2017; **6**(11): 1647-1652.

Commented [M79]: Jamkhogin Lhungdim DeK N Influence of FYM, vermicompost and biofertilizers on economics and yields of urdbean (*Vigna mungo*) under acidic soil of NEH region. *Environment and Ecology*. 2015; **33**(1) 73-77.

Commented [M80]: Papen HA, Gabler EZ Rennenbeg H. Chemolitho autotrophic nitrifiers in the phyllosphere of a spruce ecosystem receiving high nitrogen input. *Current Microbiology*. 2015; **44**: 55-60.

6. Quddus, M A, Rashid, M H, Hossain, M A, Naser H M and Abedin Mian, J Integrated nutrient management for sustaining soil fertility through chickpea-mungbean-taman cropping pattern at Madaripur region. *Bangladesh J. Agril.Res.*, 37(2), 251-262,2012.
7. Sharma J K, Jat, G Meena, R H, Purohit, H and Choudhary, R S (2017) Effect of vermicompost and nutrients application on soil, properties, yield, uptake and quality of Indian mustard (*Brassica juncea*). *Annals of Plant and Soil Research* 19(1):17-22.
8. Sipai, A H, Jat, J R and Rathore, B S Effect of Phosphorous, Sulphur and Biofertilizer on Growth Yield and Nodulation in Mungbean on Loamy Sand Soils of Kutch. *Crop Res*, 5(1), 2-4,2016.
9. Tisdale, S L, Nelson Werner L and Beaton James, D. 1985. *Soil fertility and fertilizers*. Mac publishing company, New York. pp: 437-448.

Commented [M81]: Quddus MA, Rashid MH, Hossain MA, Naser HM, Abedin MJ. Integrated nutrient management for sustaining soil fertility through chickpea-mungbean-taman cropping pattern at Madaripur region. *Bangladesh J. Agril.Res.*; 2012; 37(2): 251-262.

Commented [M82]: Sharma JK, Jat G, Meena, RH, Purohit H, Choudhary, RS. Effect of vermicompost and nutrients application on soil. properties, yield, uptake and quality of Indian mustard (*Brassica juncea*). *Annals of Plant and Soil Research*. 2017; 19(1):17-22.

Commented [M83]: Sipai AH, Jat JR, Rathore BS. Effect of Phosphorous, Sulphur and Biofertilizer on Growth Yield and Nodulation in Mungbean on Loamy Sand Soils of Kutch. *Crop Res*, 2016; 5(1): 2-4.

Commented [M84]: Tisdale S L, Nelson WL, Beaton JD. *Soil fertility and fertilizers*. Mac publishing company, New York. 1985; pp: 437-448.

UNDER PEER REVIEW