

Evaluation of physiochemical properties and WQI of the drinking water in the urban area of Bilaspur

ABSTRACT

Healthy and sustained environment is strictly dependent on the supply of clean and safe water. Present study was estimating the physicochemical parameters of water and assessment of water quality index of drinking water in residential area of Bilaspur city. Fifty water samples were collected from ten sites and analyzed six parameters of water quality by using the portable multi-parameter water quality meter (Hanna Instruments: HI98194). The data was collected in every site for selected parameters. Results were significant differs ($p < 0.001$) for sites for water quality parameters. The results were observed for water pH, water temperature, dissolved oxygen, total dissolved solids, electrical conductivity of water, and oxygen reduction potential, which ranged from 7.89 ± 0.11 to 9.3 ± 0.03 , 26.38 ± 0.25 °C to 28.86 ± 0.17 °C, 7.22 ± 0.02 to 8.14 ± 0.02 mg/l, 239.6 ± 0.68 to 496.2 ± 0.58 mg/l, 357.4 ± 0.93 to 686 ± 0.95 μ S/cm, and -38.65 to -22.82 mV, respectively. Study revealed the very strong positive correlation between EC and TDS ($r = 0.935$) and between pH and ORP ($r = 0.802$) while the significant negative correlation observed between DO and temperature. The WQI was statistically differs at ($p < 0.001$) and the range of WQI was observed from 383.67 ± 1.69 to 530.87 ± 1.25 . The WQI results were showed the water is not safe for drinking purpose.

Key words: EC, Urban water quality, water monitoring, TDS, sustainable environment.

1. INTRODUCTION

Water, the essence of life, is a vital resource on earth. 70% of the earth's mass is water, a constituent of living things, of which 2.8% is freshwater (20% groundwater) and the remaining 97.2% is saltwater [1]. Ground and surface water have been exceptionally used for drinking purpose in urban area. Water quality has been affected by human activities such as agricultural practices, urban solid waste disposal, and improper disposal of industrial effluent [2]. Quality of

water influenced human health [3]. Inappropriate uses and unsustainable waste management techniques are causing the water quality to worsen in cities [4, 5, 6, 59]. The trouble is getting dangerous because water treatment is either non-existent or insufficient. Pollutants pass through sewage systems and contaminate surfaces and groundwater [7, 8, 9, 10]. Polluted water has caused 2.2 million deaths, of which 1.4 million are children, and the number of patients suffering from waterborne diseases has been increasing worldwide [11]. A healthy and balanced world is strictly dependent on the supply of clean and safe water [12, 13]. In the meeting of UN members (held in September 2015), for the 2030 Agenda, adopted the sustainable goal “Ensure availability and sustainable management of water and sanitation for all” [14,15]. Minimizing the water contamination and uses for drinking is first priority for the nation [16]. Water quality parameters consist of physical, chemical, and microbiological characteristics, includes pH, turbidity, dissolved oxygen, total dissolved solids, oxidation reduction potential, temperature, hardness, electrical conductivity etc [17,18,19].

Bilaspur is the second largest city of Indian states of Chhattisgarh and known as the city of festivals. About 2.5 lakh people live in the city, which sharply increased in 2022 to about 3.4 [20]. Increasing population around the city creates huge pressure on water sources, which results in a decreasing groundwater level. Arpa River is main river passes between the city. It is the headquarters of South East Central Railway and South Eastern Coalfield Limited. Due to the enormous number of coal mine near the city; coal based major and minor industries (thermal power plants, iron melting industries, etc.) are existence in surroundings of the city and disposing their effluents without proper management techniques. Hence, the ground and surface water quality has been affected through it. In present study intends to assess the physio-chemical quality of drinking water from different sites of Bilaspur city (India) and to ensure that the water is either safe for drinking or not.

2. MATERIALS AND METHODS

2.1 STUDY AREA

The study was conducted during the period of March to Jun, 2023. Water samples collected from the different residential area of Bilaspur city (Table 1). It is situated at 22.09° N 82.15°E and has an average elevation of 264 meters form sea level. The climate of the area is tropical; the average temperature and rainfall was range from 20.37°C to 35.75°C and 764 mm recorded, respectively.

Table 1 Shown sites for water sample collection.

Sites	Place	Latitude	Longitude	Source of water
S1	Koni	22.134125 N	82.123842 E	Tube well, water tank
S2	SECL, Indira Vihar colony	22.106386 N	82.147844 E	Water tank
S3	Lodhipara	22.100774 N	82.140638 E	Water tank
S4	Ashok Nagar	22.102597 N	82.152661 E	Water tank
S5	Satyam Chowk	22.078887 N	82.156474 E	Water tank
S6	Mangla Chowk	22.091352 N	82.127432 E	Water tank
S7	Seepat Chowk	22.095562 N	82.144719 E	Water tank
S8	Devkinandan Street	22.088342 N	82.148754 E	Water tank, tube well
S9	DFO, colony	22.083050 N	82.140649 E	Water tank, tube well
S10	Birkona	22.140783 N	82.154384 E	Water tank, tube well

2.2 WATER SAMPLING AND ANALYSIS

Fifty samples of tap water were collected from ten sites, and five replications for each site were analyzed. The six parameters of water quality were analyzed (includes temperature, electrical conductivity, pH, total dissolved solids, dissolved oxygen, and oxygen reduction potential) using a portable multi-parameter water quality meter (Hanna Instruments Model: HI98194).

The WQI has been measured as per the drinking water quality standards recommended by the Bureau of Indian Standards (IS-10500:2012). The weighted arithmetic approach was used to create the water quality index [21, 22] and that equation has been completed into three steps which given below:

(1) To calculate the unit weight (W_i) factors from each parameters by using the formula:

$$W_i = \frac{K}{S_n}$$

Where, W_i = unit weight of i^{th} parameters

$$K = \frac{1}{\sum(\frac{1}{S_n})}$$

K = proportion constant, S_n = standard desirable value of the i^{th} parameters

(2) To calculate the sub-index value Q_i value by using the formula:

$$Q_i = \left[\frac{V_o - V_i}{S_n - V_i} \right] * 100$$

Where, V_o = concentration of i^{th} parameters at a giving sampling sites, V_i = ideal value of i^{th} parameters in pure water.

(3) To calculate the water quality index by using the formula:

$$WQI = \frac{\sum Wi * Qi}{\sum Wi}$$

With the exception of pH, dissolved oxygen, and temperature all ideal values (V_i) for drinking water are assumed to be zero [23]. Ideal Value of pH is 7.0 while maximum acceptable pH is 8.5 for drinking water. Similarly for dissolved oxygen the permissible limit is 5 mg/l while the ideal value is 14.6 mg/l. The permissible limit of water temperature range from 25 to 28°C and ideal value is 25°C. Thus, effective values for water temperature, pH and dissolved oxygen were calculated using these formulas:

$$Q_{pH} = \left[\frac{(V_{pH} - 7.0)}{(8.5 - 7.0)} \right] * 100$$

$$Q_{DO} = \left[\frac{(V_{DO} - 14.6)}{(5.0 - 14.6)} \right] * 100$$

$$Q_T = \left[\frac{(V_T - 28)}{(25 - 28)} \right] * 100$$

Where V_{DO} is observed value dissolved oxygen, V_{pH} is observed value of pH and V_T is observed value of temperature.

2.3 STATISTICAL ANALYSIS

A one-way ANOVA was performed at the 5% level of significance to analyze the variation in sampling sites for parameters using SPSS software (IBM 20.0). The matrix correlation was performed to determine the relationship among the parameters of water quality.

3. RESULTS AND DISCUSSIONS

3.1 WATERPH

The water pH is altered through electrolysis, or the splitting of the water molecule into hydrogen and hydroxide ions with an electric current. The pH of water indicates its acidity and basicity and scale of pH is <7.0 for acidic, 7.0 for neutral and >7.0 for salinity or alkanity [59, 60]. It detects the ratio of free hydroxyl and hydrogen ions present in water. The water pH was statistically significant differ ($p < 0.001$) in sites. The higher water pH values received 9.3 ± 0.03 for S1 followed by $S2 > S3 > S10 > S4 > S5 > S6 > S9$ and S8. The lower water pH was observed 7.89 ± 0.11 for S7 (Table 2). Two sites S1 and S2 showed pH value water more than 8.5 which is not considerable for drinking while other sites is safe. A study was reported in drinking water pH about more than 7.40 [24, 25, 26]. Significant higher or lower pH can be harmful for human body and not safe for drinking purposes. High pH results in an unpleasant taste; it reduces the efficacy

of chlorine disinfection, necessitating the use of extra chlorine. Metals and other materials corrode or dissolve in low-pH water [26, 27].

3.2 WATER TEMPERATURE

Water temperature is a significant influence on drinking water quality. High temperature increases the viscosity of drinking water sharply. When temperature rises from 5 to 25 °C, the level of viscosity will be falls by approximately 40%, thus lowering flow resistance. It also has effects on the mobility of copper ions, the degree of deterioration, the dissolution of metal structures, the overall chlorinated breakdown, and the production of disinfection products [28, 29]. Drinking water should have an ideal temperature of 25 °C, as per the World Health Organization[30].The reported the negative correlation in dissolved oxygen and water temperature and were dissolved oxygen of water decreased with increases the water temperature [31, 32]. The observation showed the water temperaturesignificant differs at ($P = 0.001$) level of significant. S6 was received highest water temperature at 28.86 ± 0.17 °C followed by S7> S3> S5> S8> S4> S9> S10> and S1. The minimum temperature of water was recorded 26.38 ± 0.25 °C for S2(Table 2).Similar report was found for water temperature of Wondo Genet city ranges between 28°C and 29 °C [31]. The acceptable drinking water temperature limit of WHO is 25-30 °C. Water temperature range of 29 °C was reported for drinking water in Nigeria [33]. The drinking water temperature in tropical countries could be even higher the city of Cali (Colombia) ranges of watertemperatures between 25°C and 28°C [29].

3.3 DISSOLVED OXYGEN

All aquatic species depend on dissolved oxygen. Whenever dissolved oxygen within water remains unsaturated, atmospheric oxygen can enter a water body; likewise, aquatic plants may produce dissolved oxygen during photosynthesis[34, 35]. The dissolved oxygen of water is inversely proportional to the water temperature; when the water temperature increases, the dissolved oxygen decreases. DO level is the sign the water self purification [36]. The dissolved oxygen of water for study areas was significant differs at($P = 0.001$) level of significant. The highest dissolved oxygen (mg/l) was received 8.14 ± 0.02 mg g-1for S2 followed by S7> S10> S9> S4> S8> S3> S5> and S7, while the lowest was 7.22 ± 0.02 mg g-1 for S6 recorded (Table 2).Bwire et al. (2020) reported the dissolved oxygen of 6.38 mg/l at temperature of 26.57 °C for domestic drinking water [37].

3.4 ELECTRICAL CONDUCTIVITY

Electrical conductivity ($\mu\text{S}/\text{cm}$) of water for different sites was differ significantly at ($P = 0.001$) level of significant. S9 was recorded higher electrical conductivity of water for $686 \pm 0.95 \mu\text{S}/\text{cm}$ followed by $S5 > S1 > S4 > S7 > S6 > S8 > S2 > S3$, while the lowest was $357.4 \pm 0.93 \mu\text{S}/\text{cm}$ for S10 observed. Pure water is an excellent insulator but nevertheless a strong conductor of currents [60]. As the ions in the water increases, the electrical conductance also increases. EC in water often determined by the concentrated of dissolved suspended particles in water [31, 38, 39]. Present study was reported electrical conductivity ranges from $569 - 569 \mu\text{Scm}^{-1}$. $300 \mu\text{Scm}$ electrical conductivity of the water is acceptable limit for drinking. An investigation was reported higher range of EC for drinking water quality of city Pogradec, Albania [40]. Similar data was reported higher range of EC $770 \mu\text{Scm}^{-1}$ [41, 42, 43].

3.5 TOTAL DISSOLVED SOLIDS

The TDS (mg/l) of water varied statistically ($P = 0.001$). The maximum TDS was observed $496.2 \pm 0.58 \text{ mg}/\text{l}$ for S5 followed by $S9 > S4 > S1 > S7 > S2 > S6 > S8$ and than S3. The $239.6 \pm 0.68 \text{ mg}/\text{l}$ for S10 was observed minimum water total dissolved solid (Table 2). The entirely of the dissolved component in water that remain dehydrated are known as total dissolved solids. Soluble salts compose almost all of the nonvolatile or fixed components of TDS that exist in natural waters [44]. This part might be known as water salinity. Inland water salinity tends to contain four major cations (Ca, Mg, Na, and K) and three anions (bicarbonate, sulphate, and chloride) [43, 45]. The TDS ranges between $500 - 2000 \text{ mg}/\text{l}$ is maximum permissible limit [46]. These limits are not always observed, however, and many municipalities use water containing from $2,000$ to $4,000 \text{ mg}/\text{l}$ TDS [47]. Present study maximum TDS observed $496 \text{ mg}/\text{l}$ for drinking water. Similar investigation was reported of TDS for surface drinking water $> 490 \text{ mg}/\text{l}$ [48, 49, 50].

3.6 OXIDATION REDUCTION POTENTIAL

Electrons are transferred among atoms, molecules, or ions in a process known as a redox reaction. The electrical force generated within water in the presence of an oxidant or reductant is determined by the test ORP [51]. Drinking water should have an ORP of about -50 mV . When ORP is positive, situations are oxygenating; often used as a gauge for disinfectant concentrations in untreated discharge water, like chlorine. The ORP was influenced by the existence of chlorine water, which can cause value to move towards positive [52, 53]. The correlation between ORP and pH was also reported by Vongvichiankul et al. (2017) the degree of deference was decreased

with increasing pH [54]. Present study observed the ORP (mV) of municipal tap water was significant differs for sites at ($P = 0.001$) level of significance (Table 2). The range of ORP was recorded from -38.65 (S10) to -22.82 (S1). Similar result was reported by Luccarini et al. and Azis et al. [55, 56].

Table 2 Analyzed value of different samples for water parameters

Sites	pH	Temperature (°C)	DO (mg/l)	EC (µS/cm)	TDS (mg/l)	ORP (mV)
S1	9.3±0.03 ^a	26.56±0.34 ^{de}	8.07±0.04 ^g	586.2±1.85 ^c	437.8±1.16 ^c	-22.82±0.25 ^a
S2	8.85±0.04 ^b	26.38±0.25 ^e	8.14±0.02 ^f	475.6±1.94 ^g	405±1.55 ^d	-26.38±0.21 ^b
S3	8.45±0.06 ^c	27.77±0.22 ^b	7.68±0.01 ^e	391.6±2.01 ^h	280.2±1.77 ^g	-34.28±0.13 ^e
S4	8.23±0.08 ^d	27.18±0.15 ^{bcd}	7.86±0.01 ^{de}	574.8±2.6 ^d	439.6±1.66 ^c	-34.04±0.36 ^{de}
S5	8.18±0.06 ^{de}	27.55±0.18 ^b	7.57±0.14 ^{cd}	608±3.75 ^b	495±1.82 ^a	-32.28±0.26 ^c
S6	8.17±0.07 ^{de}	28.86±0.17 ^a	7.22±0.02 ^{bc}	527.2±3.22 ^e	375±2.02 ^e	-37.24±0.23 ^f
S7	7.89±0.11 ^f	28.51±0.22 ^a	7.37±0.02 ^{ab}	570.2±2.15 ^d	409±1.52 ^d	-33.44±0.19 ^d
S8	7.95±0.1 ^f	27.28±0.2 ^{bc}	7.81±0.02 ^a	494.2±1.02 ^f	357.6±1.69 ^f	-34.64±0.19 ^e
S9	7.97±0.07 ^{ef}	26.81±0.11 ^{cde}	7.99±0.03 ^a	686±0.95 ^a	481±1.45 ^b	-33.38±0.11 ^d
S10	8.27±0.05 ^{cd}	26.59±0.23 ^{de}	8.04±0.03 ^a	357±2.55 ⁱ	240.4±1.44 ^h	-38.65±0.2 ^g

Mean±SE in the same column with different letters in superscript is significantly different ($P = 0.001$)

Table 3 Correlation matrix between physiochemical parameters

Parameters	pH	Temperature (°C)	DO (mg/l)	EC (µS/cm)	TDS (mg/l)	ORP (mV)
pH	1					
Temperature (°C)	-0.5165	1				
DO (mg/l)	0.524321	-0.987	1			
EC (µS/cm)	-0.13946	0.068883	-0.10866	1		
TDS (mg/l)	0.011174	-0.03286	-0.03423	0.935806	1	
ORP (mV)	0.802626	-0.47455	0.467226	0.316448	0.475399	1

Correlation between the physiochemical parameters of water quality has been analyzed using correlation matrix by MS Excel (Table 3). The correlation value was categorized into four classes as very strong correlation (r value greater than 0.75), moderate correlation (r value between 0.50 - 0.75), lower correlation (r value between 0.30 - 0.50), and weak correlation (r

value is <0.30). Present results showed significantly strong positive correlation between EC and TDS ($r = 0.935$) and between pH and ORP ($r = 0.802$). A moderate positive correlation was recorded between DO to pH ($r = 0.524$) and ORP ($r = 0.467$), between ORP to EC ($r = 0.316$) and TDS ($r = 0.475$). While the strong negative correlation was recorded between DO and temperature ($r = -0.987$). Comparison between mean study values with different parameters of quality of water and Bureau of Indian Standard (2022) as shown in Fig. 1.




Figure 1 Shown the comparison between mean values and BIS standard

3.7 WATER QUALITY INDEX

The water quality index of selected sites was analyzed on the basis of six types of water quality parameters includes pH, water temperature, dissolved oxygen, total dissolved solids, electrical conductivity of water, and oxygen reduction potential. A significant difference in WQI was at ($p < 0.001$) level of significance observed. The highest water quality index was at S7 for 530.87 ± 1.25 , followed by S2 (514.88 ± 1.75), S6 (512.17 ± 1.44), S3 (500.46 ± 1.44), S5 (495.08 ± 1.44), S4 (472.27 ± 1.04), S8 (462.57 ± 1.21), S9 (454.01 ± 1.58), and S10 (415.3 ± 1.37) (Fig. 2). While S1 showed the lowest water quality index value, about 383.67 ± 1.69 . Different researchers give different categorization of water quality index. According to Al-Musawi et al. (2018) the WQI was divided into six categories i.e., excellent (<50), good (50-100), poor (100-200), very poor (200-300), polluted (300-400), and very polluted (>400) [57]. This study was revealed all the collected water samples were fell into the

polluted category and not safe for drinking. Khudair Al-Musawi reported WQI of Tigris and Euphrates Rivers, Iraq was more than 300 for years from 2013 to 2014 which were not safe for drinking [58].




Figure 2 WQI for different sampling sites

4. CONCLUSION

The physiochemical parameters of drinking water were analyzed and compared with BIS (2012). However, electrical conductivity not in satisfactory level as per national standard, while the pH, TDS, DO, temperature and ORP values were within considerable limit. On the basis of selected physiochemical water quality parameters the water quality index (WQI) was fall into polluted category for all sites therefore it is not safe for drinking.

ABBREVIATIONS

- ORP = oxidation reduction potential
- DO = dissolved oxygen
- WQI = water quality index
- EC= electrical conductivity
- TDS = total dissolved solids
- ANOVA = analysis of variance
- WHO= world health organization
- BIS = bureau of Indian standards
- mV = millivolts
- mg/l = milligram per liter

$\mu\text{S}/\text{cm}$ = microSiemens per centimeter

mg g⁻¹ = milligram per gram

⁰C = degree Celsius

mm= millimeter

REFERENCES

1. Gupta, S. K., and Gupta, I. C. Management of saline & waste water in agriculture. Scientific Publishers. 2015; 1- 41.
2. Howladar, M. F., Chakma, E., Koley, N. J., Islam, S., Al Numanbakhth, M. A., Ahmed, Z. and Akter, S. The water quality and pollution sources assessment of Surma river, Bangladesh using, hydrochemical, multivariate statistical and water quality index methods. Groundwater for sustainable developent. 2021;12, 100523.<https://doi.org/10.1016/j.gsd.2020.100523>
3. Kulakova, E. S. Evaluation of the exposure of the influence of chloride ions in the water of r. Belaya in the area of the Sterlitamak city of the republic of Bashkortostan on human health. In Сборник материалов международной научно-практической конференции "Здоровье и окружающая среда. 2019;48-49.
4. Putra, F. G., Sari, A. P., Qurotunnisa, A., Rukmana, A., Darmayanti, R., and Choirudin, C. What are the advantages of using leftover cooking oil waste as an aromatherapy candle to prevent pollution?. Jurnal Inovasi Dan Pengembangan Hasil Pengabdian Masyarakat. 2023; 1 (2), 59-63.<https://doi.org/10.61650/jip-dimas.v1i2.230>
5. Ahsan, W. A., Ahmad, H. R., Farooqi, Z. U. R., Sabir, M., Ayub, M. A., Rizwan, M., & Ilic, P. Surface water quality assessment of Skardu springs using Water Quality Index. Environmental Science and Pollution Research. 2021; 28, 20537-20548.<https://doi.org/10.1007/s11356-020-11818-5>
6. Kumari, A., and Gupta, Y. K. Physico-chemical assessment of tube wells water quality of Pilani town district Jhunjhunu, Rajasthan, India. Research Journal of Science and Technology. 2022; 14 (2), 91-94.<http://dx.doi.org/10.52711/2349-2988.2022.00014>
7. Akhtar, N., Syakir Ishak, M. I., Bhawani, S. A., and Umar, K. 2021. Various natural and anthropogenic factors responsible for water quality degradation: A review. Water. 13(19), 2660.<https://doi.org/10.1007/s11356-020-11818-5>
8. D'alessandro, D., Gola, M., Appolloni, L., Dettori, M., Fara, G. M., Rebecchi, A. and Capolongo, S. COVID-19 and living space challenge. Well-being and public health

recommendations for a healthy, safe, and sustainable housing. *Acta Bio Medica: AteneiParmensis*. 2020; 91(9-S), 61. doi: [10.23750/abm.v91i9-S.10115](https://doi.org/10.23750/abm.v91i9-S.10115)

9. Naidu, R., Espana, V. A. A., Liu, Y., and Jit, J. Emerging contaminants in the environment: Risk-based analysis for better management. *Chemosphere*. 2016; 154, 350-357. <https://doi.org/10.1016/j.chemosphere.2016.03.068>
10. Bitton G. *Microbiology of Drinking Water Production and Distribution*. 1st ed. John Wiley & Sons, Inc.; Hoboken, NJ, USA. 2014; 312. DOI:10.1002/9781118743942
11. World Health Organization (WHO) *Water Sanitation and Health*. 2015. http://www.who.int/water_sanitation_health/diseases
12. Rosborg, I., and Kozisek, F. *Drinking water minerals and mineral balance*. Springer International Pu. 2016; 127-148 <https://doi.org/10.1007/978-3-030-18034-8>
13. Magana-Arachchi, DN and Wanigatunge R.P., Ubiquitous waterborne pathogens. *Waterborne Pathogen*. 2020; 15–42. doi: [10.1016/B978-0-12-818783-8.00002-5](https://doi.org/10.1016/B978-0-12-818783-8.00002-5)
14. United Nations General Assembly Resolution (UNSD). *Transforming Our World: The 2030 Agenda for Sustainable Development*. 2015; A/RES/70/1
1. Weststrate, J., Dijkstra, G., Eshuis, J., Gianoli, A., and Rusca, M. The sustainable development goal on water and sanitation: learning from the millennium development goals. *Social Indicators Research*. 2019; 143, 795-810. <https://doi.org/10.1007/s11205-018-1965-5>
15. Jaramillo, F., Desormeaux, A., Hedlund, J., Jawitz, J. W., Clerici, N., Piemontese, L., ... and Åhlén, I. Priorities and interactions of sustainable development goals (SDGs) with focus on wetlands. *Water*. 2019; 11 (3), 619. <https://doi.org/10.3390/w11030619>
16. Daud M.K., Nafees M, Ali S, Rizwan M, Bajwa R.A. and Shakoor M.B. *Drinking Water Quality Status and Contamination in Pakistan*. 2017. <https://doi.org/10.1155/2017/7908183>
17. Ford L., Bharadwaj L., Mcleod L. and Waldner C. Human health risk assessment applied to rural populations dependent on unregulated drinking water sources: a scoping review. 2017. <https://doi.org/10.3390/ijerph14080846>
18. Post, C.J., Cope, M.P., and Gerard, P.D. Monitoring spatial and temporal variation of dissolved oxygen and water temperature in the Savannah River using a sensor network. *Environ Monit Assess*. 2018; 190, 272 <https://doi.org/10.1007/s10661-018-6646-y>.

19. Shah, M., Vijayshankar, P. S., and Rural, B. Tribal Development Report: Human Development and Governance. Taylor & Francis.2022.
20. Unigwe, C. O., andEgbueri, J. C. Drinking water quality assessment based on statistical analysis and three water quality indices (MWQI, IWQI and EWQI): a case study. Environment, Development and Sustainability.2023; 25(1), 686-707. <https://doi.org/10.1007/s10668-021-02076-7>
21. Adimalla, N., and Qian, H. Groundwater quality evaluation using water quality index (WQI) for drinking purposes and human health risk (HHR) assessment in an agricultural region of Nanganur, south India. Ecotoxicology and environmental safety. 2019; 176, 153-161.<https://doi.org/10.1016/j.ecoenv.2019.03.066>
22. Ram, A., Tiwari, S. K., Pandey, H. K., Chaurasia, A. K., Singh, S., and Singh, Y. V. Groundwater quality assessment using water quality index (WQI) under GIS framework. Applied Water Science. 2021; 11, 1-20.<https://doi.org/10.1007/s13201-021-01376-7>
23. Yousefi, M., Saleh, H. N., Mahvi, A. H., Alimohammadi, M., Nabizadeh, R., and Mohammadi, A. A. Data on corrosion and scaling potential of drinking water resources using stability indices in Jolfa, East Azerbaijan, Iran. Data in brief. 2018; 16, 724-731.<https://doi.org/10.1016/j.dib.2017.11.099>
24. Alsulaili, A., Al-Harbi, M., & Al-Tawari, K. Physical and chemical characteristics of drinking water quality in Kuwait: tap vs. bottled water. Journal of engineering research. 2015; 3, 1-26.<https://doi.org/10.7603/s40632-015-0002-y>
25. Pantelić, N. Đ., Dramićanin, A. M., Milovanović, D. B., Popović-Đorđević, J., and Kostić, A. Ž. Evaluation of the quality of drinking water in Rasina district, Serbia: physicochemical and bacteriological viewpoint. Romanian Journal of Physics. 2017; 62 (9-10).<https://cherry.chem.bg.ac.rs/handle/123456789/2070>
26. Islam, R., Faysal, S. M., Amin, R., Juliana, F. M., Islam, M. J., Alam, J., and Asaduzzaman, M. Assessment of pH and total dissolved substances (TDS) in the commercially available bottled drinking water. IOSR Journal of Nursing and health Science. 2017; 6 (5), 35-40.
27. Ljiljana Z., Andreas M., Jan Peter V., Jan V.andMirjam B. Development and validation of a drinking water temperature model in domestic drinking water supply systems. Urban Water Journal. 2017; 14 (10): 1031-1037, DOI: [10.1080/1573062X.2017.1325501](https://doi.org/10.1080/1573062X.2017.1325501)

28. Agudelo-Vera, C., Avvedimento, S., Boxall, J., Creaco, E., De Kater, H., Di Nardo, A. and Blokker, M. Drinking water temperature around the globe: understanding, policies, challenges and opportunities. *Water*. 2020; 12(4), 1049.<https://doi.org/10.3390/w12041049>
29. World Health Organization (WHO) and the United Nations Children's Fund (UNICEF); Geneva. Progress on Drinking Water, Sanitation and Hygiene: Update and Sustainable Development Goal Baselines. License: 2017; CC BY-NC-SA 3.0 IGO. <https://policycommons.net/artifacts/421754/progress-on-drinking-water-sanitation-and-hygiene/1392737/>
30. Meride, Y., and Ayenew, B. Drinking water quality assessment and its effects on residents health in Wondo genet campus, Ethiopia. *Environmental Systems Research*.2016; 5 (1), 1-7.<https://doi.org/10.1186/s40068-016-0053-6>
31. Danladi Bello, A. A., Hashim, N. B., and Mohd Haniffah, M. R. Predicting impact of climate change on water temperature and dissolved oxygen in tropical rivers. *Climate*. 2017; 5(3), 58.
32. Bamigboye, C. O., Amao, J. A., Ayodele, T. A., Adebayo, A. S., Ogunleke, J. D., Abass, T. B. and Oyedemi, A. A. An appraisal of the drinking water quality of groundwater sources in Ogbomoso, Oyo state, Nigeria. *Groundwater for Sustainable Development*. 2020; 11, 100453.<https://doi.org/10.1016/j.gsd.2020.100453>
33. Suplee, M. W., Sada, R., and Feldman, D. L. Aquatic plant and dissolved oxygen changes in a reference-condition prairie stream subjected to experimental nutrient enrichments. *JAWRA Journal of the American Water Resources Association*. 2019; 55 (3), 700-719.<https://doi.org/10.1111/1752-1688.12736>
34. Shamsudin, M. S., Sapingi, H. H. J., and Aziz, M. S. A. Optical glass micro-fibre based transducers for dissolved oxygen sensing and monitoring: an overview. In *Journal of Physics: Conference Series*.2020; 1484 (1): 012004DOI 10.1088/1742-6596/1484/1/012004
35. Wei, Y., Jiao, Y., An, D., Li, D., Li, W., and Wei, Q. Review of dissolved oxygen detection technology: From laboratory analysis to online intelligent detection. *Sensors*. 2019; 19 (18), 3995.<https://doi.org/10.3390/s19183995>
36. Bwire, G., Sack, D.A., and Kagirita, A. The quality of drinking and domestic water from the surface water sources (lakes, rivers, irrigation canals and ponds) and springs in cholera prone communities of Uganda: an analysis of vital physicochemical parameters. *BMC Public Health*. 2020; 20, 1128 <https://doi.org/10.1186/s12889-020-09186-3>.

37. Bhateria, R., and Jain, D. Water quality assessment of lake water: a review. *Sustainable Water Resources Management*. 2016; 2, 161-173.<https://doi.org/10.1007/s40899-015-0014-7>
38. Corwin, D. L., and Yemoto, K. Salinity: Electrical conductivity and total dissolved solids. *Soil science society of America journal*. 2020; 84 (5), 1442-1461. <https://doi.org/10.2136/sssabookser5.3.c14>
39. Eftimi, R., and Zojer, H. Human impacts on karst aquifers of Albania. *Environmental Earth Sciences*. 2015; 74, 57-70.
40. Hurra, W. A., and Bhawsar, A. Assessment of ground water quality near solid waste dumping site Bhanpur, Bhopal. *IJAR*. 2021; 7 (8), 314-319.<http://www.allresearchjournal.com/>
41. Tyor, A. K., and Devi, P. Ecological Studies of Barwala Link Canal in Narwana Region, Haryana. *Current World Environment*. 2017; 12 (1), 116.
42. Hussien, B. M., Lattoofi, N. F., Abd-alghafour, N. M., Muslim, R. F., Zaidan, T., and Mahmood, M. Space-time hydrochemical variations and water quality of Euphrates river. *Arabian Journal of Geosciences*. 2021; 14, 1-19.<https://doi.org/10.1007/s12517-021-07794-w>
43. Wilson, J. M., Wang, Y., and VanBriesen, J. M. Sources of high total dissolved solids to drinking water supply in southwestern Pennsylvania. *Journal of Environmental Engineering*. 2014; 140 (5), B4014003.
44. Dugan, H. A. Salinity and Ionic Composition of Inland Waters. In *Wetzel's Limnology*. 2024; 75-299.<https://doi.org/10.1016/B978-0-12-822701-5.00012-4>
45. BIS, Standard, D. I. Bureau of Indian Standards. 2022.
46. Iloms, E., Ololade, O. O., Ogola, H. J., and Selvarajan, R. Investigating industrial effluent impact on municipal wastewater treatment plant in Vaal, South Africa. *International journal of environmental research and public health*. 2020; 17(3), 1096.<https://doi.org/10.3390/ijerph17031096>
47. Aksever, F., Davraz, A., and Bal, Y. Assessment of water quality for drinking and irrigation purposes: a case study of Başköy springs (Ağlasun/Burdur/Turkey). *Arabian Journal of Geosciences*. 2016; 9, 1-18.

48. Ullah, S., Javed, M. W., Rasheed, S. B., Jamal, Q., Aziz, F., and Ullah, S. Assessment of groundwater quality of district Dir Lower Pakistan. *International Journal of Biosciences*. 2014; 4 (8), 248-255.. <http://dx.doi.org/10.12692/ijb/4.8.248-255> A
49. Princela, M. A., Jose, J. M. A., Gladis, E. E., Arthi, D., and Joseph, J. 2021. Regional assessment of groundwater quality for drinking purpose. *Materials Today: Proceedings*. 45, 2916-2920. <https://doi.org/10.1007/s10661-011-2171-y>
2. Taghizadeh, A., Taghizadeh, M., Jouyandeh, M., Yazdi, M. K., Zarrintaj, P., Saeb, M. R., and Gupta, V. K. Conductive polymers in water treatment: A review. *Journal of Molecular Liquids*. 2020; 312, 113447. <https://doi.org/10.1016/j.molliq.2020.113447>
50. Copeland, A., and Lytle, D. A. Measuring the oxidation–reduction potential of important oxidants in drinking water. *Journal-American Water Works Association*. 2014; 106 (1) E10-E20. <https://doi.org/10.5942/jawwa.2014.106.0002>
51. Ahuja, S. *Evaluating water quality to prevent future disasters*. Academic Press. 2019.
52. Vongvichiankul, C., Deebao, J., and Khongnakorn, W. Relationship between pH, oxidation reduction potential (ORP) and biogas production in mesophilic screw anaerobic digester. *Energy Procedia*, 2017; 138, 877-882.
53. Azis, K., Ntougias, S., and Melidis, P. NH₄⁺-N versus pH and ORP versus NO₃⁻-N sensors during online monitoring of an intermittently aerated and fed membrane bioreactor. *Environmental Science and Pollution Research*. 2021; 28, 33837-33843.
54. Luccarini, L., Pulcini, D., Sottara, D., Di Cosmo, R., and Canziani, R. Monitoring denitrification by means of pH and ORP in continuous-flow conventional activated sludge processes. *Desalination and Water Treatment*. 2017; 61, 319-325. <https://hdl.handle.net/11311/1038571>
55. Al-Musawi, N. O., Al-Obaidi, S. K., and Al-Rubaie, F. M. Evaluating water quality index of al Hammar Marsh, south of Iraq with the application of GIS technique. *Journal of engineering science and technology*. 2018; 13(12), 4118-4130.
56. Khudair, B. H., and Al-Musawi, N. O. Water quality assessment and total dissolved solids prediction using artificial neural network in Al-Hawizeh Marsh South of Iraq. *Journal of Engineering*. 2018; 24 (4), 147-156.
57. Kamesh, Singh, B.P., Misra, M., Verma, K.K., Singh, C.K. and Kumar, R. An emerging adsorption technology and its applicability on trees as an adsorbent for the remediation of

water pollution: a review. Ecology, Environment and Conservation. 2023; 29, (2): 627-640. DOI: [10.53550/EEC.2023.v29i02.014](https://doi.org/10.53550/EEC.2023.v29i02.014)

58. Boyd, C. E., Boyd, C. E. Carbon dioxide, pH, and alkalinity. Water Quality: An Introduction. 2020; 177-203. https://doi.org/10.1007/978-3-030-23335-8_9

UNDER PEER REVIEW