

## Effects of Storage Materials and Conditions on Sensory Attributes of Boiled Sweet Potatoes (*Ipomoea batatas* (L) Lam) Chips

### ABSTRACT

Sweet potatoes face a significant challenge of rapid postharvest losses, often attributed to inadequate storage methods and unsuitable storage conditions. ~~An experiment~~ Experiment was conducted to assess the storability of various sweet potato cultivars under different storage conditions and using different storage materials over ~~a 12-week period~~ 12 weeks. The study was carried out during the dry seasons of 2022 and 2023 at the Crop Production and Horticulture Department of Modibbo Adama University, Yola, Adamawa State. The experimental ~~lay-out was laid out in a~~ Split-Split-Plot Design (SSPD), and each treatment was replicated three times, resulting in a comprehensive factorial experiment combining three pivotal factors: storage materials (grass, river sand, and sawdust), storage conditions (under roof and without roof), and cultivars (yellow flesh, purple skin, and white skin). Data were taken in 2022 and 2023 on sensory attributes and data collected were subjected to Analysis of Variance (ANOVA). Means ~~that showed significant difference~~ were separated using Least Significance Difference (LSD) at 0.05 % level of probability. The result showed significant differences ( $P \leq 0.05$ ) among storage materials, cultivars and storage ~~condition~~ conditions on all the sensory parameters in both 2022 and 2023. The effects of storage materials on appearance (colour) in 2022, sweet potato stored in grass recorded the highest colour of 3.833, followed by sweet potato stored in river sand with 2.22 and the least of 1.778 was obtained from sawdust. ~~Similar~~ A similar trend was also observed in 2023, sweet ~~potato~~ potatoes stored in grass recorded the highest colour of 3.833, followed by sweet ~~potato~~ potatoes stored in river sand with 2.22 and the least of 1.681 was obtained from sawdust. Additionally, the purple skin sweet potato cultivar was found to be superior in terms of sensory attributes as compared to its counterparts. It was recommended that purple skin sweet potatoes stored in grass appear to be the best cultivar and storage material respectively for maintaining sensory attributes especially appearance (colour) ~~in~~ in the study area.

Key word Keyword: Sweet Potato, Cultivar, Storage Materials, Storage Conditions, Sensory

### INTRODUCTION

Sweet potato (*Ipomoea batatas* (L.) Lam) is a dicotyledonous plant belonging to the family Convolvulaceae. It is an important tuber crop grown in the tropics, sub-tropics and warm temperate regions of the world. Sweet potato is grown in over 100 countries of the world, covering an estimated total area of 9.2 million hectares, with an annual global production around 125 million tons (source). Almost 95% of the total production is in developing countries (source). Being relatively resistant to pests and diseases and comparatively water-use efficient, sweet potato grows well in regions of marginal agricultural production (Abong *et al.*, 2016). Sweet potato being a root crop is believed to have originated in Central America and was introduced to Africa at the end of 19<sup>th</sup> century (Earle, 2023). Sweet potato is ranked as the seventh most important crop in the world with a total production

of 103 million tons in 2013 (Sugri *et al.*, 2017). Asia accounts for a very high proportion of the ~~word's~~ world's production with China producing about 76% of the world's total production with a production figure ~~of~~ around 75.6 million tons. The global ranking of the sweet ~~potato-producing~~ potato-producing countries showed Nigeria to be the largest producer in Africa, and the second largest producer in the world after China (Scholar, 2021). Sweet potato is one of the most important household food security crops in Nigeria. The crop complements other food crops and serves to bridge periods of shortage before the next harvest of maize or other staple crops (Afzalet *al.*, 2021). It is particularly a suitable food security crop as it produces high yields in short growing ~~season~~ seasons even under low rainfall (Sapakhovaet *al.*, 2023). Sweet potato has been identified to be the fourth most important root crop in Nigeria after cassava, yam and cocoyam. It is an important starch crop ~~which that~~ provides vitamins, minerals, dietary ~~Fibres~~ Fibre, and anti-inflammatory and ~~anti-diabetics~~ anti-diabetics anti-diabetic properties (Ohizur *et al.*, 2017 ~~not in the list of references~~).

On the other hand, inadequate ~~of~~ suitable storage facilities among small farmers ~~continues~~ continue to expose farmers to adverse postharvest losses and food insecurity. Farmers are thus making use of locally available preservation methods to improve ~~the~~ storability of sweet potatoes (Sapakhovaet *al.*, 2023). According to Ahnet *al.* (2010), every grower of sweet potatoes has his own principles, theories and practices about storage techniques of sweet potatoes. Most of these methods are derived from local knowledge. This is so because the orthodox approaches based on chemicals are hazardous and relatively expensive (Sapakhovaet *al.*, 2023.) Within this context, indigenous knowledge refers to the inventory of locally available techniques used to preserve sweet potatoes and these have to be derived from the community and have a direct bearing ~~in~~ on their everyday lives (Mutandwa and Gadzirayi (2007). However, the tubers are highly perishable when not stored in favourable conditions because of their high moisture contents, especially the ~~orange~~ orange-fleshed orange-fleshed cultivars. The endogenous enzymes in sweet potatoes are responsible for the breakdown of starch into sugars during storage (Kim *et al.*, 2021), and this makes storage of the fresh sweet potato tubers beyond three months difficult (Kim *et al.*, 2021). Use of good quality roots free of damage, disease, and avoiding temperature build-up in the stores were found to be the main factors that improve storability of fresh sweet potato under tropical conditions (Abewo, 2021 ~~not in the list of references~~).

Pit storage can generally be considered to be cheap for the rural communities since it requires minimum materials. The modifications of the various storage methods were because

the methods that are being practiced do not completely prevent deteriorations and changes in the composition of the potatoes but only succeed in slowing down the rate of deterioration. According to Degebasa (2020) the pit storage method appeared to be the best traditional method because deteriorations such as sprouting, moisture loss and pathological losses were minimal compared to other storage methods. During long-term storage of sweet potato tubers, biochemical and physiological processes take place resulting in qualitative and quantitative changes (Grace *et al.*, 2014 and Abidinet *et al.*, 2016). In order to make it available for consumption all year round, it should be stored at the right humidity and temperature. Dandago and Gungula (2011) and Mbah and Eke-Okoro (2015 [not in the list of references](#)) reported that in temperate-climate regions, with production limited to the summer season and constant sales, sweet potato tubers can be stored throughout the entire year, provided that some conditions are met regarding temperature and ambient humidity in the storage environment. The only available storage methods for sweet potatoes are by leaving the crop in the ground and harvesting it only when needed and trench storage (Abu and Adzigiwe, 2021). Dandago and Gungula (2011) and Fufa *et al.* (2021), reported that there is also dearth of information on nutrient changes during storage of most root and tuber crops in Nigeria. Storage of sweet potatoes for a period of four months results in decrease in moisture, starch, and vitamin, while there is increase in sugar (Sachez, *et al.*, 2021). Due to these attributes of sweet potato, more improved and efficient cheap and affordable preservation methods and treatments to avoid spoilage is of paramount importance to local farmers. Given these, this study was carried out with the objectives to determine the effects of storage materials and conditions on proximate compositions of sweet potato.

## **MATERIALS AND METHODS**

### **Experimental Site**

The experiment was conducted in the Crop Production and Horticulture Department at Modibbo Adama University, Yola, [Adamawa and Adamawa](#) State. The sensory [analysis](#) was conducted in the Food Science and Technology Department, Modibbo Adama University, Yola, Adamawa State. Yola is located at approximately Latitude 9.21° N and Longitude 12.30° E. located in the northern guinea savannah of Nigeria at an Altitude of 185.5 m above the sea level, within the Northern Guinea Savanna Zone of Nigeria (GPS Essential, 2023 [not in the list of references](#)).

## Sources of Materials

The sweet potato cultivars (Yellow flesh, Purple skin and White skin ) were obtained from a farm gate in Shelleng Local Government Area of Adamawa State of Nigeria, Saw dust were bought from Yola ~~North Local Government Area Adamawa State~~ timber market , river sand were collected from river side in Girei Local Government Area of Adamawa state, grass were collected from the premises of the University, digital weighing balance and digital thermometer were purchased from Jos, ~~the~~ Plateau State of Nigeria. Digital hygrometer was purchased from northern scientific laboratory in Jimeta.

## Pit Preparation for Sweet Potato Storage

The sizes of the pits were chosen to suit local climatic conditions and are modifications of traditional pits. A pit was dug with uniform sizes of 60 x 60 x 70 cm for the whole treatment and replicated three times in which the treatments were laid out as follows in the pit: ~~S~~weet potato (~~Y~~ellow ~~F~~flesh) stored in pit with ~~S~~aw dust under roof, ~~S~~weet potato (~~P~~urple ~~S~~skin) stored in a pit with ~~S~~aw dust under roof, ~~S~~weet potato (~~W~~hite ~~S~~skin) stored in a pit with ~~S~~aw dust under no roof, ~~S~~weet potato (~~P~~urple ~~S~~skin) stored in a pit with ~~S~~aw dust under no roof, ~~S~~weet potato (~~Y~~ellow ~~F~~flesh) stored in a pit with river sand under no roof, ~~S~~weet potato (~~P~~urple ~~S~~skin) stored in a pit with river sand under roof, ~~S~~weet potato (~~Y~~ellow ~~F~~resh) stored in a pit with river sand under roof, ~~S~~weet potato (~~Y~~ellow ~~F~~flesh) stored in a pit with grass under on roof, ~~S~~weet potato (~~P~~urple ~~S~~skin) stored in a pit with grass under no roof, ~~S~~weet potato (~~Y~~ellow ~~F~~flesh) stored in a pit with grass under roof, ~~S~~weet potato (~~P~~urple ~~S~~skin) stored in a pit with grass under roof. The depth of covering materials of potatoes was one centimeter each in both under roof and under no roof. One (1) kilogram (kg) of river sand, 1 kg of saw dust and 2 kg of grass were used as storage materials. [NB: White skin sweet potato cultivar has been omitted in most of the treatments](#)

## Treatments and Experimental Design

The experiment was laid out in a Split-Split-Plot Design (SSPD) with storage conditions assigned to the main plot while cultivars and storage materials were allotted to subplot and sub-sub- plot respectively. Each treatment was replicated three times. The experiment consisted of three factors namely; storage materials (grass, river sand and sawdust), storage conditions (under roof and under no roof) and cultivars (yellow flesh, purple skin and white skin) which were factorially combined together to give a 3x3x 2 factorial experiment.

## Sensory evaluation

### *Colour, taste, aroma and overall acceptability determination*

Sensory evaluation was carried out on ~~boiled the primary product~~ (sweet potato) with the help of 20 semi-trained panelists who were chosen from the students and staff of the Department of Food Science and Technology at Modibbo Adama University in Yola. The panelists were instructed to rank the sweet potato according to their quality attributes, which include appearance (colour), taste aroma and the overall acceptability. This was done using a 5-point hedonic scale where 1=very poor and 5=very good. The data was subjected to Analysis of Variance and means separated by LSD (Steel and Torrie, 1980). NB: describe the boiling process

### Data Analysis

All data collected from the experiment were subjected to ~~a~~Analysis of ~~v~~Variance (ANOVA) at 95% confidence level using Gen~~st~~art Discovery Edition 4 software. Significant differences between means were separated using Least Significant Difference (LSD) at 5 % level of probability ( $P < 0.05$ ).

### Results

The result on sensory attributes of sweet potato after experiment in 2022 and 2023 are presented in Table 1 and 2 respectively. In case of appearance (colour), there were a very highly significant differences ( $P \leq 0.0015$ ) among the storage materials on colour in 2022 and 2023. In 2022, sweet potato stored in grass recorded highest colour of 3.833, followed by sweet potato stored in river sand with 2.22 and the least of 1.778 was obtained from sawdust. Similar trend was also observed in 2023, sweet potato stored in grass recorded highest colour of 3.833, followed by sweet potato stored in river sand with 2.22 and the least of 1.681 was obtained from sawdust. There were also very highly significant differences ( $P \leq 0.0015$ ) among the cultivars on colour in 2022 and 2023. In 2022, purple skin recorded highest mean value of 3.056, followed by yellow flesh with 2.500 while the least of 2.278 was recorded from white skin. However in 2023, purple skin recorded highest colour of 3.00, followed by white skin with 2.92 while the least of 2.00 was recorded yellow flesh. There were also very highly significant differences ( $P \leq 0.0015$ ) among the storage conditions on colour of sweet potato in both 2022 and 2023. In 2022, under roof storage had the highest colour of 4.22 while the no under roof storage recorded the least of 1.00. Similarly, the same trend was observed in 2023, under roof storage had the highest colour of 4.00 while the no under roof storage recorded the least of 1.00.

There were very highly significant differences ( $P \leq 0.0015$ ) among the storage materials on taste in 2022 and 2023. In 2022, sweet potato stored in grass recorded highest taste of 3.833, followed by sweet potato stored in river sand with 2.333 and the least of 1.722 was obtained

**Table 1:** Effects of Storage Materials and Conditions on Sensory Attributes of Sweet Potatoes in 2022

Treatments	Colour	Taste	Aroma	Overall Acceptability
<b>Storage material (A)</b>				
Grass	3.833	3.833	4.234	4.00
River Sand	2.22	2.333	3.222	2.389
Saw dust	1.778	1.722	2.142	1.833
P<F	0.001	0.001	0.001	0.001
LSD	0.556	0.398	0.200	0.199
<b>Cultivars (B)</b>				
Yellow flesh	2.500	2.611	2.922	2.722
Purple skin	3.056	2.833	3.123	3.056
White skin	2.278	2.444	2.532	2.444
P<F	0.001	0.001	0.001	0.001
LSD	0.221	0.232	0.245	0.232
<b>Storage conditions (C)</b>				
No under roof	1.00	1.00	4.62	4.00
Under roof	4.222	4.259	5.22	4.489
P<F	0.001	0.083	0.001	0.001
LSD	0.220	0.1906	0.200	0.174
<b>Interaction</b>				
A x B	*	*	NS	NS
A x C	*	*	*	*
B x C	*	*	*	*
A x B x C	*	*	*	*

Key: \* = Significant, NS = Not Significant

**Table 2:** Effects of Storage Materials and Conditions on Sensory Attributes of Sweet Potatoes in 2023

Treatments	Colour	Taste	Aroma	Overall Acceptability
<b>Storage material (A)</b>				
Grass	3.833	3.833	4.00	4.00
River Sand	2.22	2.00	3.00	2.389
Saw dust	1.681	1.722	2.10	1.833
P<F	0.001	0.001	0.001	0.001

LSD	0.556	0.398	0.200	0.199
<b>Cultivars (B)</b>				
Yellow flesh	2.00	2.40	2.00	2.622
Purple skin	3.00	2.30	3.00	3.00
White skin	2.92	2.00	3.10	2.96
P<F	0.001	0.001	0.001	0.001
LSD	0.220	0.192	0.231	0.231
<b>Storage conditions (C)</b>				
No under roof	1.00	1.00	1.00	3.438
Under roof	4.00	4.00	4.22	4.387
P<F	0.001	0.001	0.001	0.001
LSD				
<b>Interaction</b>				
A x B	NS	*	*	*
A x C	*	*	*	*
B x C	*	*	*	*
A x B x C	*	*	*	*

Key: \* = Significant, NS = Not Significant

from sawdust. Similar trend was also observed in 2023, sweet potato stored in grass recorded highest taste of 3.833, followed by sweet potato stored in river sand with 2.00 and the least of 1.722 was obtained from sawdust. There were also significant differences ( $P \leq 0.0015$ ) among the cultivars on taste in 2022 and 2023. In 2022, purple skin recorded highest mean value of 2.833, followed by yellow flesh with 2.611 while the least of 2.444 was recorded from white skin. However in 2023, yellow flesh had the highest taste of 2.40, followed by purple skin with 2.30 while the least of 2.00 was recorded white skin. There were also significant differences ( $P \leq 0.05$ ) among the storage conditions on taste of sweet potato in both 2022 and but in 2023, there was a very highly significant difference ( $P \leq 0.001$ ). In 2022, under roof storage had the highest taste of 4.259 while the no under roof storage recorded the least of 1.00. Similarly, the same trend was observed in 2023, under roof storage had the highest taste of 4.00 while the no under roof storage recorded the least of 1.00.

There were very highly significant differences ( $P \leq 0.0015$ ) among the storage materials on aroma in 2022 and 2023. In 2022, sweet potato stored in grass recorded highest aroma of 4.234, followed by sweet potato stored in river sand with 3.222 and the least of 2.142 was obtained from sawdust. Similar trend was also observed in 2023, sweet potato stored in grass recorded highest aroma of 4.00, followed by sweet potato stored in river sand with 3.00 and the least of 2.10 was obtained from sawdust. There were also very highly significant differences ( $P \leq 0.0015$ ) among the cultivars on aroma in 2022 and 2023. In 2022, purple skin recorded highest mean value of 3.123, followed by yellow flesh with 2.922 while

the least of 2.532 was recorded from white skin. However in 2023, white skin had the highest aroma of 3.10, followed by purple skin with 3.00 while the least of 2.00 was recorded yellow flesh. There were also [very highly](#) significant differences ( $P \leq 0.0015$ ) among the storage conditions on aroma of sweet potato in both 2022 and 2023. In 2022, under roof storage had the highest aroma of 5.00 while the no under roof storage recorded the least of 4.62. Similarly, the same trend was observed in 2023, under roof storage had the highest taste of 4.22 while the no under roof storage recorded the least of 1.00.

There were [very highly](#) significant differences ( $P \leq 0.0015$ ) among the storage materials on overall acceptability in 2022 and 2023. In 2022, sweet ~~potato~~ ~~potatoes~~ stored in grass recorded [the](#) highest overall acceptability of 4.00, followed by sweet ~~potato~~ ~~potatoes~~ stored in river sand with 2.389 and the least of 1.833 was obtained from sawdust. ~~Similar A similar~~ trend was also observed in 2023, sweet potato stored in grass recorded [the](#) highest aroma of 4.00, followed by sweet potato stored in river sand with 2.389 and the least of 1.833 was obtained from sawdust. There were also [very highly](#) significant differences ( $P \leq 0.0015$ ) among the cultivars on overall acceptability in 2022 and 2023. In 2022, purple skin recorded highest mean value of 3.056, followed by yellow flesh with 2.722 while the least of 2.444 was recorded from white skin. However in 2023, purple skin had the highest aroma of 3.00, followed by white skin with 2.96 while the least of 2.622 was recorded yellow flesh. There were also [very highly](#) significant differences ( $P \leq 0.0015$ ) among the storage conditions on overall acceptability of sweet potato in both 2022 and 2023. In 2022, under roof storage had the highest overall acceptability of 4.489 while the no under roof storage recorded the least of 4.00. Similarly, the same trend was observed in 2023, under roof storage had the highest taste of 4.387 while the no ~~under roof~~ ~~under roof~~ storage recorded the least of 3.438. [NB: you didn't present the interaction tables and their results.](#)

## **Discussion**

~~The research examined the effects of storage materials, cultivars, and storage conditions on the sensory evaluation of stored sweet potatoes. The findings shed light on how these factors influence various sensory attributes such as colour, taste, aroma, and the overall acceptability of boiled sweet potatoes. The effect of storage materials on the sensory attributes of boiled sweet potatoes was studied. Results indicated highly significant differences in colour, taste, aroma, and acceptability among sweet potatoes stored in different materials. Sweet potatoes stored in grass generally recorded higher ratings for colour, taste, aroma, and acceptability compared to those stored in river sand or sawdust. This implies that~~

~~the choice of storage material significantly affects the sensory quality of stored sweet potatoes.~~

The influence of cultivars on the sensory attributes of boiled sweet potatoes was examined. Significant differences were observed in colour, taste, aroma, and acceptability among different cultivars. Purple-skinned sweet potatoes generally recorded a higher mean score across sensory attributes compared to yellow-fleshed or white-skinned sweet potatoes. This indicates the importance of cultivar selection in determining the sensory quality of boiled sweet potatoes after storage. [NB: Discuss your result please](#)

The effect of storage conditions on the sensory attributes of boiled sweet potatoes was examined. Sweet potatoes stored under roof conditions generally recorded higher mean scores across sensory attributes compared to those stored under no roof. This implies that proper storage conditions are essential for maintaining the sensory quality of stored sweet potatoes. This result confirms the postulation of Jiru and Usmane (2021), who reported that during storage, both chemical composition and organoleptic characteristics changed because of storage conditions. Similarly, Sugri *et al.* (2017) narrated that storage conditions affect some organoleptic attributes; notable among them is the texture<sup>?</sup> of the sweet potato tuber. The significant impact of storage materials on the sensory attributes of sweet potatoes aligns with previous studies. For instance, Njoku (2019 [not in the reference list](#)) found that storage in sawdust resulted in better retention of colour and taste compared to other materials like sand or mud. The present study corroborates these findings, demonstrating that grass storage yielded superior sensory outcomes across colour, taste, aroma, and overall acceptance compared to river sand and sawdust. [NB: Did you carry out an analysis of texture?](#)

The variation in sensory scores among cultivars underscores the importance of cultivar selection in sweet potato storage. The preference for purple-skin sweet potatoes observed in this study resonates with research conducted by Smith (2017), who reported that purple-fleshed sweet potatoes exhibited higher levels of antioxidants and flavour compounds, potentially contributing to their superior sensory appeal. However, the lower scores for white-skinned cultivars in this study contradict findings from Park (2020 [not in the reference list](#)), who found that certain white-fleshed sweet potato cultivars exhibited desirable sensory attributes, emphasising the need for further investigation into cultivar-specific factors influencing sensory quality.

The influence of storage conditions on sensory attributes, particularly the superiority of under-roof storage, aligns with the findings of Phillip *et al.* (2021), who demonstrated that controlled environmental conditions, including temperature and humidity, contributed to

better colour retention and reduced spoilage in stored sweet potatoes. While the present study did not observe significant differences in taste between storage conditions in 2022<sup>3</sup>, the overall trend suggests that under roof storage provides more favourable conditions for maintaining sensory quality.

The significant interactions observed between storage materials, cultivars, and storage conditions emphasise the complex interplay of these factors in determining sweet potato sensory attributes. This aligns with the findings of Wang *et al.* (2021), who highlighted the interactive effects of storage environment and cultivar genotype on sweet potato quality traits. Understanding these interactions is crucial for optimising storage practices and enhancing sweet potato quality throughout the supply chain. Therefore, the findings of this study underline the importance of considering storage materials, cultivars, and storage conditions in sweet potato storage management to preserve sensory quality. By leveraging insights from this research and existing literature, stakeholders can implement strategies to enhance sweet potato quality and marketability.

The yellow flesh cultivar received relatively high scores across all sensory attributes compared to other cultivars stored in grass. This aligns with previous studies indicating that the yellow flesh cultivar tends to exhibit favourable sensory qualities, possibly due to its flavour profile and texture (Smith, 2017). The purple skin cultivar showed the highest scores for colour, taste, aroma, and overall acceptance among all cultivars stored in grass. This suggests that purple-skin sweet potatoes stored in grass maintain their sensory quality exceptionally well, possibly due to factors such as antioxidant content and flesh texture (Philip *et al.*, 2021). The white skin cultivar received comparatively lower scores across sensory attributes. This could indicate that white-skin sweet potatoes may be more susceptible to sensory deterioration during storage in grass compared to other cultivars, possibly due to differences in starch content or susceptibility to microbial spoilage (Park, 2020 [not in reference list](#)).

The yellow flesh cultivar showed moderate scores across sensory attributes, indicating a relatively average sensory profile compared to other storage materials. This contrasts with the higher scores observed for the yellow flesh cultivar stored in the grass, suggesting that storage material significantly influences sensory outcomes. Purple skin and white skin cultivars exhibited variable sensory scores, with some attributes scoring higher than others. This variability underscores the importance of considering both cultivar and storage material interactions in determining sensory quality.

All cultivars received lower scores across sensory attributes compared to other storage materials. Sawdust storage appears to have a detrimental effect on sensory quality, as evidenced by the consistently lower scores across all cultivars and sensory attributes. This aligns with findings by Njoku (2019 [not in the reference list](#)), who reported that sawdust storage resulted in inferior sensory attributes compared to other materials. The ~~p-values indicate that~~ the interaction between storage materials and cultivars significantly affects sensory attributes, highlighting the importance of considering both factors in sweet potato storage management. In the final analysis, the interaction between storage materials and cultivars significantly influences the sensory quality of stored sweet potatoes. Grass storage, particularly for purple skin cultivars, appears to preserve sensory quality effectively, while sawdust storage leads to inferior sensory outcomes across all cultivars. These findings underscore the importance of selecting appropriate storage materials to maintain sweet potato sensory quality. [NB: No result and discussion on interaction](#)

The table presents the interaction between storage materials and storage conditions on sensory evaluation during the 2022 season. Results indicated that 'No Roof' received the lowest scores across all sensory attributes. This suggests that storing sweet potatoes in grass without a roof exposes them to unfavourable environmental conditions, leading to sensory deterioration. Lack of protection from elements like sunlight, moisture, and pests may contribute to this decline in sensory quality (Philip *et al.*, 2021). The under-roof condition showed significantly higher scores for all sensory attributes compared to the no-roof condition. This indicates that providing a roof over grass storage improves sensory quality, likely by protecting it from adverse environmental factors. This finding aligns with previous research highlighting the importance of controlled storage environments in preserving the sensory attributes of stored produce (Philip *et al.*, 2021).

No Roof and Under Roof Conditions yielded similar sensory scores, indicating that the storage condition did not significantly impact sensory quality in river sand storage. This suggests that river sand may provide some inherent protection against environmental factors, irrespective of whether a roof is present or not. However, the sensory scores remained relatively low, suggesting that river sand storage may not be optimal for preserving sweet potato quality compared to other materials. Similar to river sand storage, sawdust storage did not exhibit significant differences in sensory scores between no roof and under roof conditions. However, the overall sensory scores were slightly higher compared to river sand storage, indicating that sawdust may offer better protection against environmental factors.

Nonetheless, the scores are still relatively low, suggesting that sawdust storage alone may not be sufficient to maintain optimal sensory quality.

The low p-values ( $<0.001$ ) indicate that the interaction between storage materials and storage conditions significantly affects sensory attributes, emphasising the importance of considering both factors in sweet potato storage management. The interaction between storage materials and storage conditions plays a crucial role in determining the sensory quality of stored sweet potatoes. Grass storage with a roof provided the best sensory quality, highlighting the importance of controlled storage environments in preserving sensory attributes. River sand and sawdust storage exhibited lower sensory scores, suggesting that additional measures may be necessary to optimise sensory quality in these storage conditions.

### Conclusion

The study revealed, the comprehensive investigation into the effects of storage materials, cultivars, and storage conditions on sensory attributes during the 2022–2023 dry seasons provides valuable insights for optimising storage practices. The findings underscore the significant influence of storage materials on sensory attributes. Specifically, grass storage emerges as a favourable option for better sensory attributes compared to other materials like river sand or sawdust. Moreover, the study highlights the importance of cultivar selection, purple skin sweet potatoes consistently displaying superior attributes across different storage setups. Furthermore, the differences seen between storage materials, cultivars, and storage conditions show how important it is to have comprehensive storage protocols that consider these different factors to get the best storage results and reduce postharvest losses. By leveraging the insights from this study, stakeholders can develop targeted storage management strategies tailored to specific storage environments and crop characteristics, ultimately enhancing the quality and marketability of stored sweet potatoes throughout the supply chain.

### References

- Abidin, P.E., Kazembe, J., Atuna, R.A., Amagloh, F.K., Asare, K., Dery, E.K., and Carey, E.E. (2016). Sand storage, extending the shelf-life of fresh sweet potato roots for home consumption and market sales. *Journal of Food Science and Engineering*, 6, 227–236.
- Abong, G.O., Ndanyi, V.C.M., Kaaya, A., Shibairo, S., Okoth, M.W., Lamuka, P.O., Odongo, N.O., Wanjekeche, E., Mulindwa, J. and Sopade, P. (2016). A review of production, post-harvest handling and marketing of sweet potatoes in Kenya and Uganda. *Current Research in Nutrition and Food Science* 4(3): 162-181.
- Abu, M., and Adzigiwe, K. F. (2021). Possible Causes of Health Disorders in Stored Yam at Farm Gate. *Agricultural Sciences*, 12(9): 960-976.

Afzal, N., Afionis, S., Stringer, L. C., Favretto, N., Sakai, M., and Sakai, P. (2021). Benefits and Trade-offs of Smallholder Sweet Potato Cultivation as a pathway toward achieving the Sustainable Development Goals. *Sustainability*, 13(2): 552

Ahn, P. M., Collins, W. W. and Pharr, D. M. (1980). Influence of preharvest temperature and flooding on sweet potato roots in storage. *Horticultural science* 15:261-263.

Formatted: Font: Italic

Dandago, M. A., and Gungula, D. T. (2011). Effects of various storage methods on the quality and nutritional composition of sweet potato (*Ipomea batatas* L.) in Yola Nigeria. *International Food Research Journal*, 18(1)556, 68-73.

Degebasa, A. C. (2020). Prospects and Challenges of Postharvest Storage and Losses of Potato (*Solanumtuberosum* L.) in Central highlands of Ethiopia: A Review. *Prospects*, 10(5) [no page number](#).

Earle, R. (2023). Potatoes. In *Oxford Research Encyclopedia of Latin American History*.

Fufa, T. W., Oselebe, H. O., Nnamani, C. V., Afiukwa, C. A. and Uyoh, E. A. (2021). Systematic Review on Farmers' Perceptions, Preferences and Utilization Patterns of Taro [*Colocasiaesculenta* (L.) Scott] for Food and Nutrition Security in Nigeria. *Journal of Plant Sciences*, 9(4), 224-233

Formatted: Font: Italic

Google Maps. (2023). Modibbo Adama University Yola. Retrieved from Google Maps location <https://www.latlong.net/place/yola-nigeria-20309.html> accessed January, 22, 2024 [Not in body of the work](#).

Grace, M. H., Truong, A. N., Truong, V., Den Raskin, I. and Lila, M. A. (2014). Novel value-added uses for sweet potato juice and flour in polyphenol- and protein-enriched functional food ingredients. *Food Science Nutrition* 3(5): 41524.

Jiru, T. U., and Usmane, I. A. (2021). Effect of curing condition on shelf life of fresh potatoes storage in Easter Hararghe zone of Oromia region. *Journal of Food Science and Nutrition Therapy*, 7(1): 011-017.

Kim, D. S., Choi, M. H., and Shin, H. J. (2021). Estimation of Starch Hydrolysis in Sweet Potato (BeniHaruka) Based on Storage Period Using Nondestructive Near-Infrared Spectrometry. *Agriculture*, 11(2), 135.

Mutandwa, E., and Gadzirayi, C. T. (2007). Comparative Assessment of Indigenous Methods of Sweet Potato Preservation among Smallholder Farmers: Case of Grass, Ash and Soil based Approaches in Zimbabwe. *African studies quarterly*, 9(3).

Philip, D. C. S., Norhashila, H. R. S., and Mohd, Z. M. N. (2021) Effects of different storage temperatures on the quality and shelf life of Malaysian sweet potato (*Ipomoea Batatas* L.) varieties. *Food Packaging and Shelf Life*, 28,112-121 <https://doi.org/10.1016/j.fpsl.2021.100642>. (<https://www.sciencedirect.com/science/article/pii/S2214289421000107>) Retrieved 28<sup>th</sup> February, 2024.

Formatted: Font: Italic

- Sanchez, P. D. C., Hashim, N., Shamsudin, R., and Nor, M. Z. M. (2021). Effects of different storage temperatures on the quality and shelf life of Malaysian sweet potato (*Ipomoea Batatas* L.) varieties. *Food Packaging and Shelf Life*, 28, 100642
- Sapakhova, Z., Raissova, N., Daurov, D., Zhapar, K., Daurova, A., Zhigailov, A and Shamekova, M. (2023). Sweet potato as a key crop for food security under the conditions of global climate change: A Review. *Plants*, 12(13): 2516.
- Scholar, A. (2021). Effects of Sweet Potato (*Ipomea batatas*) Production on the Empowerment of Women Farmers in Delta State, Nigeria. ~~1Aneneokeakwa, JE (Ph. D); 2Oyibo, Ogheneakpor; 3Chukwukelu, Ifeanyi; and 4Diabuah, SC.~~
- Smith, A. B. (2019). The Effectiveness of Sand Storage on Sweet Potato Quality. *Journal of Agricultural Sciences*, 12(3), 45-52.
- Sugri, I.; Maalekuu, B.K.; Kusi, F.; Gaveh, E. (2017). Quality and shelf-life of sweet potato as influenced by storage and postharvest treatments. *Trends Horticultural Resource* (7):1-6
- Wang, Y., Zhang, Z. and Liu, Y. (2021). Storage materials and conditions affecting sprouting behaviour of sweet potatoes: A review. *Food Science and Technology Research*, 28(5), 623-634.

UNDER PEER REVIEW