

Case report

Renal and hemorrhagic complications following bee sting envenomation: a case report from the hospital of Zinder

ABSTRACT

Renal and hemorrhagic complications following envenomation by bee stings are rare. A 44-year-old patient was admitted to the infectious and tropical diseases department of the Zinder national hospital with cervicofacial edema following bee stings. Respiratory symptoms included dyspnea and 91% desaturation on room air. The patient was put on oxygen, doxycycline and prednisone. The clinical picture was complicated by a worsening of the patient's condition due to the appearance of haematemesis, and a deterioration in renal function. A renal biopsy revealed acute tubulointerstitial nephropathy. He was discharged after 35 days. We report a rare case of bee sting complicated by hemorrhagic syndrome and renal involvement.

Keywords: bees, envenomation, kidney complications, haemorrhages, Zinder, Niger Republic

1. INTRODUCTION

Renal and hemorrhagic complications following bee stings with envenomation are most commonly observed as a result of venom inoculation[1,2]. They constitute a medical emergency, necessitating prompt intervention. Bee venom inoculation can lead to pain, fever, swelling, and multi-visceral involvement in affected individuals. The injected venoms are harmful either due to their quantity (multiple stings) or because the stung individual is allergic[3,4]. Bee stings can induce various clinical manifestations that can jeopardize the patient's life. Localized pain, tissue necrosis, and anaphylactic reactions followed by wasp stings are well-documented[5]. Besides these, they can produce systemic reactions and organ dysfunction[1,2]. Among these, rhabdomyolysis has been identified as one of the main mechanisms. Mellitin is believed to be the primary molecule responsible for rhabdomyolysis, as well as hemolysis[6]. Phospholipase A2, to a lesser extent, may play a role in the occurrence of rhabdomyolysis[2]. Acute renal failure results from tubular obstruction caused by myoglobin and the direct nephrotoxic effect of the venom[3]. Different cases may present varying severity levels, including mild cases (local edema with acute pain and skin reaction, sometimes accompanied by fever), moderate cases (syncope with dizziness, nausea, severe headaches, palpitations, skin reaction, and allergic reactions in certain organs), and fatal cases that can occur following anaphylactic shock [1–3].

Regarding hymenopterans, particularly bee venom, the presence of histamine, a polypeptide called apitoxin, and riboflavin have been identified. Several properties of bee venom have been elucidated, such as its hemolytic action, decreased blood coagulability, hormonal balance alterations, histamine effects, and the potency of apitoxin responsible for allergic reactions. Feldberg and Kellaway also demonstrated in 1937 that bee venom causes a significant release of histamine in the lungs and perfused livers of guinea pigs[6,7].

2. METHODS

This study examines a case of renal and hemorrhagic complications following bee stings at the Department of Infectious and Tropical Diseases of the National Hospital of Zinder. The study involved analyzing the patient's medical record.

3. CASE PRESENTATION

The patient is a 44-year-old individual admitted to the Department of Infectious and Tropical Diseases of the National Hospital of Zinder for cervicofacial edema following bee stings. Upon interrogation, fever and vomiting were reported after the sting. Physical examination revealed multiple painful sting sites with embedded stingers, cervicofacial edema, and pale mucous membranes and conjunctivae.



Figure 1 Patient stung by bees

Upon arrival, the farmer is found to be covered with numerous stings on the face, skull, neck, upper limbs, and thorax. The most affected areas are the face, neck, and thorax. In the impossibility of counting each individual sting, the Wallace's rule is applied, which divides the body surface into proportional zones:

- 9% for the head and neck
- 9% for each upper limb
- 18% for the anterior trunk
- 18% for the posterior trunk
- 18% for each lower limb
- 1% for the genitals

Based on this estimation, we can calculate that the total surface area affected by stings for this farmer is 27% (head/face = 9% and anterior trunk = 18%). According to the formula used to calculate body surface area taking into account height (in cm) and weight (in kg):

$X = 1 + (\text{height} + \text{weight} - 160) / 100$ Patient constants: Weight=83 kg, Height=178 cm. Area=X/100

This amounts to 20,100 cm², with 27% receiving at least one sting per cm², the total number of stings can thus be estimated between 6000 – 7000 or 70 stings per Kg of body weight. This number is comparable to that of bees in a hive (about 50,000 insects). It corresponds to an inoculation of 2.1 to 4.9 mg of venom in total (one sting releases 0.03-0.07 mg of venom and the reservoir contains 3-4 mg of liquid).

The patient was fully conscious, with a Glasgow score of 15/15. Additionally, he had a fever with a temperature of 38.6°C, his heart rate was within normal range (97 beats/min), SPO₂ at 97% in ambient air, and his blood pressure (BP) was (143/70 mmHg). He received paracetamol intravenously at a dose of 60 mg/kg per day, divided into four doses, Solumedrol 120mg in 250cc of solution, Saline serum 40mg/kg/d, Ceftriaxone 50 mg/kg/d.

The hematological assessment showed leukocytosis at 42,900 cells/mm³, anemia at 9.6 g/dl, and thrombocytopenia at 126,000 cells/mm³. The biochemistry revealed a blood glucose level of 5.63 mmol/l, uremia at 3.30 mmol/l, and a creatinine level at 82.54 μmol/l.

The hemostasis assessment included a prothrombin time of 84.9%, an activated partial thromboplastin time of 30 sec, and C-reactive protein at 384 g/l. SRV serology and HbS antigenemia were negative. Transaminases and creatine kinase could not be performed due to technical limitations.

On the second day, the patient was dyspneic and desaturated to 91% in ambient air. Oxygen, doxycycline, and prednisone were administered.

On the third day, the clinical picture was complicated by worsening patient's condition with the appearance of hematemesis and sleep disturbances as well as deterioration of renal function, uremia at 49.97 mmol/l, creatinine at 1265.63 μmol/l (GFR at 4.22 ml/min). On urine dipstick, proteinuria at one cross and hematuria at 4 crosses were noted. Faced with this severe uremia, the indication for emergency dialysis was raised. Renal biopsy revealed acute

tubulointerstitial nephropathy. The patient's condition improved after four dialysis sessions but he developed a catheter infection during hospitalization which required catheter removal. Culture of the catheter tip isolated *Staphylococcus aureus* sensitive to vancomycin. The patient was treated with vancomycin and gentamicin.

The onset of hematemesis and sleep disturbances, along with deterioration of renal function, uremia at 49.97 mmol/l, creatinine at 1265.63 $\mu\text{mol/l}$ (GFR at 4.22 ml/min), and findings of proteinuria at one cross and hematuria at 4 crosses on urine dipstick prompted the indication for emergency dialysis due to severe uremia. Renal biopsy revealed acute tubulointerstitial nephropathy. The patient's condition improved after four dialysis sessions, but during hospitalization, he developed a catheter infection necessitating catheter removal. Culture of the catheter tip isolated *Staphylococcus aureus* sensitive to vancomycin. The patient was treated with vancomycin and gentamicin.

On the tenth day, the patient experienced a low-grade fever (38.5°C), moderate hypoxia with SPO₂ at 93% in ambient air, and spontaneous hematemesis. Hematological assessment revealed anemia at 6.7 g/dl, requiring a blood transfusion.

Treatment involved administering oxygen at 5 l/min, followed by the transfusion of 2 units of packed red blood cells (450 cc each) matched for ABO and RhD using delta Hb, along with gastric dressing with Maalox 2 sachets three times a day.

On the thirteenth day, follow-up blood tests showed normalization of various parameters.

By the twenty-fourth day, esophagogastroduodenoscopy revealed pangastropathy. A treatment protocol was initiated, including ampicillin 2g IV twice daily, metronidazole 500 mg three times a day, gentamicin 160mg IM, intravenous Gaspral 2 vials as a bolus followed by 1 vial twice daily, saline serum 500mg twice daily, and Perfalgan 1g.

On the thirty-second day, hematological and renal function tests were all normal. The patient was discharged cured after 35 days.

3. DISCUSSION

Venomous bee stings are rare but often serious due to the occurrence of various complications[8]. According to literature data, Africanized bee species are the most aggressive[9]. Bee stings are considered a public health problem in Latin America, where they are frequent and responsible for a significant number of deaths[10]. Even in Africa, cases of bee stings have been reported, notably in Morocco. The lethal threshold generally ranges between 300 and 500 stings; however, in some cases, simultaneous stings can be deadly. In the United States, bee stings are the second leading cause of anaphylactic reactions after penicillin[11]. The extreme toxicity of massive bee envenomation is due to the direct toxic effect of large quantities of injected venom. Bee venom is a complex combination of enzymes and amines. Mellitin, a component that initiates pain, constitutes 50% of the venom's weight and acts in conjunction with phospholipase A2 as a cytolytic agent. Hyaluronidase increases capillary permeability and facilitates toxin diffusion[12,13].

Immediate reactions following numerous stings typically include localized pain, sweating, and erythema at the sting site. Early systemic signs include fatigue, confusion, nausea, vomiting, and diarrhea. Within 24 hours, hemolysis, hemoglobinuria, rhabdomyolysis, and hepatic injury may develop[7]. Neurological and cardiac involvement is much rarer[14]. Acute renal involvement is often linked to the direct toxicity of the venom[3,15]. Tissue necrosis appears to result from the cytolytic effects of three venom components, namely phospholipase A2, hyaluronidase, and serine proteases[11]. A hemorrhagic syndrome was observed 24 hours after admission. Hemostasis and coagulation disorders have been reported by Przybilla and Ruëff[9]. Rhabdomyolysis, which can lead to acute renal failure, is a complication of massive envenomations[3]. Creatine phosphokinase, lactate dehydrogenase, and aspartate aminotransferase levels would be elevated[16].

In this case, therefore, muscle enzyme assays could not be performed. Nevertheless, given the clinical picture (myalgia, dark urine), it is highly likely that the patient suffered from massive rhabdomyolysis, complicated by renal failure. Although renal involvement may go unnoticed[14], cases of renal failure requiring dialysis have been reported[1,2]. The location of the stings is an important prognostic factor, particularly stings to the head and neck as well as those in the pharynx and larynx, which can lead to death due to airway obstruction[17]. Ocular lesions from bee stings can occur[13,18], but this patient did not present with direct ocular injuries. Urgent anti-anaphylactic measures should be instituted, such as fluid resuscitation, corticosteroid therapy, and subcutaneous administration of adrenaline, due to the similarity of clinical manifestations with acute anaphylactic reactions[19]. Early management is essential, with prompt patient admission, massive fluid resuscitation, high-dose corticosteroid therapy, and removal of stingers. Local treatment involves gentle scraping to remove stingers, followed by local disinfection with an antiseptic such as hydrogen peroxide or bleach[6]. The use of analgesics such as paracetamol is recommended. The prognosis of envenomation is related to venom plasma concentrations[20], and in severe cases, early plasmapheresis may be useful to reduce circulating venom concentrations and eliminate venom-mediated mediators[19].

This patient experienced a massive envenomation resulting in multi-organ failure, particularly acute renal failure. Furthermore, there was a satisfactory dynamic in the management of the patient despite the absence of a specific protocol for managing envenomations in general and bee stings in particular. Additionally, specific antivenom immunotherapy is not yet available.

4. CONCLUSION

Bee stings with envenomations are serious and can lead to various complications that can result in death. The prognosis depends on the severity of the stings. Adequate management is necessary.

CONSENT

All authors declare that written informed consent was obtained from the patient for publication of this case report and accompanying images.

ETHICAL APPROVAL (WHEREEVER APPLICABLE)

This study was conducted in accordance with ethical principles and received approval from the Institutional Review Board (IRB) of Faculty of Medical Sciences University of Zinder (FSS-UAS), Niger. The IRB reviewed the research plan, ensuring that it adhered to ethical standards and guidelines for conducting research involving human subjects. The approval from the IRB signifies compliance with the principles of confidentiality, informed consent, and the protection of participants' rights.

REFERENCES

1. Chowdhury FR, Bari MS, Shafi AM, Ruhan AM, Hossain ME, Chowdhury S, et al. Acute Kidney Injury Following Rhabdomyolysis due to Multiple Wasp Stings (*Vespa affinis*). *Asia Pacific Journal of Medical Toxicology*. 1 mars 2014;3(1):41-3.
2. Deshpande PR, Farooq AK, Bairy M, Prabhu RA. Acute Renal Failure and/or Rhabdomyolysis due to Multiple Bee Stings: A Retrospective Study. *N Am J Med Sci*. mars 2013;5(3):235-9.
3. Almeida RAM de B, Olivo TET, Mendes RP, Barraviera SRCS, Souza L do R, Martins JG, et al. Africanized honeybee stings: how to treat them. *Rev Soc Bras Med Trop*. déc 2011;44:755-61.
4. Berdai MA, Labib S, El Balbal M, Harandou M. Massive stings by a swarm of bees in a child. *Pan African Medical Journal*. 2011;10.
5. Sullerot I, Birnbaum J, Girodet E. Massive envenomation syndrome about a clinical case of multiple bites discussed by the SFA "insects" group. *French Journal of Allergology*. 2013;53(1):50-2.
6. Schumacher MJ, Tveten MS, Egen NB. Rate and quantity of delivery of venom from honeybee stings. *J Allergy Clin Immunol*. mai 1994;93(5):831-5.
7. Betten DP, Richardson WH, Tong TC, Clark RF. Massive honey bee envenomation-induced rhabdomyolysis in an adolescent. *Pediatrics*. janv 2006;117(1):231-5.
8. LoVecchio F, Cannon RD, Algier J, Ruha AM, Curry SC, Wallace KL, et al. Bee swarmings in children. *The American Journal of Emergency Medicine*. 1 oct 2007;25(8):931-3.
9. Przybilla B, Ruëff F. Insect Stings. *Dtsch Arztebl Int*. mars 2012;109(13):238-48.
10. Prado M, Quirós D, Lomonte B. Mortality due to Hymenoptera stings in Costa Rica, 1985-2006. *Rev Panam Salud Publica*. mai 2009;25(5):389-93.
11. Golden DBK, Moffitt J, Nicklas RA, Freeman T, Graft DF, Reisman RE, et al. Stinging insect hypersensitivity: A practice parameter update 2011. *Journal of Allergy and Clinical Immunology*. 1 avr 2011;127(4):852-854.e23.
12. Mass envenomations by honey bees and wasps - PubMed [Internet]. [cité 6 janv 2024]. Available on: <https://pubmed.ncbi.nlm.nih.gov/10344177/>
13. Abayi DM, Mengome EM. Ocular lesions by bee sting: about a case. *French Journal of Ophthalmology*. 2021;44(4):e191-3.
14. BOURGAIN C, PAUTI MD, FILLASTRE JP, GODIN M, FRANCOIS A, LEROY JP, et al. Mass envenomation after African bee stings. *Press med (1983)*. 1998;27(22):1099-101.
15. Azad C, Parmar VR, Jat KR. Unusual fatal multiple-organ dysfunction and pancreatitis induced by a single wasp sting. *J Venom Anim Toxins incl Trop Dis*. 2011;17:108-10.
16. Guimarães JV, Costa RS, Machado BH, Reis MA dos. Cardiovascular profile after intravenous injection of Africanized bee venom in awake rats. *Rev Inst Med trop S Paulo*. févr 2004;46:55-8.
17. Shah D, Tsang TK. Bee sting dysphagia. *Ann Intern Med*. 1 août 1998;129(3):253.
18. [Ocular lesions after bee sting of the cornea. A case report] - PubMed [Internet]. [cité 6 janv 2024]. Disponible sur: <https://pubmed.ncbi.nlm.nih.gov/19769859/>

19. Survival after massive (>2000) Africanized honeybee stings - PubMed [Internet]. [cité 6 janv 2024]. Disponible sur: <https://pubmed.ncbi.nlm.nih.gov/9570180/>
20. França FO, Benvenuti LA, Fan HW, Dos Santos DR, Hain SH, Picchi-Martins FR, et al. Severe and fatal mass attacks by « killer » bees (Africanized honey bees--*Apis mellifera scutellata*) in Brazil: clinicopathological studies with measurement of serum venom concentrations. *Q J Med.* mai 1994;87(5):269-782.

UNDER PEER REVIEW