

# Effect of planting methods and weed control treatments on weed dynamics, yield and economics of green gram (*Vigna radiata* L.)

## ABSTRACT

An experiment was conducted at the Agronomy Research Farm, College of Agriculture, Gwalior (M.P.), during the 2019 *kharif* season to investigate the effects of different planting methods and weed control treatments on the weeds, growth, yield, and economics of green gram. The field experiment followed a split-plot design with three planting methods as main plot treatments (broadcasting on a flat bed, line sowing on a flat bed, and ridge and furrow) and five weed management practices as sub-plot treatments (Pendimethalin 1000 g/ha, Diclosulam 26 g/ha, Imazethapyr 75 g/ha, hand weeding at 20 and 40 DAS and weedy check), replicated thrice. The experimental field was primarily infested with narrow-leaved weeds such as *Cyperus rotundus*, *Cynodondactylon*, *Echinochloa spp.*, and *Dactyloctenium aegyptium*, and broad-leaved weeds like *Digera arvensis*, *Celosia argentea*, *Commelinabenghalensis*, and *Phyllanthus niruri*. Among the herbicidal treatments, Imazethapyr at 75 g/ha effectively controlled both narrow-leaved and broad-leaved weeds, resulting in the lowest weed index. The ridge and furrow planting method recorded the highest values in growth parameters, yield attributes, and seed yield, proving to be more remunerative compared to other methods. These findings indicate that the combination of ridge and furrow planting with effective weed management, particularly using Imazethapyr, optimizes green gram productivity and economic returns.

**Keywords:** Green gram, Herbicides, Imazethapyr, Planting method, Ridge and furrow, Weeds

## 1. INTRODUCTION

Pulses serve as the primary protein source for India's vegetarian populace and stand as a crucial crop group within the nation's agricultural sector. They not only supply sustenance and forage but also enhance soil fertility and its physical attributes[1]. India holds the distinction of being both the largest consumer and producer of pulses, with its production accounting for 24% globally and cultivated area covering 34% [2]. Cultivation of pulses is adaptable to diverse climatic and soil conditions, playing a pivotal role in ecological balance by facilitating soil phosphorus release and atmospheric nitrogen

fixation, thus bolstering soil fertility and promoting sustainability across various farming systems[3]. Furthermore, pulses can serve multiple purposes, functioning as seeds, fodder, or green manure crops.

Green gram (*Vigna radiata* L.) stands out as a significant and widely cultivated pulse crop in India, particularly thriving in arid and semi-arid regions[4]. Locally referred to as "moong," it boasts a nutritional profile comprising approximately 25% protein, 1.3% fat, 3.5% minerals, 4.1% fiber, and 56.7% carbohydrates[5]. Despite its pivotal role in the daily diet, the average productivity of green gram remains notably low in India. Green gram (*Vigna radiata* L.) holds the third position in India in terms of both production, totaling 1.52 million tons, and cultivated area, covering 3.77 million hectares, following chickpea and pigeon pea[6]. Despite its significant importance in the diet, green gram exhibits low productivity in India. The farming community tends to favor spring season cultivation of green gram, finding it more favorable. Conversely, kharif season cultivation is less embraced due to heightened vulnerability to excessive rainfall, insect pests, and diseases. Spring season cultivation of green gram is deemed superior as it ensures a more reliable crop, with fewer incidences of insect pest attacks and no risk from rainfall compared to the rainy season crop[7].

Several factors contribute to the low yield of green gram, including improper planting methods, limited knowledge regarding herbicide application, failure to adhere to appropriate sowing dates, imbalanced fertilizer use, and inadequate pest control measures[8]. Among these, planting pattern and timing of sowing are pivotal. Planting pattern significantly impacts crop yield and growth by influencing moisture utilization and radiation interception[9]. While broadcasting remains a common method of sowing, it is recognized as a major limitation in yield and growth. Optimal production is achieved through line sowing in rows, with raised bed planting methods effectively reducing weed populations and maximizing control efficiency[10]. Cross sowing is also practiced to enhance yield. Various planting patterns significantly affect growth and yield attributes such as plant height, dry matter accumulation, number of branches per plant, pods per plant, seeds per plant, stover, and seed yield [11].

Weed infestation also poses a significant challenge, accounting for yield losses ranging from 50 to 90%, particularly during spring season cultivation[12]. Studies indicate that if weed infestation remains unchecked beyond 20 days after sowing (DAS), it can lead to severe yield reductions of up to 38% or more[13,14]. Losses due to uncontrolled weed growth were reported at 95% in wet seasons and 77% in dry seasons [15]. In more developed agricultural systems, herbicides have largely supplanted

mechanical weed control methods[16,17]. However, labor shortages during weeding periods lead to severe field infestations, rendering mechanical weeding ineffective, laborious, and costly[18,19]. Consequently, chemical weed control emerges as a viable and cost-effective alternative for this crop[20,21]. Employing effective herbicides at the appropriate rates can serve as an efficient weed control measure, potentially replacing conventional methods[22,23]. Minimizing weed growth during the crop-weed competition period can result in crop yields comparable to those of weed-free crops[24,25]. Hence, it is imperative to control weeds through any means during crop-weed competition. This paper aims to investigate the effects of planting methods and various weed control practices on the growth and yield of green gram.

## **2. MATERIALS AND METHODS**

The experiment was carried out at the Agronomy Research Farm, College of Agriculture, Gwalior (M. P.), during the 2019 *kharif* season. The experimental site, situated at 26°13' North latitude and 78°14' East longitude, stands at an elevation of 206 meters above mean sea level. It lies in the Northern tract of M.P. and experiences a subtropical climate characterized by extreme temperatures, reaching up to 48°C in summer and dropping to 4.1°C in winter. Annual rainfall typically ranges between 750 to 800 mm, primarily falling from the end of June to the end of September, with occasional showers in the winter months. Weather conditions remained normal throughout the crop season, with average maximum and minimum temperatures of 35.2°C and 24.5°C, respectively. The total rainfall received during the crop-growing period from July to October 2019 was 907.7 mm, although it was noted to be scanty and unevenly distributed.

The soil at the experimental site is sandy clay loam with a pH of 7.57. It has medium levels of organic carbon (0.42%), available nitrogen (183.50 kg/ha), phosphorus (14.48 kg/ha), and potassium (223 kg/ha).

The experiment utilized a split-plot design replicated thrice, with three planting methods as main plot treatments (broadcasting in flat bed, line sowing in flat bed, ridge and furrow) and five weed management practices as sub-plot treatments (pendimethalin 1000 g/ha, diclosulam 26 g/ha, imazethapyr 75 g/ha, hand weeding at 20 and 40 days after sowing, and a weedy check). Each experimental plot measured 5 m x 3.60 m.

The green gram variety employed in the experiment was TJM 3, with a recommended seed rate of 20 kg/ha. Seeds were manually sown in the field on July 24, 2019. After proper field preparation and layout, fertilizers were applied at the rate of 20 kg/ha of nitrogen, 50 kg/ha of P<sub>2</sub>O<sub>5</sub>, and 20 kg/ha of

K<sub>2</sub>O, with the entire quantity applied as basal dose to the crop. Urea, single super phosphate, and muriate of potash served as the sources of nitrogen, phosphorus, and potassium, respectively. Pre-emergence application of pendimethalin 1000 g/ha and diclosulam 26 g/ha was conducted within 24 hours after sowing, while post-emergence application of imazethapyr 75 g/ha was performed at 19 days after sowing to control associated weeds. Hand weeding was carried out at 20 and 40 days after sowing using khurpi. The weeds in the weedy check treatment were left uncontrolled and allowed to grow alongside the crop until harvest. Irrigation was applied as per the crop's requirement. Plant height was recorded for five plants per plot in centimeters from the ground level to the tip of the growing point using a scale. Weed species were randomly counted using one square meter quadrates from each plot. The weed index, expressed as a percentage, was calculated at harvest according to the formula by [26].

$$\text{Weed index (\%)} = \frac{(X - Y) \times 100}{X}$$

Where,

X = Yield from maximum weed free plot

Y = Yield from other treated plot

To determine the number of pods per plant, all pods were collected from three tagged plants, carefully removed by hand, and the seeds were separated from the straw and counted to obtain an average. After threshing, the seed and stover were separated and weighed per plot, and the seed yield per hectare in kilograms was calculated by multiplying with the conversion factor. Harvest index was calculated using the following formula:

$$\text{Harvest Index (\%)} = \frac{\text{Economic yield (seed)}}{\text{Biological yield (seed+straw)}} \times 100$$

### 3. RESULTS AND DISCUSSION

#### 3.1 Weed flora

The experimental field was entirely invaded with mixed weed flora consisting of narrow and broad-leaved weeds. Among the total weeds, narrow-leaved weeds were more prominent than broad-leaved weeds. Major weeds observed in the experimental field were *Cyperus rotundus*, *Cynodondactylon*, *Echinochloa Spp.*, *Dactylacteniumaegyptium* among narrow-leaved weeds while *Digera arvensis*, *Celosia argentea*, *Commelinabenghalensis*, *Phyllanthus niruri* were the most common in broad-leaved weeds. Similar findings were reported by [27, 28].

#### 3.2 Weed density (no./m<sup>2</sup>)

There was no significant variation observed in the density of all weed species at 40 days after sowing (DAS) across different planting methods (**Table 1**). In contrast, significant differences were observed in weed density among various weed control treatments. The population of weeds was notably higher in the weedy check treatment compared to all herbicidal treatments, including two hand-weeding sessions at 20 and 40 DAS. This finding underscores the importance of implementing effective weed control measures to mitigate weed competition and minimize yield losses in green gram cultivation. Among the herbicidal treatments, Imazethapyr 75 g/ha demonstrated the highest efficacy in reducing the population of both narrow and broad-leaved weeds, with results similar to Diclosulam 26 g/ha and Pendimethalin 1000 g/ha. These findings are consistent with previous research highlighting the effectiveness of these herbicides in weed suppression in various crops [29]. However, it is essential to consider factors such as herbicide resistance and environmental impact when selecting herbicidal weed control strategies [30]. Interestingly, complete control of all weeds was achieved with two hand-weeding sessions at 20 and 40 DAS. The hand-weeding method proved highly effective in reducing weed populations to negligible levels. While hand weeding can be labor-intensive and costly, particularly in large-scale agricultural operations, these results highlight its efficacy as a weed management option, especially in situations where herbicide use may be limited or undesirable [31,32]. The non-significant interaction effect of planting methods and herbicides on weed population further supports the results that planting methods did not significantly interact with the efficacy of herbicidal weed control treatments in influencing weed density.

UNDER PEER REVIEW




(PxW)								
-------	--	--	--	--	--	--	--	--

UNDER PEER REVIEW

### 3.3 Weed index (%)

The weed index varied significantly with different planting methods (**Figure 1**). Crops planted using the ridge and furrow method had the lowest weed index at 14.01%, outperforming other sowing techniques. Line sowing on flat beds also had a relatively low weed index of 20.52%, compared to broadcasting on flat beds. Among the weed control treatments, the weed index was highest at 47.54% in the untreated (weedy check) plots, significantly higher than in all herbicide-treated plots, including those with two hand weeding sessions at 20 and 40 days after sowing (DAS). Among the herbicides, Imazethapyr at 75 g/ha was the most effective, with a weed index of 18.38%, closely followed by Diclosulam at 26 g/ha and Pendimethalin at 1000 g/ha. The interaction between planting methods and herbicides on the weed index was found to be non-significant.




**Figure 1. Effect of planting methods and weed control methods on weed index (%) in green gram**

### 3.4 Herbicidal efficiency index (%)

Crops sown using the broadcasting method on a flat bed exhibited the highest herbicidal efficiency index at 2.196%, surpassing all other treatments (**Figure 2**). Line sowing on a flat bed also showed a relatively high herbicidal efficiency index of 1.362%, compared to the 0.829% observed with the ridge and furrow method. This indicated that the broadcasting method may provide more uniform herbicide distribution and better coverage, resulting in more effective weed control. The even dispersion of seeds in broadcasting likely enhances herbicide contact with the weed flora, thereby increasing the

herbicidal efficiency index. Various weed control treatments demonstrated no significant variation in herbicidal efficiency index. Additionally, the interaction between planting methods and herbicides on the herbicidal efficiency index was found to be non-significant.



**Figure 2. Effect of planting methods and weed control methods on herbicidal efficiency index (%) in green gram**

### 3.5 Growth parameters and yield attributing trait

The different planting methods significantly influenced growth parameters such as plant height, number of branches, number of root nodules, and yield traits like pods per plant at 40 DAS (**Table 2**). The ridge and furrow method resulted in the highest plant height (32.88 cm), number of branches (5.62), number of root nodules (25.59), and pods per plant (6.68) compared to other methods. The improved soil aeration and drainage in the ridges likely enhance root development and nutrient uptake, leading to more robust plant growth. Additionally, the distinct spatial arrangement reduces competition among plants for resources such as light, water, and nutrients, thereby promoting better overall growth and yield attributes[33]. Line sowing on a flat bed showed significantly higher values for these parameters than broadcasting on a flat bed[34]. All growth parameters and yield traits were also significantly affected by the different weed control treatments at 40 DAS. The results clearly indicate that all weed control treatments led to significantly higher growth parameters and yield traits compared to the untreated control. The highest values were observed with two hand weedings at 20 and 40 days after sowing (DAS). Among herbicide treatments, Imazethapyr at 75 g/ha resulted in the

highest plant height (32.03 cm), number of branches (5.70), number of root nodules (25.68), and pods per plant (6.93), comparable to the other three herbicide treatments but significantly superior to the untreated control. The efficacy of Imazethapyr can be attributed to its broad-spectrum activity, which effectively controls a wide range of weed species, thereby reducing competition and promoting better crop growth. The comparable performance of other herbicide treatments indicates that they are also effective in managing weed pressure, but Imazethapyr's specific mode of action and residual activity might offer a slight advantage in sustaining crop growth[35,36]. The lowest values for these parameters were found in the untreated control, which was significantly lower than all other herbicide treatments[37]. The interaction between planting methods and weed control treatments on all growth parameters and yield traits was found to be non-significant at 40 DAS.

**Table 2. Effect of planting methods and weed control methods on growth parameters and yield attributing trait in green gram at 40 DAS**

Treatments	Plant height (cm)	No. of branches/plant	Root nodules/plant	Pods/plant
<b>Planting Methods (P)</b>				
Broadcasting in flat bed	26.6	4.85	21.4	5.50
Line sowing in flat bed	30.4	5.62	25.1	6.62
Ridge and furrow	32.8	5.41	25.5	6.68
SEm±	0.51	0.14	0.56	0.09
C.D. (at 5%)	1.99	0.55	2.19	0.35
<b>Weed control methods (W)</b>				
Pendimethalin 1000 g/ha	29.2	5.30	23.3	6.09
Diclosulam 26 g/ha	31.2	5.37	25.2	6.48
Imazethapyr 75 g/ha	32.0	5.70	25.6	6.93
Hand weeding	31.0	5.52	26.1	6.90
Weedy check	26.3	4.59	19.8	4.93
SEm±	0.95	0.18	0.76	0.21
C.D. (at 5%)	2.76	0.53	2.20	0.60
Interaction (PxW)	NS	NS	NS	NS

### **3.6 Seed yield (kg/ha)**

Different planting methods and weed control treatments significantly influenced seed yield (**Table 3**). The highest seed yield (808 kg/ha) was observed with the ridge and furrow method, which was significantly higher than the line sowing on a flat bed (723 kg/ha) and broadcasting on a flat bed (615 kg/ha). The ridge and furrow method likely promotes better root development and nutrient uptake due to improved soil aeration and drainage. These conditions can enhance plant health and productivity, resulting in higher seed yields. The spatial arrangement in this method also reduces plant competition for resources such as light, water, and nutrients, allowing each plant to grow more robustly[38]. Conversely, the broadcasting method often leads to uneven seed distribution and greater intra-species competition, which can inhibit growth and reduce yield. Weed control treatments resulted in higher seed yields per hectare compared to the untreated control. The highest seed yield (825 kg/ha) was achieved with two hand weedings at 20 and 40 days after sowing (DAS). Among the herbicides, Imazethapyr at 75 g/ha was the most effective, comparable to Diclosulam at 26 g/ha and Pendimethalin at 1000 g/ha. The broad-spectrum activity of Imazethapyr and residual control of a wide range of weed species likely contribute to its effectiveness. By effectively managing weed populations, herbicides minimize competition and enhance resource availability for the crop, leading to higher yields. The lowest seed yield (493 kg/ha) was recorded in the untreated control. The interaction between planting methods and weed control treatments on seed yield was found to be non-significant. These findings are in confirmation with[39,40].

### **3.7 Harvest index (%)**

The ridge and furrow method resulted in the highest harvest index (34.72%), surpassing the other two sowing methods (**Table 3**). Line sowing on a flat bed also achieved a higher harvest index (33.64%) compared to broadcasting on a flat bed (32.44%). Weed control treatments led to a higher harvest index compared to the untreated control. The highest harvest index was observed with Imazethapyr at 75 g/ha (34.27%), followed by Diclosulam at 26 g/ha and Pendimethalin at 1000 g/ha. The lowest harvest index (31.58%) was recorded in the untreated control. Similar findings have been reported by [41].

### **3.8 Economics**

Crops sown using the ridge and furrow method achieved the highest net monetary return (Rs. 15,296/ha) and B:C ratio (1.64), outperforming the other two sowing methods. Line sowing on a flat

bed also recorded a significantly higher net monetary return (Rs. 11,273/ha) and B:C ratio (1.48) compared to broadcasting on a flat bed (Rs. 6,111/ha; 1.26). Among weed control treatments, Imazethapyr at 75 g/ha resulted in the highest net monetary return (Rs. 15,181/ha) and B:C ratio (1.67), followed by Diclosulam at 26 g/ha and Pendimethalin at 1000 g/ha. The lowest net monetary return (Rs. 3,537/ha) and B:C ratio (1.17) was observed in the untreated control. These findings corroborate with[42].

**Table 3. Effect of planting methods and weed control methods on seed yield, harvest index and economics in green gram**

Treatments	Seed yield (kg/ha)	Harvest index (%)	Net monetary returns (Rs/ha)	B:C ratio
<b>Planting Methods (P)</b>				
Broadcasting in flat bed	615	32.44	6111	1.26
Line sowing in flat bed	723	33.64	11273	1.48
Ridge and furrow	808	34.72	15296	1.64
SEm±	16.63	-	-	-
C.D. (at 5%)	63.87	-	-	-
<b>Weed control methods (W)</b>				
Pendimethalin 1000 g/ha	738	33.58	13827	1.61
Diclosulam 26 g/ha	755	33.81	13967	1.60
Imazethapyr 75 g/ha	767	34.27	15181	1.67
Hand weeding	825	34.76	7954	1.24
Weedy check	493	31.58	3537	1.17
SEm±	14.09	-	-	-
C.D. (at 5%)	41.29	-	-	-
Interaction (P×W)	NS	-	-	-

#### 4. CONCLUSION

In conclusion, Imazethapyr at 75 g/ha proved to be the most effective weed control treatment for moongbean, resulting in the lowest weed index. Among the planting methods, the ridge and furrow method demonstrated superior performance, achieving the highest values in growth parameters, yield attributes, seed yield, net monetary returns, and benefit-cost ratio. This combination of effective weed

management and optimal planting techniques maximizes productivity and profitability for moongbean cultivation.

UNDER PEER REVIEW

## REFERENCES

1. Kumar, A., Kadam, S., Arif, M., Meena, R., & Verma, T. (2020). Legumes an alternative land use options for sustaining soil health. *Agriculture & Food e-newsletter*, 6.
2. Ajatasatru, A., Prabhu, V., Pal, B. D., & Mukhopadhyay, K. (2024). Economy-wide impact of climate smart agriculture in India: a SAM framework. *Journal of Economic Structures*, 13(1), 4.
3. Rani, K., Rani, A., Sharma, P., Dahiya, A., Punia, H., Kumar, S., ... & Banerjee, A. (2022). Legumes for agroecosystem services and sustainability. In *Advances in Legumes for Sustainable Intensification* (pp. 363-380). Academic Press.
4. Kumar, D. (2005). Status and direction of arid legumes research in India. *The Indian Journal of Agricultural Sciences*, 75(7).
5. Muchomba, M. K., Muindi, E. M., & Mulinge, J. M. (2023). Overview of greengram (*Vigna radiata* L.) crop, its economic importance, ecological requirements and production constraints in Kenya. *Journal of Agriculture and Ecology Research International*, 24(2), 1-11.
6. IIPR, 2023. <https://iipr.icar.gov.in/mungbean/>
7. Singh G, Kaur H, Aggarwal N, Sharma P. (2015). Effect of herbicides on weeds growth and yield of green gram. *Indian Journal of Weed Science*, 47:38-42.
8. Khairnar, K. Y., Pokharkar, V. G., Kadam, S. A., & Yadav, D. B. (2019). Green gram production technology: An economic analysis. *Journal of Pharmacognosy and Phytochemistry*, 8(3), 2491-2494.
9. Rachaputi, R. C., Chauhan, Y., Douglas, C., Martin, W., Krosch, S., Agius, P., & King, K. (2015). Physiological basis of yield variation in response to row spacing and plant density of mungbean grown in subtropical environments. *Field Crops Research*, 183, 14-22.
10. Pooniya, V., Choudhary, A. K., Dass, A., Bana, R. S., Rana, K. S., Rana, D. S., ... & Puniya, M. M. (2015). Improved crop management practices for sustainable pulse production: An Indian perspective. *The Indian Journal of Agricultural Sciences*, 85(6), 747-458.
11. Fraz RA, Iqbal J, Bakhsh MAAHA. Effect of sowing dates and planting patterns on growth and yield of mungbean (*Vigna radiata* L.) cv. M-6. *Int. J Agri. Biol*, 8(3), 363-365. *International Journal of Chemical studies*. 2006;6(5):2850-2853.
12. Singh, R., Das, T. K., Kaur, R., Raj, R., & Shekhawat, K. (2018). Weed management in dryland agriculture in India for enhanced resource use efficiency and livelihood security. *Proceedings of the National Academy of Sciences, India Section B: Biological Sciences*, 88, 1309-1322.
13. Kumara, P., Pakeerathan, K., & Deepani, L. P. (2021). Assessment of yield loss in green gram (*Vigna radiata* (L.) R. Wilczek) cultivation and estimation of weed-free period for eco-friendly weed management. In *Biology and Life Sciences Forum*, 3(1): 22-28.
14. Sinchana, J. K., & Raj, S. K. (2023). Weed management in pulses: a review. *Legume Research-An International Journal*, 46(5), 533-540.
15. Dash M, Tandon M, Mohapatra S. A Review on Integrated Weed Management in Green Gram. *Indian Journal of Weed Science*. 2018;7(06):1865-1871.
16. Verma, B., Bhan, M., Jha, A. K., & Porwal, M. (2023). Influence of weed management practices on direct-seeded rice grown under rainfed and irrigated agroecosystems. *Environment Conservation Journal*, 24(3), 240-248.
17. Pahade, S., Jha, A. K., Verma, B., Meshram, R. K., Toppo, O., & Shrivastava, A. (2023). Efficacy of sulfentrazone 39.6% and pendimethalin as a pre emergence application against weed spectrum of soybean (*Glycine max* L. Merrill). *International Journal of Plant & Soil Science*, 35(12), 51-58.
18. Sahu, V., Kewat, M. L., Verma, B., Singh, R., Jha, A. K., Sahu, M. P., & Porwal, M. (2023). Effect of carfentrazone-ethyl on weed flora, growth and productivity in wheat. *The Pharma Innovation Journal*, 12(3), 3621-3624.
19. Verma B, Bhan M, Jha AK, Agrawal KK, Kewat ML and Porwal M. 2023. Weed management in direct-seeded rice (*Oryza sativa*) in central India. *Indian Journal of Agronomy*. 68 (2): 217-220.
20. Kaur, R., & Kaur, S. (2018). Controlling Weeds in Major Pulse Crops by integrated weed management. *Indian Farming*, 66(8).

21. Raghav, P., Jha, A. K., Badal, V., Rahul, K., & Richa, S. (2023). Bioefficacy of pinoxaden as post-emergence herbicide against weeds in wheat crop. *Pollution research*, 42(1), 115-117.
22. Tiwari, R.K., Khan I.M. , Singh Nirmla and Amit Jha. 2011a. Chemical weed control in wheat through on farm demonstration in Rewa district of Madhya Pradesh. *Indian Journal of Weed Science*. 43(3&4): 215–216.
23. Tomar, D. S., Jha, A. K., Porwal, M., Verma, B., Tirkey, S., Khare, Y., ... & Chouhan, M. (2023). Efficacy of Halauxifen-methyl+ Florasulam against Complex Weeds in Wheat under Kymore Plateau and Satpura Hill Zone of Madhya Pradesh, India. *International Journal of Plant & Soil Science*, 35(15), 161-171.
24. Kaur, R., & Kaur, S. (2021). Strategies for integrated weed management in major pulse crops. *Indian Farming*, 71(8).
25. Sahu M P, Kewat ML, Jha AK, Sondhia S, Choudhary VK, Jain N and Verma B. 2022. Weed prevalence, root nodulation and chickpea productivity influenced by weed management and crop residue mulch. *AMA, Agricultural Mechanization in Asia, Africa and Latin America* 53(6): 8511-8521.
26. Gill, H. S. (1969). Weed index-a new method for reporting weed control trials. *Indian Journal of Agronomy*, 14, 96-98.
27. Kade, S. K., Sethi, H. N., Goud, V. V., & Patil, A. N. (2014). Effect of herbicides on weed, nutrient uptake, soil micro flora and yield of mungbean. *Panjabrao Deshmukh Krishi Vidyapeeth Research Journal*, 38, 37-41.
28. Singh, S. P., & Yadav, R. S. (2015). Effect of weed management on growth, yield and nutrient uptake of greengram. *Indian Journal of Weed Science* 47(2): 206–210.
29. Samant, T. K., & Mohanty, T. R. (2017). Effect of sowing date and weed management on productivity and economics of rainfed mungbean (*Vigna radiata*). *Indian journal of agronomy*, 62(3), 332-337.
30. Peterson, M. A., Collavo, A., Ovejero, R., Shivrain, V., & Walsh, M. J. (2018). The challenge of herbicide resistance around the world: a current summary. *Pest management science*, 74(10), 2246-2259.
31. Kaur, G., Brar, H. S., & Singh, G. (2009). Effect of weed management on weeds, growth and yield of summer mungbean [*Vigna radiata* (L.) R. Wilczek]. *Indian Journal of weed science*, 41(3and4), 228-231.
32. Verma B, Bhan M, Jha AK, Porwal M and Patel R. 2023. Assessment of different herbicides for effective weed management in direct seeded rice technology. *Ecology, Environment and Conservation Journal*. 29 (3): 211-217.
33. Karotiya, N., & Gangwar, B. (2019). Effect of diverse sowing methods on organic mungbean production in Bundelkhand region of India. *Journal of Food Legumes*, 32(4), 268-271.
34. Zaryal, K., Barai, S. M., Mohammadi, N., Stori, R. M., Nasrat, N. A., Sayedi, S. A., ... & Ahmadi, H. (2020). Effects of Crop Establishment Methods and Phosphorus Fertilization on Growth and Yield of Mungbean (*Vigna radiata* L. Wilczek.) in Kandahar Region of Afghanistan.
35. Khairnar, C.B., Goud, V.V. and Sethi, H.N. 2014. Pre and post emergence herbicides for weed management in mungbean. *Indian Journal of Weed Science* 46(4):392-395.
36. Yadav R, Kumar S, Dhaka AK, Kumar N. Effect of planting methods and weed management practices on yield of green gram (*Vigna radiata* (L.) R. Wilczek), weed dynamics vis a vis phytotoxicity in green gram. *Indian Journal of Agricultural Research*. 2019;53(2):158- 1.
37. Kundu, R., Bera, P. S., & Brahmachari, K. (2009). Effect of different weed management practices in summer mungbean (*Vigna radiata* L.) under new alluvial zone of West Bengal. *Journal of Crop and Weed*, 5(2), 117-121.
38. Yadav, M. K., Yadav, A., Singh, A. K., Mahajan, G., Singh, M. K., Singh, R. S., ... & Babu, S. (2012). Ridge Planted Pigeonpea and Furrow Planted Rice in an Intercropping System as Affected by Nitrogen and Weed Management. INTECH Open Access Publisher.
39. Singh, G., Virk, H. K., & Khanna, V. (2020). Effect of land configuration and weed management on productivity of greengram (*Vigna radiata*). *The Indian Journal of Agricultural Sciences*, 90(5), 947-951.
40. Verma, L., & Kushwaha, H. S. (2020). Evaluation of different herbicides against weeds in mungbean (*Vigna radiata* L.). *Legume Research-An International Journal*, 43(6), 866-871.

41. Mirjha, P. R., Prasad, S. K., Singh, M. K., Paikra, R. H., Patel, S., & Majumdar, M. (2013). Effect of weed control measures on weeds, nodulation, growth and yield of mungbean (*Vigna radiata*). *Indian Journal of Agronomy*, 58(4), 615-617.
42. Bahar, F. A., Dar, S. A., Lone, A. A., Ansarul Haq, S., Alie, B. A., Dar, Z. A., ... & Zaffar, G. (2017). Effect of land configuration and weed management on mungbean productivity under temperate conditions of Kashmir, India. *Inter J Curr Microbiol Appl Sci*, 6(10), 863-70.

UNDER PEER REVIEW