

NUTRITIONAL COMPOSITION OF “OSE-OKU”, A TRADITIONAL FOOD OF ABATETE IN ANAMBRA STATE

ABSTRACT

Aims: This study assessed the nutritional composition of “Ose-Oku”, a traditional food of Abatete in Anambra State, Nigeria.

Study Design: The research involved the preparation of “Ose-Oku” as traditionally done, followed by comprehensive laboratory analyses to determine its nutritional composition. Proximate, mineral, vitamin, antinutrient, fatty acid, and amino acid analyses were conducted using established scientific methods.

Place and Duration of Study: The study was carried out in a laboratory setting with samples collected from Abatete in Anambra State. The duration of the study spanned several months, allowing for thorough analysis and validation of results.

Methodology: The food was prepared as it is done in the traditional setting. The methods of AOAC (1984) were used in the proximate analysis of the dried food sample. Mineral and vitamin analyses were carried out using Atomic Absorption Spectrophotometer. Antinutrient and fatty acid analyses were done using Gas chromatography. HPLC apparatus was used to determine the amino acid composition.

Results: Among the nutrients found in the studied food sample were carbohydrates ($82.56 \pm 0.009\%$), crude protein ($7.67 \pm 0.042\%$), crude fat ($1.49 \pm 0.004\%$), crude fibre ($0.44 \pm 0.011\%$), moisture ($6.05 \pm 0.005\%$), ash ($1.78 \pm 0.002\%$), and total amino acids (72.64%). Also total SFA (26.62%), total UFA (85.09%), and total PUFA (70.60%) were found. Vitamins A (0.71mg/100g), C (72.67mg/100g), D (0.73mg/100g), E (3.92mg/100g), K (7.83mg/100g), B1 (0.04mg/100g), B3 (0.58mg/100g), B6 (0.20mg/100g), B9 (3.48mg/100g), and B12 (0.25mg/100g) were also found. Minerals such as iron (0.02mg/100g), manganese (0.02mg/100g), zinc (0.16mg/100g), sodium (0.20mg/100g), potassium (3.00mg/100g), magnesium (2.08mg/100g), calcium (3.61mg/100g), and copper (0.47mg/100g) were present. Antinutrients such as Kaempferol,

Cyanogenic Glycosides, Narigenin, Ammodendrine, Tannins, Flavonones, Flavone, Proanthocyanidin, and Spartein were present in the food.

Conclusion: This study has shown that the proportionate combination of the food materials enhanced the nutrient quality of this traditional food sample. Besides the low content of moisture and fats with high carbohydrate, appreciable amount of protein, fibre and vitamins qualify its complimentary use to tackle food insecurity and malnutrition.

Keywords: *Ose-Oku, Proximate, mineral, vitamin, antinutrient, fatty acid, and amino acid analyses.*

1. INTRODUCTION

Traditional foods are the origin of diet in various cultures and societies. Since the onset of food industrialization, that is, the mass manufacture that began in the mid-twentieth century, a clear difference in quality was observed, especially by consumers, which divided food into two large groups: those manufactured in mass, homogeneous and from which the source of the raw material with which they are manufactured is not known, neither the method of explanation; and those that are manufactured in small scale, to a certain point heterogeneous and we could say that artisan, and of which the link producer–consumer is closer, since the method of explanation is not absolutely known, the source of the raw material can be indirect, therefore resulting in a better assurance for the consumers. Consequently, it can be said that, ironically, globalization inhibited and encouraged the consumption and manufacture of these foods. Consequently, interest in these foods has increased since the 1990s, especially focused on enhancing their significance and protecting them [1,2,3].

“Ose Oku”, a traditional food of the Abatete people, is among the foods of plant origin that contains nutrients and phytochemicals. Though, this very traditional food, Ose Oku, has been extincted, since after the Biafra-Nigeria civil war, when it was predominantly consumed. Abatete is a community in Idemili North L.G.A. of Anambra State, South-eastern Nigeria. The people of Abatete are of Igbo ethnicity, and speak Igbo as their dialect. Ose Oku was predominantly consumed during the Biafra-Nigeria civil war due to the fact that it was very

easy and quick to prepare ipso-facto as an emergency traditional food. Amadi et al., 2017 worked on nutritional, phytochemical and sensory evaluation of "Mberiaagworagwo", a traditional food of Uruagu-Nnewi people in Anambra State, Nigeria [4]. Also, Okwu & Okwu, 2004 researched on phytochemicals and vitamin content of indigenous spices of South-eastern Nigeria [5]. Another research was carried out by Duru et al., 2012, on nutrient composition of "Nduduagworagwo", a traditional food of Akokwa people in Ideato L.G.A. of Imo State, Nigeria [6]. Then Majesty et al., 2013 worked on fatty acid composition and sensory evaluation of "Nduduagworagwo", a local food of Akokwa people in Imo State, Nigeria [7]. Yet some traditional diets in the South-eastern part of Nigeria, such as Ose Oku, are going into extinction, thereby reducing the availability and variety of foods needed to cushion the effect of the food insecurity in Nigeria. Consequently, it is imperative to assess the nutritional composition of Ose Oku, a traditional food of the Abatete people in Anambra State, in the South-eastern part of Nigeria. A survey of available literature shows that nutritional composition of Ose Oku has not been studied.

The present study is geared towards the investigation of the nutritional composition of "Ose Oku", a traditional food. This will help to expose the possible health benefits on consuming the food, and its possible acceptability.

2. MATERIALS AND METHODS

2.1 Sample Collection

The major raw materials and ingredients such as yam, dried fish, biscuit bone of beef, etc, used in the preparation of the traditional diet used in this study, were purchased from "Eke Agu" market in Abatete, Idemili North Local Government Area of Anambra State, South-eastern Nigeria.

2.2 Preparation of "Ose Oku"

Recipe: About 200g of peeled yam (a thickner), 12.31g of fresh pepper, 1g of salt, 850g of dried fish, 2.5g of fermented sesame seed (ogiri), 200ml of water

Procedure: About 200ml of water, together with 850g of dried fish, was put in 2000mL pot and boiled for 2 minutes, using gas burner. Then 12.31g of fresh pepper, 1g of salt, 2.5g of fermented sesame seed (ogiri) were added. Also, 200g of sliced peeled yam was washed,

boiled and pounded, to produce a paste. The yam paste was added to the boiling soup. It was allowed to boil for about 15 minutes, and then put down from fire.

2.3 Processing of traditional food sample for analysis

The prepared sample of "Ose Oku" was dried in an oven at 70°C for 48 hours. The dried food sample was ground, with a hand mill, into powdered form and stored in air-tight sample containers at 4°C, until required for analysis.

2.4 Proximate analysis of traditional food sample

The methods of AOAC (1984) were used in the proximate analysis of the food sample [8].

2.5 Determination of amino acid composition of the sample

A 0.1g lyophilized sample was weighed into a 16 × 125mm screw-cap Pyrex (Barcelona, Spain) tube, 15 ml of 6N hydrochloric acid was added, and the tube was thoroughly flushed with N₂, quickly capped, and placed in an oven at 110°C for 24 h (17). After hydrolysis, the tube contents were vacuum filtered (Whatmann #541, Maidstone, England) to remove solids, the filtrate was made up to 25 ml with water, and an aliquot of this solution was further filtered through a 0.50-µm pore-size membrane (Millipore, Madrid, Spain). A standard solution containing 1.25 µmol/ml of each amino acid in 0.1N hydrochloric acid was created.

2.6 Determination of fatty acid profile analysis of the sample

One millilitre (1ml) of filtered residue was dissolved in 50ml of chloroform and transferred to a 100ml volumetric flask and was then diluted to the mark. Most of the chloroform was evaporated at room temperature. 1 ml of the reagent (20% vol. benzene and 55% vol. methanol) was added into the volumetric flask, sealed, and heated at 40°C in a water bath for 30 minutes. After heating, the organic sample, with hexane and water, was extracted, so that the final mixture of the reagent, hexane and water, was in proportion of 1:1:1 (i.e. 1ml each of hexane and water to the reaction mixture). The mixture was shaken vigorously by hand for 2min. Due to a stable emulsion that was formed, it was broken by centrifugation. About half of the top hexane phase was transferred to a small test tube for injection. Only the organic layer was carefully removed. Injection was not done directly from the reaction vial because of the risk of injecting water, as it can ruin the GC column.

2.7 Determination of vitamins

Vitamin A was estimated by the method of Bayfield and Cole, 1980 [9]. Vitamin E was estimated in the samples by the Emmerie-Engel reaction as reported by Emmerie & Engel, 2010 [10]. Vitamin C was analyzed by the spectrophotometric method described by Roe and Kuether, 1943 [11]. Vitamin D was assayed according to the method reported by Holick, 2009 [12].

2.8 Determination of Heavy Metals

Heavy metal analysis was conducted using Varian AA240 Atomic Absorption Spectrophotometer according to the method of APHA, 1995 (American Public Health Association) [13].

2.9 Determination of the phytochemical and anti-nutrient contents of the traditional food sample

One gram (1g) of the ground food sample was weighed and transferred into a test tube and 15ml ethanol and 10ml of 50% potassium hydroxide was added. The test tube was allowed to stay in a water bath at 60°C for 60mins. After the reaction time, the reaction product contained in the test tube was transferred to a separatory funnel. The tube was washed successfully with 20ml of ethanol, 10ml of cold water, 10ml of hot water and 3ml of hexane, which were all transferred to the separatory funnel. These extracts were combined and washed three times with 10ml of 10%v/v ethanol aqueous solution. The solution was dried with anhydrous sodium sulfate and the solvent was evaporated. The sample was solubilized in 1000µl of pyridine of which 200µl was transferred to a vial for analysis. The analysis of anti-nutrient was performed on a BUCK M910 Gas chromatography equipped with a flame ionization detector. A RESTEK 15 meter MXT-1 column (15m x 250µm x 0.15µm) was used. The injector temperature was 280°C with split-less injection of 2µl of sample and a linear velocity of 30cms⁻¹, Helium 5.0pa.s was the carrier gas with a flow rate of 40 mlmin⁻¹. The oven operated initially at 200°C, it was heated to 330°C at a rate of 3°C min⁻¹ and was kept at this temperature for 5min. the detector operated at a temperature of 320°C. Anti-nutrients were determined by the ratio between the area and mass of internal standard and the area of the identified anti-nutrients. The concentration of the different anti-nutrients was expressed in µg/g.

2.10 Statistical analyses

All values were reported as mean \pm standard deviation. Data obtained from the various parameters were analyzed statistically by analysis of variance (ANOVA) using the IBM Statistical Package for Social Sciences (SPSS) 24.0 software programme. Significance testing was done at 95% confidence interval.

3. RESULTS AND DISCUSSION

The results of the proximate composition of "Ose Oku" are presented in Table 1. This showed the carbohydrate ($82.56 \pm 0.009\%$), crude protein ($7.67 \pm 0.042\%$), crude fat ($1.49 \pm 0.004\%$), crude fibre ($0.44 \pm 0.011\%$), and moisture ($6.05 \pm 0.005\%$). The moisture content obtained in this study was very much lower, when compared to values obtained by Amadi (2011) (22.73%) [14], and Ejiogu (2019) (11.50%) [15]. The low moisture content found in "Ose Oku" traditional food could be due to the method of drying undertaken on the traditional food sample, to prevent microbial growth, and aid for longer storage. High moisture content in foods has been shown to encourage microbial growth [16]. Ash represents the mineral matter left after feeds are burnt in oxygen [17]. It is used as a measure of the mineral content in any sample [18]. The ash obtained was $1.78 \pm 0.002\%$. The energy value was 57.98 Kcal/100g. The energy value of "Ose Oku" traditional food sample in this study was very much lower, when compared to the values reported by Amadi (2011) (348 - 499 Kcal/100g), and the values obtained in Beniseed soup (572.97 - 666.05 Kcal/100g), a local delicacy cherished in the Northern Cross Rivers State of Nigeria, by Agiang *et al.* (2010) [14,19].

Table 1: Proximate composition (%) of "Ose Oku"

Proximate (%)	Ose Oku
Carbohydrate	82.56 ± 0.009
Crude Protein	7.67 ± 0.042
Crude Fat	1.49 ± 0.004
Crude Fiber	0.44 ± 0.011
Moisture	6.05 ± 0.005
Ash	1.78 ± 0.002
Energy value (Kcal/100g)	57.98 ± 0.165

Values are means \pm standard deviations of triplicate determinations.

The amino acid composition of "Ose Oku" traditional food showed that the food sample contains some quantity of essential and non-essential amino acids, as presented in Table 2. Arginine, lysine, aspartate and glutamate were very much high in "Ose Oku". According to studies, arginine boosts immune function, hastens healing process, promotes cell division, and triggers the release of hormone [20,21,22]. Lysine aids in cardiac arrest [23] and its deficiency results to immunodeficiency [24]. Onyeike *et al.* (2005) reported that arginine and histidine are required for the growth of infants [25]. The total saturated fatty acid (SFA) content of "Ose Oku" (26.62%), as revealed in Table 4, is comparably higher than those of some Middle Eastern traditional dishes in Lebanon that ranged from 2.8% to 24.8%, as reported by Hoteit *et al.* (2021) [26]. Linoleic acid was the most abundant fatty acid in "Ose Oku" traditional diet. The total unsaturated fatty acids (UFA) content of "Ose Oku" was 85.09%; linoleic acid and oleic acid were the major UFA and PUFA in the traditional food sample, but linoleic acid was most abundant (41.75%). MUFA and PUFA have beneficial health and dietary values. They contribute to a decrease in the occurrence of vascular syndrome, carcinoma of the uterine cervix and immunological diseases [27]. Knowledge of the ratio of PUFA to SFA is essential in most nutritional studies. The British Department of Health and Social Security considers PUFA/SFA values lower than 0.45 as inappropriate for human health due to their association with cardiac diseases [28]. The PUFA/SFA ratio of Ose Oku is 2.65, ipso-facto can be said to be very much appropriate for human health with respect to cardiac disease.

Table 2: Amino acid content of "Ose Oku" traditional diet

Aminoacids (mg/g)	Ose Oku
ESSENTIAL	
Arginine	5.69
Histidine	2.68
Isoleucine	4.20
Leucine	2.27
Lysine	6.66
Methionine	1.37
Phenylalanine	3.95
Threonine	3.70

Tryptophan	1.68
Valine	2.46
NON-ESSENTIAL	
Alanine	1.75
Aspartate	10.35
Cysteine	2.47
Glutamate	14.34
Glycine	2.60
Proline	1.50
Serine	2.40
Tyrosine	2.57
TOTAL	72.64

Table 3: Fatty acid content of "Ose Oku" traditional diet

Fatty acid components (%)	Ose Oku
StearicAcid (C _{18:0})	5.69
LauricAcid (C _{12:0})	10.25
PalmiticAcid (C _{16:0})	10.68
OleicAcid (C _{18:1})	14.49
LinoleicAcid (C _{18:2})	41.75
LinolenicAcid (C _{18:3})	5.04
DocosahexaenoicAcid (C _{22:6})	2.56
AdrenicAcid (C _{22:4})	21.25
TOTAL SFA	26.62
TOTAL UFA	85.09
TOTAL PUFA	70.60
UFA/SFA	3.20
PUFA/SFA	2.65

SFA = Saturated Fatty Acids;

UFA = Unsaturated Fatty Acids;

PUFA = Polyunsaturated Fatty Acids;

PUFA/SFA value below 0.45 is inappropriate for the body system.

The concentrations of some of the phytochemicals and antinutrients present in "Ose Oku" traditional food are presented in Table 5. Antinutrients are natural or artificial compounds that meddle with the assimilation of nutrients [29]. The "Ose Oku" traditional food sample was analyzed qualitatively and quantitatively for the presence of saponins, tannins, phytates, flavonoids, oxalates, cyanogenic glycosides, cardiac glycosides, kaempferol, catechin, proanthocyanidin and sapogenin. Tannins are protective substances produced majorly by plants used for treatment of wound, inflamed mucous membrane and astringent in nature [30]. Tannins also have some pharmacological activity like soothing relief, anti-inflammatory, diuretics, frostbite, burn and regeneration of skin hemorrhoids [5,31]. Butler, 1989 reported that tannins have possible anticarcinogenic effect. Dietary phytate could be associated to the prevention of dental decay, kidney stones, and calcification of blood vessels [29]. Oxalates and phytates bind calcium, magnesium and iron in the body; acts as pathogenic factor, thereby enhancing the acquisition of nutrients [32]. There is concern about oxalates because diets with high level of oxalates can be a risk for renal calcium absorption [33]. The toxic level of soluble oxalate is, however, reported to range from 2.0g to 5.0g [34].

Table 4: Phytochemical and antinutrient content of "Ose Oku" traditional diet

Antinutrients ($\mu\text{g/g}$)	Ose Oku
Kaempferol	5.59
CardiacGlycosides	3.32
Steroid	4.14
Catechin	3.80
Anthocyanin	2.29
Dihydrocystidine	3.68
CyanogenicGlycosides	9.46
Aphyllidine	3.66
Narigenin	8.45
Ammodendrine	5.99

Tannins	18.88
Flavonones	9.37
Flavone	6.77
Proanthocyanidin	8.75
Phytates	4.15
Sparteine	7.63
Oxalates	2.53
Ephedrine	3.48
Sapogenins	0.02

4. CONCLUSION

The proximate and nutritional composition of "Ose Oku" traditional food from Abatete in Anambra State cannot be overemphasized. This indicates their abundance in nutrients, and encourages their usage as complementary diets for children, adults, and people suffering from malnutrition, especially when taken at 100%.

5. REFERENCES

1. Cerjak, M., Haas, R., Brunner, F., & Tomic, M. (2014). What motivates consumers to buy traditional food products? Evidence from Croatia and Austria using word association and laddering interviews. *British Food Journal*, 116(11), 1726-1747. <https://doi.org/10.1108/BFJ-02-2-14-0090>.

2. Trichopoulou, A., Vasilopoulou, E., Georga, K., Soukara, S., & Dilis, V. (2006). Traditional foods: why and how to sustain them. *Trends in Food Science & Technology*, 17(9), 498-504.
3. Trichopoulou, A., Soukara, S., & Vasilopoulou, E. (2007). Traditional foods: a science and society perspective. *Trends in Food Science & Technology*, 18(8), 420-427. <https://doi.org/10.1016/j.tifs.2007.03.007>.
4. Amadi, B. A., Duru, M. K., Agomuo, E., Amadi, P., & Onedibe, O. (2017). Nutritional, phytochemical and sensory evaluation of “Mberiagworagwo”, a traditional food of Uruagu-Nnewi people in Anambra State, Nigeria. *Journal of Advances in Biology and Biotechnology*, 14(1), 1-8. ISSN: 2394-1081. <https://doi.org/10.9734/JABB/2017/27901>.
5. Okwu, D. E. & Okwu, J. P. (2004). Phytochemical and vitamin content of indigenous spices of South-Eastern Nigeria. *Journal of Sustainable Agriculture and Environment*, 6(2), 30-34.
6. Duru, M. K. C., Agomuo, E. A., & Amadi, B. A. (2012). Nutrient composition of “Nduduagworagwo”, a traditional food of Akokwa people in Ideato L. G. A. of Imo State, Nigeria. *Continental Journal of Science and Technology*, 6(3), 27-32. <https://doi.org/10.5707/cjfst.2012.6.3.19.26>.
7. Majesty, D., Amadi, B. A., Arukwe, U., Adindu, E., Amadi, C., & Onuoha, N. (2013). Fatty acid composition and sensory evaluation of “Nduduagworagwo”, a local food of Akokwa people in Imo State, Nigeria. *Global Research Journal of Science*, 2(2), 67-72.
8. AOAC (1984). *Official method of analysis* (14th Edition). Association of Official Analytical Chemists, AOAC, Arlington, Washinton DC, 249-252.
9. Bayfield, R. F., & Cole, E. R. (1980). Colorimetric estimation of vitamin A with trichloroacetic acid. *Methods in enzymology*, 67, 189-195. [https://doi.org/10.1016/s0076-6879\(80\)67026-8](https://doi.org/10.1016/s0076-6879(80)67026-8).
10. Emmerie, A., & Engel, C. (2010). Colorimetric determination of tocopherol (vitamin E): III estimation of tocopherol in blood serum. *Recueil des Travaux Chimiques des Pays-Bas*, 58(10), 895-902. <https://doi.org/10.1002/recl.19390581007>.
11. Roe, J. H., & Kuether, C. A. (1943). The determination of ascorbic acid in whole blood and urine through the 2,4-dinitrophenylhydrazine derivatives of dehydroascorbic acid. *Journal of Biological Chemistry*, 147(2), 399-407. [https://doi.org/10.1016/S0021-9258\(18\)72395-8](https://doi.org/10.1016/S0021-9258(18)72395-8).
12. Holick, M. F. (2009). Vitamin D status: measurement, interpretation, and clinical application. *Annals of epidemiology*, 19(2), 73-78. <https://doi.org/10.1016/j.annepidem.2009>.
13. APHA (1995). *Standard methods for the examination of water and wastewater* (19th Edition). American Public Health Association Inc., New York.
14. Amadi, B. A. (2011). Nutritional evaluation of selected traditional diets of the Ikwerre people of Niger Delta, Nigeria. Ph.D Thesis, Department of Biochemistry, University of Port Harcourt

15. Ejiogu, A. G. (2019). Nutritional evaluation of diets formulated from cocoyam, three leaf yam, saba banana and their effect on selected biochemical parameters of wistar albino rats. M.Sc. dissertation, Department of Biochemistry, Faculty of Science, University of Port Harcourt
16. Temple, V. J., Badamosi, E. J., Ladeji, O. & Solomon, M. (1996). Proximate chemical composition of three locally formulated complementary foods. *West African Journal of Biological Sciences*, 5(2), 134-143.
17. Bingham, S. (1978). *Nutrition: A Consumer's Guide to Good Eating*. Transworld Publishers, London. 123-127.
18. Pearson, D. A. (1981). *Chemical Analysis of Foods* (8th Edition). Churchill Livingstone, Edinburg, London. 313-316.
19. Agiang, M. A., Umoh, I. B., Essien, A. I. & Eteng, M. U. (2010). Nutrient changes and anti-nutrient contents of beniseed soup during cooking using a Nigerian traditional method. *Pakistan Journal of Biological Sciences*, 13(20), 1011-1015. <https://doi.org/10.3923/pjbs.2010.1011.1015>
20. Tapiero, H., Mathe, G., Couvreur, P. & Tew, K. D. (2002). L-Arginine (review). *Biomedicine and Pharmacotherapy*, 56(9), 439-445.
21. Stechmiller, J. K., Childress, B. & Cowan, L. (2003). Arginine supplementation and wound healing (review). *Nutrition in Clinical Practice*, 20(1), 52-61.
22. Witte, M. B. & Barbul, A. (2003). Arginine physiology and its implication for wound healing (review). *Wound Repair and Regeneration*, 11(6), 419-423.
23. Flodin, N. W. (1997). The metabolic roles, pharmacology, and toxicology of lysine. *Journal of American College of Nutrition*, 16(1), 7-21.
24. Chen, C., Sander, J. E. & Dale, N.M. (2003). The effect of dietary lysine deficiency on the immune response to Newcastle disease vaccination in chicken. *Avian Discovery*, 47(4), 1346-1351.
25. Onyeike, E. N., Ayalogu, E. O. & Okaraonye, C. C. (2005). Nutritive value of the larvae of *Raphia* palm beetle (*Oryctes rhinoceros*) and weevil (*Rhyncophorus pheonicis*). *Journal of the Science of Food and Agriculture*, 85, 1822-1828.
26. Hoteit, M., Zoghbi, E., Rady, A., Shankiti, I. & Al-Jawaldeh, A. (2021). Fatty acids quality in middle Eastern traditional dishes, Arabic sweets and market foods frequently consumed in Lebanon: *Nutrients*, 13(7), 2462
27. Tonial, I. B., Aguiar, A. C., Oliveira, C. C., Bonaafe', E. G., Visentainer, J. V. & de Souza, N. E. (2009). Fatty acid and cholesterol content, chemical composition and sensory evaluation of horse meat: *South African Journal of Animal Science*, 39(4), 328-332.
28. HMSO, UK (1994). Nutritional aspects of cardiovascular disease. Department of Health. *London Report on Health and Social Subject*, 46, 37-46.

29. Cammack, R., Atwood, T., Campbell, P., Parish, H., Smith, A., Vella, F., & Stirling, J. (2006). *Oxford dictionary of biochemistry and molecular biology* (Rev. Ed.). 47. Oxford University Press. ISBN 9780198529170.
30. Obichi, E. A., Monago, C. C. & Belonwu, D. C. (2015). Effect of *Cnidoscolus aconitifolius* (Family: *Euphorbiaceae*) aqueous leaf extract on some antioxidant enzymes and haematological parameters of high fat diet and streptozotocin induced diabetic wistar albino rats. *Journal of Applied Science and Environmental Management*, 19(1), 201-209.
31. Osuntokun, O. T. & Oluwafoise, B. O. (2015). Phytochemical screening of ten Nigerian medicinal plants. *International Journal of Multidisciplinary Research and Development*, 2(4), 390-396.
32. Agoreyo, B. O., Obansa, E. S. & Obanor, E. O. (2012). Comparative nutritional and phytochemical analyses of two varieties of *Solanum melongena*. *Science World Journal*, 7(1), 1597-6343.
33. Osagie, A. U. (1998). *Nutritional quality of plant foods*. Post Harvest Research Unit, Department of Biochemistry, University of Benin. 221-244.
34. Oke, O. L. (1966). Chemical studies on the more commonly used leafy vegetables in Nigeria. *Journal of West African Science Association*, 11, 42-48.