

ASSESSMENT OF PHYSICO-CHEMICAL PROPERTIES OF SOIL FROM DIFFERENT BLOCKS OF MIRZAPUR DISTRICT, UTTAR PRADESH, INDIA

ABSTRACT

An Assessment of Physico-chemical properties of soil from different blocks of Mirzapur district, Uttar Pradesh carried out 2023-24. The prime objectives of this study were to carried out the physico-chemical properties of soil at different Blocks of Mirzapur District, Uttar Pradesh to determine the availability of macro nutrient in the soil. Soil samples were collected with depth of 0-15 cm, 15-30 cm and 30-45 cm respectively. Soil textural classes were sandy loam. Bulk Density (1.17Mg m⁻³ to 1.44Mg m⁻³). Particle Density (2.22 Mg m⁻³ to 2.48 Mg m⁻³). % Pore Space (42.78% to 48.00) %. It clearly indicated that soil has good Water Holding Capacity (39.92 to 45.76) % and good physical condition, The pH of soil is alkaline nature (7.43 to 8.15) and the Electrical Conductivity (0.21 to 0.34 dSm⁻¹) was suitable for all crops. These soils have low Organic carbon (0.27 to 0.38%). Nitrogen range is medium (250 kg ha⁻¹ to 380 kg ha⁻¹) in all villages. Phosphorus (22.53 kg ha⁻¹ to 37.49 kg ha⁻¹) content is medium in all villages. Potassium (148.43 kg ha⁻¹ to 282.4 kg ha⁻¹). is medium at eight locations and high in one location.

Keywords:

INTRODUCTION

Soil suitability for agriculture is mostly determined by its physical characteristics. The physical characteristics of the soil are closely related to its capacity to support life, to move, retain, and make water and nutrients available to plants, to facilitate root penetration, and to allow for the passage of heat and air. Chemical and biological qualities are influenced by physical attributes as well. an explanation of the physical characteristics of soils and their significance for nutrient and water mobility as well as the growth of vegetation cover.(Phogat *et al.*, (2015). Soil is the base for the existence of many life forms and an indispensable medium for plant growth. The overarching definition that resonates today is "Soil is a natural independent body which like any other natural body or organism, has a specific origin, history of development, and external appearance" (Hartemink *et al.*, 2016). Soil formation is a constructive as well as destructive process. Soil is composed of particles of broken rock that have been altered by chemical and mechanical processes that weathering and erosion. Soil has a complex function which is beneficial to human and other living organism Soil is not merely a group of mineral particles. It has also a biological system of living organism as well as some other components. The climate and other factor largely affect the soil formation. (Tale and Ingole,2015) The study of soil is known as Soil Science or pedology (pedo means Soil) and (logy means to study). This study is also known as Edaphology (Edaphos means Soil). Soil may also be defined as the part of the earth crust in which humus is present as in organic part (Das, 2021).

Commented [a1]: -The general methodology has not been provided briefly.
- For keywords, they have not been mentioned

Commented [a2]: The introduction is only one paragraph. Please describe the background and problems why this study was important.
In the end of introduction, please also provide the objective of this study and what is the implementation of the output of this study

MATERIALS AND METHODS

Sampling site and collection

Mirzapur is located at 25.15°N 82.58°E. It has an average elevation of 80 metres (265 feet). The District of Mirzapur lies between the parallels of 23.52 & 25.32 North latitude and 82.7 and 83.33 East longitude. It forms a portion of the Varanasi district. On the north and north-east it is bounded by the Varanasi district; on the south bounded by Sonbhadra district; on the north-west by Allahabad district. The shape to the north and west is totally regular. In no direction, except for about 13 km. in the north-east where the Ganges separates the Tehsil of Chunar from the district of Varanasi, has Mirzapur a natural frontier. The Chanvar fields, considered to be one of the most fertile lands tracts in India, are located on Gangetic flood plains of the district..

Mirzapur district lies in the South-East part of Uttar Pradesh. With an area of 4521 km² and population of more than 20 lacs, the district is divided in four sub division, 12 blocks and 973 gramsabhas containing 1698 villages. Agro-climatically, the district falls under two zones, viz. Indo-Gangetic Plains covering only 30-40 percent of the total area and Vindhyan Zone covering remaining area. The area under Gangetic Plains is endowed with rich alluvial and fertile soil and good irrigation facilities while the Vindhyan Zone has meagre resource of water and the land is mostly degraded.

Soil samples were collected from 3 different Blocks of Mirzapur district in Uttar Pradesh.. Samples were collected randomly using soil auger by composite sampling method at depths of 0-15, 15-30 and 30-45 cm. All the samples were divided into four parts and then among them two samples were collected and only half kg sample is being taken for the soil analysis by the conning and quartering method.

Table 1. The methods of analysis for different soil parameters

S. No.	Particulars	Scientist Name	Methods	Unit
PHYSICAL PROPERTIES				
1.	Bulk density	Muthuval <i>et al.</i> (1992)	Graduated measuring cylinder	Mg m ⁻³
2.	Particle density	Muthuval <i>et al.</i> (1992)	Graduated measuring cylinder	Mg m ⁻³
3.	Textural class (Sand, Silt, Clay)	Bouyoucos (1927)	Bouyoucos hydrometer	Percentage (%)
4.	Pore space	Muthuval <i>et al.</i> (1992)	Graduated measuring cylinder	Percentage (%)
5.	Water Holding capacity	Muthuval <i>et al.</i> (1992)	Graduated measuring cylinder	Percentage (%)
CHEMICAL PROPERTIES				
1.	Soil pH (1:2.5)	Jackson (1958)	Digital pH meter	
2.	Electrical conductivity (1:2.5)	Wilcox (1950)	Digital conductivity meter	dS m ⁻¹
3.	Organic carbon	Walkley and Black (1947)	Wet oxidation method	Percentage (%)
4.	Available nitrogen	Subbiah and Asija (1956)	Soil alkaline permanganate method	kg ha ⁻¹
5.	Available phosphorus	Olsen <i>et al.</i> (1954)	Photometric colorimeter method	kg ha ⁻¹

Commented [a3]: Please change this section to be more systematic. It is important for further studies that require the replication of this methods.

6.	Available potassium	Toth and Prince (1949)	Flame photometric method	kg ha ⁻¹
----	---------------------	------------------------	--------------------------	---------------------

UNDER PEER REVIEW

RESULT AND DISCUSSION

Physical Properties

The textural classification of soil in different villages Mirzapur block. The texture classification of soil samples was shown **Sandy loam** in all villages.

Soil Bulk Density (Mg m^{-3})

The bulk density in a soil ranges from 1.17 -1.48 Mg m^{-3} . V_6 – Indi Parwatpur was reported as highest bulk density *i.e.*, 1.48 (Mg m^{-3}) while V_3 Amirati was reported as lowest bulk density *i.e.* 1.17 (Mg m^{-3}). Significant results were observed by (Ghosh *et al.*, 2021).

Soil Particle Density (Mg m^{-3})

Soil particle density ranges from 2.22– 2.55 Mg m^{-3} . V_1 – Aksauli was reported as highest particle density *i.e.* 2.55 (Mg m^{-3}), and lowest was found in V_3 - Amirati *i.e.* 2.22 (Mg m^{-3}). Similar results were reported by (Ghosh *et al.*, 2021)

Soil Porosity (%)

Soil porosity in a soil sample ranges from 40.61 – 48.00 %. V_1 -Aksauli reported as highest porosity of 48.00 % while V_6 – Indi Parwatpur reported as lowest porosity *i.e.* 40.61%. Significant results were observed by (Ghosh *et al.*, 2021)

Water Holding Capacity (%)

The values of water holding capacity of soil ranges from 39.37– 45.17%. Highest water holding capacity was observed in the V_1 - Aksauli which was 45.17% ,whereas the lowest water holding capacity was observed in V_6 – Indi Parwatpur *i.e.* 39.37% . Similar results were reported by (Ghosh *et al.*, 2021).

Table 2. Soil physical properties

	bulk density(Mg m^{-3})			particle density(Mg m^{-3})			Pore space(%)			water holding capacity(%)		
	0-15 cm	15-30 cm	15-45 cm	0-15 cm	15-30 cm	15-45 cm	0-15 cm	15-30 cm	15-45 cm	0-15 cm	15-30 cm	15-45 cm
V_1	1.29	1.31	1.35	2.48	2.52	2.55	48.00	47.82	46.45	45.17	43.19	41.78
V_2	1.31	1.33	1.35	2.32	2.35	2.38	43.51	43.00	42.80	42.76	40.78	39.37
V_3	1.17	1.25	1.29	2.22	2.25	2.27	47.33	44.50	42.70	44.19	42.81	40.18
V_4	1.34	1.39	1.44	2.42	2.47	2.50	44.63	43.74	42.36	43.39	41.78	40.17
V_5	1.30	1.35	1.39	2.44	2.48	2.52	46.74	45.54	44.85	44.15	42.03	40.35
V_6	1.44	1.48	1.48	2.41	2.44	2.44	42.78	40.61	40.61	40.18	39.37	39.37
V_7	1.25	1.29	1.33	2.29	2.33	2.35	45.62	44.64	43.37	43.07	41.35	39.75
V_8	1.29	1.33	1.36	2.35	2.38	2.40	45.09	44.11	43.26	43.31	41.17	49.17
V_9	1.31	1.35	1.42	2.45	2.46	2.50	46.52	45.03	43.18	44.13	42.30	40.39
F- test	S	S	S	S	S	S	S	S	S	S	S	S
S.E.m. (\pm)	0.023	0.020	0.020	0.037	0.382	0.323	0.656	0.767	0.645	0.610	0.610	0.683
C. D. @ 5%	0.070	0.061	0.060	0.118	0.113	0.096	1.949	2.280	1.919	1.814	1.813	2.030

Commented [a4]: More discussion is required rather than the result of this study. What is the main output of this study and what is the benefit for land management

Commented [a5]: More discussion is required




Fig. 1. Bulk density and Particle density (Mg m⁻³)

Commented [a6]: More discussion is required based on this table




Fig. 2. Pore space and water holding capacity (%)

Soil Chemical Properties

Soil pH

pH of a soil samples ranges from 7.43 – 8.20. The highest pH value was observed in the V₈ - Leduki *i.e.* 8.20 . the lowest pH was found in V₂ – Dewahi *i.e.* 7.43. The results shown the pH is alkaline in nature . The pH is significant and appropriate for the nutrient availability. Similar significant results were reported by (Ghosh *et al.*, 2021).

Soil EC (dS m⁻¹)

EC in a soil samples ranged from 0.21-0.39 dS m⁻¹. The results were found to be significant. Highest EC content was reported in V₉- Sirsi *i.e.* 0.39 dS m⁻¹ , whereas the lowest EC content was observed in V₃–Amirati *i.e.* 0.21 dS m⁻¹. Similar results were reported by (Ghosh *et al.*, 2021).

4.7 Soil Organic Carbon (%)

Organic carbon soil samples value ranges from 0.21 – 0.38 % The results were found to be significant. Highest organic carbon reported in V₂- Dewahi *i.e.* 0.38 % , whereas the lowest organic carbon was observed in V₃– Amirati *i.e.* 0.21%. The organic carbon is low as there is less vegetation is used as residue and due to tropical climate, degradation is quick in whole block. Similar results were reported by (Ghosh *et al.*, 2021).

Table 3. Variation in Soil pH, electric conductivity and organic carbon

S. No.	Soil pH			Soil EC (dS m ⁻¹)			Soil organic Carbon (%)		
	0-15 cm	15-30 cm	15-45 cm	0-15 cm	15-30 cm	15-45 cm	0-15 cm	15-30 cm	15-45 cm
V ₁	7.45	7.50	7.88	0.24	0.28	0.31	0.34	0.30	0.28
V ₂	7.43	7.63	7.76	0.26	0.29	0.32	0.38	0.36	0.33
V ₃	7.83	7.95	7.95	0.21	0.24	0.26	0.27	0.25	0.22
V ₄	8.03	8.15	8.19	0.33	0.35	0.37	0.28	0.26	0.21
V ₅	8.06	8.10	8.12	0.33	0.37	0.39	0.29	0.27	0.24
V ₆	8.07	8.13	8.16	0.31	0.35	0.35	0.32	0.30	0.28
V ₇	8.04	8.10	8.12	0.34	0.35	0.37	0.38	0.36	0.32
V ₈	8.15	8.18	8.20	0.22	0.25	0.27	0.33	0.30	0.27
V ₉	8.10	8.15	8.15	0.33	0.35	0.39	0.31	0.27	0.24
F- test	S	S	S	S	S	S	S	S	S
S.Em. (±)	0.134	0.108	0.083	0.006	0.004	0.004	0.002	0.004	0.004
C. D. @ 5 %	0.399	0.322	0.234	0.018	0.014	0.14	0.008	0.013	0.012

Commented [a7]: More discussion is required based on this information




Fig. 3. pH, EC (dS m⁻¹) and Organic Carbon (%)

Soil Nitrogen (kg ha⁻¹)

Nitrogen content in soil sample ranges from 160 – 380 kg ha⁻¹. The results were found to be significant. Highest nitrogen content in soil was observed in V₅- Birshahpur *i.e.*, 380kg ha⁻¹ and the lowest nitrogen content was reported in V₁- Aksauli *i.e.* 160 kg ha⁻¹. Similar results were observed (Ghosh *et al.*, 2021).

Soil Phosphorus (kg ha⁻¹)


Phosphorus in soil samples ranged from 22.53 – 37.49 kg ha⁻¹. The results were found to be significant. Highest phosphorus was reported in V₉– Ghorī *i.e.* 37.49 kg ha⁻¹, whereas the lowest phosphorus content was observed in V₄- Bhūli *i.e.* 22.53 kg ha⁻¹. Significant results were observed (Ghosh *et al.*, 2021).

Soil Potassium (kg ha⁻¹)

Potassium content in the soil samples ranges from 105.54 – 282.4kg ha⁻¹. The results were found to be significant. Highest potassium content was reported in V₆–Indī parwatpur *i.e.* 282.4 kg ha⁻¹, whereas the lowest potassium content was observed in V₈– Leduki *i.e.* 105.54 kg ha⁻¹. Similar results were observed with (Ghosh *et al.*, 2021).

Table 4. Variation in soil nitrogen, phosphorous and potassium

UNDER PEER REVIEW



S. No.	Soil Nitrogen (kg ha ⁻¹)			Soil Phosphorus (kg ha ⁻¹)			Soil Potassium (kg ha ⁻¹)		
	0-15 cm	15-30 cm	15-45 cm	0-15 cm	15-30 cm	15-45 cm	0-15 cm	15-30 cm	15-45 cm
V ₁	250	205	160	26.22	25.50	24.96	200	180.47	150.32
V ₂	344	285	200	24.22	23.33	22.82	156.8	148.32	122.22
V ₃	297	232	170	25.95	24.82	24.23	190.4	172.25	140.22
V ₄	338	265	195	24.45	23.08	22.53	170	150.45	125.25
V ₅	380	320	275	26.76	25.79	24.70	244	222.44	180.44
V ₆	350	280	220	26.60	25.64	24.89	282.4	260.22	220
V ₇	275	220	160	35.75	34.88	34.33	156.8	140.56	110.55
V ₈	280	250	190	36.71	35.20	34.59	148.43	130.44	105.54
V ₉	290	210	140	37.49	34.44	32.44	222	185.55	135.52
F- test	S	S	S	S	S	S	S	S	S
S.Em. (±)	5.295	2.823	2.622	0.403	0.401	0.534	3.540	2.743	2.082
C. D. @ 5 %	15.732	8.388	7.792	1.198	1.193	1.587	10.519	8.151	6.186




Fig. 4. Available nitrogen, available phosphorous and available potassium (kg ha⁻¹)

Commented [a8]: More discussion is required. What is the implementation for agricultural practices and land management

CONCLUSION

It can be concluded that the soils of Mirzapur block of Uttar Pradesh are in moderately good physical condition which favors the cultivation of most of the crops, especially rice and wheat. Soil texture showed high sand percentage, pH of soil is alkaline in nature, Electrical conductivity was suitable for all crops, Organic carbon was found low in all the villages, soils have medium Nitrogen in all villages. Phosphorus content was found medium in all villages.. Potassium content was found medium to high in all villages.. The deficiency of nutrients can be mitigated by the use of some inorganic fertilizers or organic fertilizers. Tolerant varieties can be used and Integrated Nutrient Management can be adopted.

REFERENCES

1. **Anonymous.** Munsell colour chart. Munsell colour company inc. 2241 N. calveri street, Baltimore, Maryland 21212, USA; 1971.

Bouyoucos, G.J. (1927) The hydrometer as new method of mechanical analysis of soil. *Soil Science*.23:343-353.

Das, R., Kumar , M., Singh, S. K., Jha, S., and. Sahoo, S. (2021). Soil Physico-chemical properties as affected by Longterm Application of Organic and Inorganic NPK Fertilisers under Rice-wheat Cropping System in Calcareous Soil of Bihar. *International Journal of Environment and Climate Change*, 100-107

Ghosh, A. K., Yadav, S., Singh, P., & Thakur, A. (2021). Characterization and Fertility Assessment of Soils of Mirzapur District of Eastern Uttar Pradesh for Sustainable Land Use Planning. *Journal of the Indian Society of Soil Science*, 69(4), 347-353

Hartemink, A. E., McBratney, A. B. and Mendonça-Santos, M. L. (2016) Digital soil mapping. In Soil mapping and process modeling for sustainable land use management (3- 11). Springer, Cham

Jackson. M. L. (1958). Soil Chemical Analysis. *Prentice Hall, Inc. Englewood.cliffe.N.J.*

Joffe, J. S. (1949). *Pedology* (Vol. 68, No. 4, p. 346). LWW.

John, M. S, Mohammad, Y., Ullahshah, H. and Bat, J. (2010). *Soil health*. 42(3): 1909-1922.

Olsen. S.R., Cole. C.V., Watanabe. F.S. and Dean. L.A. (1954). Estimation Of Available Phosphorus in Soils by Extraction with Sodium Bicarbonate. *U. S. Department of Agriculture, Circular No. 939.*

Muthuvel P, Udayasoorian C, Natesan R, Ramaswami PR(1992). Introduction to soil analysis. Tamil Nadu Agricultural University Coimbatore; 1992

Phogat, V. K., Tomar, V. S., and Dahiya, R. I. T. A. (2015). Soil physical properties. *Soil science: An introduction*, 135-171.

Tale, K. S., and Ingole, S. (2015). A review on role of physico-chemical properties in soil quality. *Chemical Science Review and Letters*, 4(13), 57-66.

Toth, S.J and Prince, A.L. (1949) Estimation of Cation Exchange Capacity and Exchangeable Ca, K, Na Content of Soil by Flame Photometer Technique. *Soil Sci.*67, 439-445.

Walkley, A. and Black, (1947) An examination of the Degtjareff method for determining soil organic matter and a proposed modification of the chromic acid titration method. *Soil Sci.*, 37:29-38

UNDER PEER REVIEW