

## Original Research Article

# EFFECT OF CROP ESTABLISHMENT METHODS AND PRECISION NUTRIENT MANAGEMENT ON GROWTH, YIELD ATTRIBUTES AND YIELD OF RICE UNDER RICE-WHEAT SYSTEM

### ABSTRACT:

Rice-wheat system is the major contributor in food security of India. The significance of the system is well addressed by time to time but, excessive use of resources, stagnation in yield, environment deterioration including erosion and nutrient mining are the greater challenge it possess. To address the issue a study was carried out at Banaras Hindu University's Agricultural Research Farm in Varanasi during the *Kharif* and *Rabi* seasons of 2019-20 and 2020-21, respectively. The experiment plotted in a split plot statistical design with three replications and four crop establishment methods: CE<sub>1</sub>: Conventional till rice (puddled transplanted)-Conventional till wheat, CE<sub>2</sub>: Conventional till direct seeded rice (DSR)-Conventional till wheat, CE<sub>3</sub>: Conventional till DSR-Zero-till wheat (rice residue retained), and CE<sub>4</sub>: Zero-till DSR-Zero-till wheat (residue retention in rice and wheat) and CE<sub>4</sub>: ZTDSR-ZTW (Anchored rice and wheat residue retained) crop establishment method among the main plot treatments and RWCM-based nutrient recommendation among the sub plot treatments replicated thrice. CE<sub>4</sub> treatment among main plot and N<sub>3</sub> among nutrient management practices produced higher values of plant height and number of tillers at 30, 60, 90 days after sowing and at harvest, as well as yield attributes *i.e.* maximum number of panicles m<sup>2</sup> and grains per panicle, than the other treatments. The same treatments also resulted in increased rice grain yield. It may be inferred that conservation agriculture-based crop establishment *i.e.* ZTDSR-ZTW with residue retention of rice and wheat crops and RWCM-based nutrient application, may be favourable for improved growth, yield attributes and yield in the rice crop under the region of eastern Uttar Pradesh.

*Keywords: ZTDSR, conservation agriculture, residue retention, rice growth*

### 1. INTRODUCTION:

The rice-wheat system contributed maximum calorie in the food basket of the IGP region. Puddling has been shown to have detrimental effects on the soil environment, particularly on healthy microbes and soil aggregation (Jat *et al.*, 2014; Haque *et al.*, 2016). The drying of the soil and the emergence of cracked soil blocks impede the preparation of the land for the next upland crop. As a result, to prepare a healthy seedbed for the crop that follows rice, heavy tillage and irrigation are needed, which delays planting and ultimately lowers the yield of the dryland crop. The use of machine-driven processes including tillage, sowing/transplanting, harvesting, and threshing has become more pressing due to the growing shortage of human labour (Jat *et al.* 2013). According to Parihar *et al.* (2017), mechanisation and input-intensive agricultural practices have a negative influence on soil quality and environmental pollution. This highlights the need for alternative crop management strategies that could reduce energy use, safeguard the environment, and maintain crop productivity that is comparable to or even higher than current practices. Rice productivity in India has been stagnant and may decline in the future due to over-exploitation of natural resources (Ladha *et al.* 2009), low seed replacement rate, poor irrigation water, fertiliser, and crop residue management (Ladha *et al.* 2009), consistent cropping patterns over time, and a lack of awareness among farmers about the consequences of poor cultivation practices. According to several studies (Verhulst *et al.*, 2010; Kumar *et al.*, 2019; Gathala *et al.*, 2020b; Jat *et al.*, 2014), conservation agriculture (CA) can improve crop establishment and timely sowing, maintain or increase yield, reduce water and energy use, lower production costs and increase income, and improve soil quality. It can also increase system resilience. Rice seedlings are often put onto repeatedly tilled, puddled, and flooded fields in the standard manner of rice farming. However, it is also possible to transplant seedlings without puddling, which can reduce the cost of rice production overall as well as water, energy, labour, and other factors (Krupnik *et al.*, 2014; Haque *et al.*, 2016; Hossen *et al.*, 2018; Gathala *et al.*, 2021).

There was non-significant increase in fertilizer N efficiency in rice grown in different Asian countries during the past 30 years, the average plant recovery efficiency of fertilizer N in rice is still only about 30% (Dobermann, 2000; Dobermann and Cassman, 2002). Application of other macronutrients, such as K, has lagged behind leading to imbalanced plant nutrition and negative K input–output balances in many parts of Punjab and Asia. Environmental pollution by nutrient leaching or runoff from rice fields has become another concern across Asia (Dobermann *et al.*, 2002). The site specific nutrient management (SSNM) approach provides scientific principles for determining field-specific fertilizer nitrogen (N), phosphorus (P), and potassium (K) requirements for crops. The SSNM approach provides algorithms that can be used to determine field-specific fertilizer requirements matching the needs and conditions of individual farmers. Recent advances in information and communication technology (ICT) offer ample opportunities to use mobile phones to provide farmers with field-specific nutrient management recommendations calculated by decision making tools using algorithms based on SSNM. Use of SSNM based fertilizer recommendations were shown to increase yields, net income of farmers, and provide positive impacts on the environment when compared with existing fertilizer practices.

## **MATERIALS AND METHODS:**

The present investigation was conducted at the Agricultural Research Farm, Institute of Agricultural Sciences, Banaras Hindu University, Varanasi during *kharif* and *rabi* season of 2019-20 and 2020-21, in South-eastern part of Varanasi city of India (25°18'N latitude and 83°31'E longitude at an altitude of 128.93 meter above sea level). The experimental soil was Gangetic alluvial sandy clay loam with pH 7.22. It was moderately fertile-being low in available organic carbon (0.41%) as well as available nitrogen (213.59 kg ha<sup>-1</sup>) and medium in available phosphorus (23.23 kg ha<sup>-1</sup>) as well as available potassium (223.50 kg ha<sup>-1</sup>). The experiment was laid out in split-plot design replicated thrice with four crop establishment methods viz., CE<sub>1</sub>: Conventional till rice (puddled transplanted)-Conventional till wheat { farmers practice} [CT rice–CT wheat], CE<sub>2</sub>: Conventional till direct seeded rice-Conventional till wheat [CTDSR–CT wheat], CE<sub>3</sub>: Conventional till direct seeded rice-Zero-till wheat +rice residue retained [CTDSR–ZT wheat + R R], CE<sub>4</sub>: Zero-till direct seeded rice-Zero-till wheat + residue retention in rice and wheat [ZTDSR–ZT wheat + RW R] in main plots and three nutrient management practices viz., N<sub>1</sub>: Farmers Practices (FP), N<sub>2</sub>: Recommended fertilizer dose (RFD), N<sub>3</sub>: Rice-Wheat Crop Manager recommendation (RWCM) in sub plots during both the years. In nutrient management treatments, fertilizer applications to rice were done as per the treatment requirement. Half dose of N and full doses of P<sub>2</sub>O<sub>5</sub>, K<sub>2</sub>O and Zn were applied as basal before sowing and remaining half dose of N was top dressed in two equal splits at active tillering and panicle initiation stage in rice in farmers practice and also in RFD as per requirement/recommendation. In SSNM, fertilizer was applied as per RWCM recommendation. The sources of fertilizers were urea, Di-ammonium phosphate (DAP), muriate of potash (MOP) and ZnSO<sub>4</sub>. The doses of fertiliser for N<sub>1</sub> were 164-50-32-4 kg N-P<sub>2</sub>O<sub>5</sub> –K<sub>2</sub>O –Zn ha<sup>-1</sup> whereas in N<sub>2</sub> it was 150-60-60-5 kg N-P<sub>2</sub>O<sub>5</sub> –K<sub>2</sub>O –Zn ha<sup>-1</sup> and in N<sub>3</sub>, the doses were based on RWCM based nutrient recommendations.

## **2. RESULT AND DISCUSSION**

### **2.1 GROWTH PARAMETERS:**

Data indicates that maximum plant height and number of tillers recorded at 30 60, 90 DAS/DAT (Days after sowing/transplanting) and at harvest (Table number 1) was achieved under the CA-based crop establishment method *i.e.* CE<sub>4</sub>: ZTDSR-ZT wheat + RW R over the rest of the crop establishment method *i.e.* CE<sub>3</sub>, CE<sub>2</sub>, and CE<sub>1</sub>, but CE<sub>4</sub> and CE<sub>3</sub> were found at par with each other in term of plant height at both the stages of observation among the main plot treatments, while under the sub plot treatments, N<sub>3</sub>: Rice wheat crop manager based nutrient recommendation observed maximum plant and higher number of tillers at both the stages, However, N<sub>3</sub> was observed on par with N<sub>2</sub>: Recommended fertiliser dosage (RFD) in terms of plant height. A similar pattern was seen throughout the second year of evaluation.

The reason for rice's better performance in terms of plant height and number of tillers in the afore mentioned treatments could be due to better nutrient availability, better organic carbon, which leads to improved microbial activity, and better moisture availability, all of which contribute to better initial crop growth under CA based system (Choudhary *et al.*, 2020). Optimised nutrient usage during SSNM may result in higher cell division, enlargement, photosynthesis, and protein synthesis, leading to quantitative improvements in plant growth

traits such as tillering (Singh *et al.*, 2015b). Similar results were also obtained by Shahi *et al.*, 2022.

**Table 1: Effect of crop establishment and nutrient management on plant height (cm) on Rice**

Treatments	30 DAS/DAT		60 DAS/DAT		90 DAS/DAT		At harvest	
	2019	2020	2019	2020	2019	2020	2019	2020
<b>Crop establishment (CE)</b>								
CE <sub>1</sub> : CT rice – CT wheat	25.57	26.33	52.16	54.53	82.22	81.92	79.15	82.56
CE <sub>2</sub> : CTDSR – CT wheat	27.29	28.03	56.96	57.79	84.66	86.23	82.44	84.24
CE <sub>3</sub> : CTDSR – ZT wheat	27.61	28.38	58.06	58.94	85.45	87.56	84.02	86.15
CE <sub>4</sub> : ZT rice – ZT wheat	28.95	29.77	61.37	64.30	89.11	91.18	87.63	91.02
Sem ±	0.52	0.51	0.71	0.68	1.09	1.04	0.98	0.94
CD (P=0.05)	1.55	1.49	2.29	2.17	3.29	3.17	3.11	2.98
<b>Nutrient management (N)</b>								
N <sub>1</sub> : Farmers Practices	26.27	26.85	53.85	54.76	82.09	82.94	80.45	81.73
N <sub>2</sub> : Recommendation Fertilizer Dose	27.39	28.33	57.88	58.40	85.72	87.07	83.71	86.62
N <sub>3</sub> : RWCM Recommendation	28.41	29.20	59.70	60.51	88.26	90.15	85.76	89.63
Sem ±	0.40	0.37	0.61	0.57	0.98	0.93	0.87	0.82
CD (P=0.05)	1.20	1.12	1.94	1.79	2.85	2.73	2.65	2.51
<b>Interaction</b>	NS	NS	NS	NS	NS	NS	NS	NS

**Table 2: Effect of crop establishment and nutrient management on number of tillers (m<sup>2</sup>) on Rice**

Treatments	30 DAS/DAT		60 DAS/DAT		90 DAS/DAT		At harvest	
	2019	2020	2019	2020	2019	2020	2019	2020
<b>Crop establishment (CE)</b>								
CE <sub>1</sub> : CT rice – CT wheat	139.70	143.28	350.89	359.35	332.75	339.57	301.89	307.89
CE <sub>2</sub> : CTDSR – CT wheat	228.61	232.73	365.01	375.56	350.47	354.75	322.54	325.92
CE <sub>3</sub> : CTDSR – ZT wheat	232.70	237.77	370.82	378.09	353.77	358.90	325.87	329.24
CE <sub>4</sub> : ZT rice – ZT wheat	242.04	247.08	381.31	387.98	367.81	370.69	337.79	343.87
Sem ±	1.62	1.59	3.39	3.32	3.21	3.16	3.09	3.02
CD (P=0.05)	4.82	4.65	10.23	9.98	9.79	9.62	9.52	9.39
<b>Nutrient management (N)</b>								
N <sub>1</sub> : Farmers Practices	202.39	205.67	359.03	367.29	342.01	346.79	315.24	319.37
N <sub>2</sub> : Recommendation Fertilizer Dose	211.42	216.54	363.90	375.25	350.94	355.72	321.71	325.95
N <sub>3</sub> : RWCM Recommendation	218.47	223.43	378.10	383.21	360.66	365.44	329.11	334.87
Sem ±	1.45	1.42	3.08	3.01	2.91	2.87	2.80	2.74
CD (P=0.05)	4.42	4.27	9.40	9.17	9.00	8.84	8.75	8.63
<b>Interaction</b>	NS	NS	NS	NS	NS	NS	NS	NS

## 2.2 YIELD ATTRIBUTES AND YIELD

Table number 2 presents' data on the number of panicles per m<sup>2</sup> and the number of grains per panicle and grain yield. The number of panicles and number of grains were recorded significantly higher under the CE<sub>4</sub> crop establishment technique (ZTDSR-ZT wheat + RW R), followed by CE<sub>3</sub>, CE<sub>2</sub>, and CE<sub>1</sub>. Among the nutrient management approaches, site-specific nutrient management, *i.e.* RWCM-based nutrient recommendation (N<sub>3</sub>), observed the highest number of panicles and per m<sup>2</sup> number of grains per panicle compared to other nutrient management practices. N<sub>3</sub> were found at par with N<sub>2</sub> in terms of number of panicles in first year only. A similar pattern was seen in second year of testing.

The rice production was significantly influenced by various crop establishment methods and nutrient management strategies used throughout both years of study. The ZTDSR-ZTW + RW R treatment produced significantly greater yield than the other main plot treatments *i.e.* crop establishment methods. In the sub plot treatment, site specific nutrient management, *i.e.* nutrient prescription based on RWCM, resulted in a significantly higher yield than the other sub plot treatments but RFD was found at par with N<sub>3</sub>. The same pattern was found in the second year. The ZT residue retention plots (CE<sub>4</sub>) improved rice crop establishment and light interception, resulting in more effective use of integrated inputs which resulted in better yield attributes and grain yield (Choudhary *et al.*, 2020). Improved nutrient delivery throughout rice's active growth and development phases may be the cause of the rise in yield-contributing features under SSNM-based treatments (Singh *et al.* 2014). Similar results were observed by Shahi *et al.*, 2022.

**Table 3: Effect of crop establishment and nutrient management on yield attributes and grain yield of Rice**

Treatments	Panicles (m <sup>-2</sup> )		Grains panicle <sup>-1</sup>		Grain yield (Quintal ha <sup>-1</sup> )	
	2019	2020	2019	2020	2019	2020
<b>Crop establishment (CE)</b>						
CE <sub>1</sub> : CT rice – CT wheat	234.96	239.33	107.64	110.14	43.21	43.76
CE <sub>2</sub> : CTDSR – CT wheat	253.87	260.46	115.33	118.90	44.93	45.55
CE <sub>3</sub> : CTDSR – ZT wheat	260.91	268.28	118.87	124.08	46.28	46.96
CE <sub>4</sub> : ZT rice – ZT wheat	279.58	288.85	126.88	132.78	48.62	49.32
Sem ±	3.89	3.84	1.93	1.91	0.76	0.77
CD (P=0.05)	11.97	11.79	5.82	5.77	2.29	2.31
<b>Nutrient management (N)</b>						
N <sub>1</sub> : Farmers Practices	248.64	252.21	110.78	114.19	44.06	44.65
N <sub>2</sub> : Recommendation Fertilizer Dose	256.32	263.35	116.64	120.70	45.52	46.16
N <sub>3</sub> : RWCM Recommendation	267.02	277.14	124.13	129.45	47.70	48.39
Sem ±	3.57	3.49	1.84	1.81	0.76	0.75
CD (P=0.05)	11.09	10.77	5.59	5.47	2.29	2.33
<b>Interaction</b>	NS	NS	NS	NS	NS	NS

## CONCLUSION:

The Conservation Agriculture (CA)-based crop establishment method, *i.e.* ZTDSR-ZT Wheat + standing residue of rice and wheat crop with site-specific nutrient management based on Rice-Wheat Crop Manager recommendation (N<sub>3</sub>), produced better growth parameters, *i.e.* maximum plant height and number of tillers, better yield attributes, *i.e.* maximum number of panicles per m<sup>2</sup> and maximum number of grains per panicle, and higher yield than the other main and sub plot treatments. ZTDSR-ZTW + RW R crop setup, along with RWCM-based nutrition recommendations, can improve rice growth and yield in eastern Uttar Pradesh's rice-wheat system.

## References:

Jat RK, Sapkota TB, Singh RG, Jat ML, Kumar M, Gupta RK. Seven years of conservation agriculture in a rice–wheat rotation of Eastern Gangetic Plains of South Asia: yield trends and economic profitability. *Field Crops Research*. 2014 Aug 1;164:199-210.

Haque ME, Bell RW, Islam MA, Rahman MA. Minimum tillage unpuddled transplanting: An alternative crop establishment strategy for rice in conservation agriculture cropping systems. *Field crops research*. 2016 Jan 1;185:31-9.

Jat ML, Gathala MK, Saharawat YS, Tetarwal JP, Gupta R. Double no-till and permanent raised beds in maize–wheat rotation of north-western Indo-Gangetic plains of India: Effects on crop yields, water productivity, profitability and soil physical properties. *Field Crops Research*. 2013 Aug 1;149:291-9.

Parihar CM, Jat SL, Singh AK, Ghosh A, Rathore NS, Kumar B, Pradhan S, Majumdar K, Satyanarayana T, Jat ML, Saharawat YS. Effects of precision conservation agriculture in a maize-

wheat-mungbean rotation on crop yield, water-use and radiation conversion under a semiarid agro-ecosystem. *Agricultural Water Management*. 2017 Oct 1;192:306-19.

Jat ML, Gathala MK, Ladha JK, Saharawat YS, Jat AS, Kumar V, Sharma SK, Kumar V, Gupta R. Evaluation of precision land leveling and double zero-till systems in the rice–wheat rotation: Water use, productivity, profitability and soil physical properties. *Soil and Tillage Research*. 2009 Sep 1;105(1):112-21.

Ladha JK, Singh Y. Integrated crop and resource management in the rice-wheat system of South Asia. *Int. Rice Res. Inst.*; 2009.

Verhulst N, Govaerts B, Verachtert E, Castellanos-Navarrete A, Mezzalama M, Wall P, Deckers J, Sayre KD. Conservation agriculture, improving soil quality for sustainable production systems. *Advances in soil science: food security and soil quality*. 2010 Jun 23;1799267585:137-208.

Kumar V, Gathala MK, Saharawat YS, Parihar CM, Kumar R, Kumar R, Jat ML, Jat AS, Mahala DM, Kumar L, Nayak HS. Impact of tillage and crop establishment methods on crop yields, profitability and soil physical properties in rice–wheat system of Indo-Gangetic Plains of India. *Soil Use and Management*. 2019 Jun;35(2):303-13.

Gathala MK, Laing AM, Tiwari TP, Timsina J, Islam MS, Chowdhury AK, Chattopadhyay C, Singh AK, Bhatt BP, Shrestha R, Barma NC. Enabling smallholder farmers to sustainably improve their food, energy and water nexus while achieving environmental and economic benefits. *Renewable and Sustainable Energy Reviews*. 2020 Mar 1;120:109645.

Krupnik, T.J., Yasmin, S., Shahjahan, M., McDonald, A., Hossain, K., Baksh, E., Hossain, F., Kurishi, A.S.M.A., Miah, A.A., Mamun, M.A. and Rahman, B.M.S., 2014, June. Productivity and farmers' perceptions of rice-maize system performance under conservation agriculture, mixed and full tillage, and farmers' practices in rainfed and water-limited environments of southern Bangladesh. In *6th World Congress on Conservation Agriculture*. Winnipeg, Canada. June (Vol. 24).

Hossen MA, Hossain MM, Haque ME, Bell RW. Transplanting into non-puddled soils with a small-scale mechanical transplanter reduced fuel, labour and irrigation water requirements for rice (*Oryza sativa* L.) establishment and increased yield. *Field Crops Research*. 2018 Aug 1;225:141-51.

Gathala MK, Laing AM, Tiwari TP, Timsina J, Rola-Rubzen F, Islam S, Maharjan S, Brown PR, Das KK, Pradhan K, Chowdhury AK. Improving smallholder farmers' gross margins and labor-use efficiency across a range of cropping systems in the Eastern Gangetic Plains. *World Development*. 2021 Feb 1;138:105266.

Dobermann A. Future intensification of irrigated rice systems. In *Studies in Plant Science* 2000 Jan 1 (Vol. 7, pp. 229-247). Elsevier.

Dobermann A, Cassman KG. Plant nutrient management for enhanced productivity in intensive grain production systems of the United States and Asia. *Plant and soil*. 2002 Nov;247:153-75.

Choudhary M, Jat HS, Datta A, Sharma PC, Rajashekar B, Jat ML. Topsoil bacterial community changes and nutrient dynamics under cereal based climate-smart agri-food systems. *Frontiers in Microbiology*. 2020 Jul 28;11:542545.

Singh VK, Shukla AK, Singh MP, Majumdar K, Mishra RP, Rani M, Singh SK. Effect of site-specific nutrient management on yield, profit and apparent nutrient balance under pre-dominant cropping systems of Upper Gangetic Plains. ICAR.

Shahi UP, Singh VK, Kumar A, Upadhyay PK, Rai PK. Site-specific nutrient management: impact on productivity, nutrient uptake and economics of rice-wheat system. *Indian J. Agric. Sci*. 2022 Feb 1;92:195-8.

Singh VK, Dwivedi BS, Tiwari KN, Majumdar K, Rani M, Singh SK, Timsina J. Optimizing nutrient management strategies for rice–wheat system in the Indo-Gangetic Plains of India and adjacent region for higher productivity, nutrient use efficiency and profits. *Field Crops Research*. 2014 Aug 1;164:30-44.