

Review Article

Effect of zero tillage on soil properties, weed dynamics and performance of chickpea under rice – chickpea cropping system: A review

Abstract

Farming is a critical segment in many countries of the world. Zero tillage systems are sustainable farming practices that involve sowing seeds without disturbing the soil through tillage. Instead of ploughing, this system leaves the previous year's crop residues on the soil surface and plant directly into the undisturbed soil. This approach helps to maintain the soil structure, reduce soil erosion, increase water retention, and improve soil health. This sustainable farming practices has several benefits for farmers, including improved soil health, reduced soil erosion, increased crop yields, and environmental sustainability. This system helps to reduce soil erosion by minimizing soil disturbance, which prevents soil particles from being exposed to the wind and water. Also helps to retain soil moisture, which reduces the need for irrigation, and improves soil structure, which enhances soil fertility. Zero-tillage systems can have reduced labor and fuel costs associated with conventional tillage practices. Systems require less fuel and labor compared to conventional tillage practices, which can save farmers money and reduce their carbon footprint. This is because these systems improve soil fertility by increasing soil organic matter, which enhances soil nutrient availability. Moreover, zero tillage systems can increase water retention in the soil, which can improve crop growth and yield. By reducing soil erosion, conserving soil moisture, and improving soil health, these systems help to promote environmental sustainability. Also reduce greenhouse gas emissions by reducing fuel usage and sequester carbon in the soil through the accumulation of crop residues. Therefore, farmers should adopt zero-tillage systems to enhance their productivity, reduce costs, and promote environmental sustainability.

Key words: Zero tillage, soil properties, weed dynamics, rice – chickpea system, benefits

Introduction

Chickpea (*Cicer arietinum* L.) is grown in the winter season in sequence with different crops like rice, maize, soybean, sorghum and ~~pearlmillet~~ pearl millet in the states of Madhya Pradesh, Maharashtra, Rajasthan, Andhra Pradesh, Tamil Nadu, Karnataka, Uttar Pradesh and Gujarat (Adarshet al., 2019). Rice- chickpea cropping system (RCCS) is predominantly cultivated in the northern part of the country ~~next to~~ after rice-wheat cropping system (Arya et al., 2005). ~~Beside~~ Besides wheat, the introduction of chickpea in a ~~rice-based~~ rice-based cropping system is a feasible option to ensure sustainable food production and maintain environmental integrity. It can fix N up to 140 kg ha⁻¹ in a growing period (Poonia and Pithia, 2013) and helps in

Commented [KA1]: Can rewrite the sentences as its merely a repetition

eradicating obnoxious weeds like wild oat (*Avena fatua*) and canary grass (*Phalaris minor*) (Ali and Basu, 1992). Chickpea, sown during rabi season after harvesting of the rice, ~~it is~~ considered as a most valuable and drought tolerant crop (Gangwar and Singh *et al.*, 2010). Present agriculture has globally been facing major challenges including soil erosion which drastically lowers the crop yield. Chickpea can increase the productivity both in terms of N saving from fertilizer sources and build up soil fertility through biological source of N (Banjara *et al.*, 2017). In rice-chickpea cropping system (RCCS), the sowing of chickpea often delayed until last November or early December either due to late harvesting of rice or more time required to seedbed preparation for chickpea seeding. Late sowing put negative effects on germination, seedling establishment, reproductive stage due to the less soil moisture and low temperature. Conventional tillage (CT) affect the sustainable resources through its influence on soil properties, crop growth and the use of excessive and un-necessary tillage operations is often harmful to soil (Nazeer *et al.*, 2012). Conservation agriculture (CA) techniques involve zero tillage (ZT) which reduces risk of late sowing of chickpea. Zero tillage seeding of chickpea helps in timely sowing, because it facilitates direct sowing in previous unprepared crop ~~field~~fields. ZT also ~~reduce~~reduces the negative environmental effects of agriculture such as soil erosion and degradation of physical properties of soil leading to decrease crop productivity (Monneveux *et al.*, 2006).

Commented [KA2]: Not making proper sense

Commented [KA3]: Please rewrite for clarity

Chickpeas work perfectly ~~into~~in a lot of different cropping schemes. The main cropping sequences that are followed in India's various climatic circumstances include pearl millet/sorghum-chickpea, rice/maize-chickpea, and cotton-chickpea (AICRP-Sorghum, 2017). One of the main agricultural systems in the Bundelkhand region is fodder sorghum-chickpea (Singh and Praharaj, 2020). Similarly, Andhra Pradesh and Karnataka follow the rice-chickpea, maize/sorghum-chickpea, and pearl millet-chickpea systems. These cereal-legume cropping systems aid in interrupting monocropping sequences, conserving nitrogen, halting soil erosion, and enhancing soil health. ~~This Present~~ review assesses the impact of zero tillage on performance of chickpea as well as impact on weed dynamics within the context of rice-chickpea cropping systems.

1. What is zero tillage

Zero tillage has emerged as a promising approach in sustainable agriculture, offering benefits such as improved soil health and reduced erosion. One of the conservation tillage techniques is zero-tillage systems, where soil disturbance is limited to sowing activities. It keeps crop leftovers on the soil surface, which raises the amount of soil organic carbon and

Commented [KA4]: Add recent reference

enhances soil quality and health. In comparison to conventional tillage, zero tillage systems typically needed less maintenance and provided higher economic returns (Smart and Bradford, 1999). Zero tillage reduces cost by 3.8 % to 13.7 % and frees up at least 8 days for the growth of succeeding crops. Zero tillage farming is also an option for low-income farmers (Quddus et al., 2020)

Commented [KA5]: Recent reference

Zero tillage globally advocated to its benefits. Zero tillage in rice-based system will not only enable the earlier planting of chickpea, also reduce the land preparation cost, avert the water logging by just utilizing the residual moisture, but also it will let the weeds stay dormant below the soil for their lack of exposure to light.

Commented [KA6]: Please rewrite

2. Effect of zero tillage

Mishra et al. (2012) reported that zero tillage in chickpea had significant impact on crop growth and yield parameters such as plant population, number of branches, height of plants, number of pods per plant etc. when compared with conventional tillage. This may have been due to improved moisture conditions near the soil surface or greater seed-soil contact in ZT systems (Miller et al., 2002). Tillage systems affect soil disturbance, weed management, and weed seed production. A change in tillage systems will influence the species composition and vertical distribution of weed seeds in agricultural soils (Bhular et al., 1995).

According to Hemmat et al. (2004), the grain production of chickpea at the Dryland Agricultural Research Station in Iran was higher under the minimum tillage-sweep ploughing, reduced tillage-tandem disk, and zero tillage techniques, respectively, than under the conventional tillage-mould board method.

2.1 Effect on growth and yield attributes of chickpea

In a field experiment (Quddus et al., 2020) observed significant variation in growth and yield attributes of chickpea due to different tillage practices except 100-seed weight. They obtained tallest plant (48.9 cm and 50.1 cm) under zero tillage (T₁) treatment as compared to the rest of treatments, they stated that tallest plant under zero tillage condition observed might be due to availability of optimum soil moisture in soil for available nutrients to plant uptake. Performance of zero tillage and minimum tillage both achieved well and produced taller plant, maximum number of branches per plant, and the highest number of pods per plant. The highest number of pods plant⁻¹ was realized in zero tillage which is related with more branching (Banjara et al., 2017). The climatic condition of zero tillage plots might be favoured favourable for the growth and development of chickpea plant. They stated that zero tillage system successfully adopts the weather conditions in the growing season. In an experiment,

Commented [KA7]: Pls rewrite for clarity

Commented [KA8]: Pls rewrite

Commented [KA9R8]:

carried out in ICARDA (Syria), tillage, residue, and weed control techniques had a significant impact on the growth and yield performance of chickpea. Compared to conventional tillage, zero-till seeded chickpea had a 26.3% greater grain yield (Piggin *et al.*, 2015).

3. Effect of Zero Tillage on Soil Properties:

Zero tillage have impact on physical, chemical and biological properties of the soil (Bhatt, 2017). No-till in the context of CA can also lead to improvements in soil quality by improving soil structure and enhancing soil biological activity, nutrient cycling, soil water holding capacity, water infiltration and water use efficiency (Hobbs *et al.*, 2008).

3.1 Soil Physical Properties:

3.1.1 Effect on bulk density- Bulk density significantly influenced ($P \leq 0.05$) by the different tillage practice. Under zero tillage and minimum tillage practices lower value of bulk density (1.36 and 1.39) observed by several researchers (Lal *et al.*, 1989; Karlen *et al.*, 1994; Green *et al.*, 2005; Jat *et al.*, 2006). Under conventional tillage the bulk density was 1.39 Mgm^{-3} . They explained that ~~this~~ the increase was the result of natural reconsolidation of soil particles because of subsequent irrigation and summer drying. ~~The gradual increase the disturbances in the soil by increased number of tillage and intercultural operations resulting more compaction of soil that leads to increased bulk density.~~ The increase in bulk density under tilled soils may be due to increase in non-capillary porosity and low soil mass per unit volume. Owing to the progressive increase in bulk density after tillage, the difference between the tilled and no tilled treatments becomes smaller and smaller with the time since tillage progresses. The no tillage system maintains a significantly greater amount of residue on the soil surface, increases soil organic carbon and biotic activity, thereby decreasing bulk density, particularly near the soil surface.

Commented [KA10]: units

Commented [KA11]: Pls rewrite

Commented [KA12]: Add Reference

3.1.2. Effect on water stable aggregate (WSA): Reduced tillage practices ~~showing~~ showed the higher WSA compared ~~with to~~ the intensive tillage and more intercultural operations. The highest WSA was recorded in Zero tillage (59.32%) followed by under minimum tillage (59.22%). In an experiment, Hula *et al.* (2010), observed worsened soil structure under conventional tillage in comparison to reduced tillage. It might be due to ~~that~~ soil tillage, ~~based on ploughing can that~~ result in faster deterioration of soil structure compared to shallower tillage which ~~less disturbing the soil~~ minimizes soil disturbance. According to various authors (Daraghme *et al.*, 2009; Boguzas *et al.*, 2010), site adaptable tillage, in comparison to conventional methods, increases the amount of water stable aggregates and improves ~~(soil)~~ structure due to a combination of greater amounts of organic matter, reduced bulk weight of

soil, and a greater share of larger aggregates. Enhanced in physical parameters of soil resulting in increased porosity and water stable aggregate hence more available water content.

Commented [KA13]: Pls rewrite

3.1.3. Effect on soil aggregation: Soil aggregation is an important physical property and is affected by divergent tillage methods. According to (Mannering *et al.*, (1975) and (Edwards *et al.*, (1988), soil aggregation decreased in CT plots as tillage break-broke down the aggregates. Aggregation was highest in the 0-0.05 m layer of ZT plots. Long-term Long-term adoption of the ZT will certainly improve the aggregate stability of the topsoil (Douglas and Goss, (1982). (Lal *et al.*, (1989) reported that aggregate size tended to be around 22% higher under ZT treatments in comparison to that of tilled plots. Soil under ZT have better aggregates, aggregate stability, increased porosity which further improved rhizosphere environment for the better plant growth while intensive tillage led to decline in soil organic matter through accelerated oxidation of the organic matter (Ghuman and Sur, 2001); (Francis and Knight, 1993); (Martino and Shaykewich, 1994); (Ghosh *et al.*, (2010). Zero tillage can improve macro-aggregation (>0.25 mm) and mean weight diameter which further improved-improves the carbon sequestration potential of the soil (Franzluebbers and Arshad, 1996; McConkey *et al.*, 2003). Most studies coined that aggregation improved with the adoption of the ZT and ZT must be adopted for about 5-8 years (Bhatt, 2015; Bhatt and Kukal, 2015e).

Commented [KA14]: Chronological order of references

Formatted: Font: Bold

Formatted: Font: Bold

4. Effect on soil chemical properties

Tillage operations and soil disturbance generally cause an increase in soil aeration, residue decomposition; Organic N mineralization and availability of N for plant use (Dinnes *et al.*, 2002). (Moussa-Machraoui *et al.*, (2010) observed that some of the chemical parameters of soil were significantly modified under no-tillage (NT) when compared to conventional tillage systems. The nutrient (N, P, K, P₂O₅ and K₂O) contents were as more under no tillage than conventional tillage.

4.1. Effect on EC: Lower electrical conductivity of soil under the ZT system compared with CT pertains to the enhanced water movement in the soil and improved soil aggregate development. The inconsistent effect of zero tillage on soil electrical conductivity (EC) is observed and reported by (Singh and Singh (2014) and (Chatterjee and Lal, (2009). (Patni *et al.*, (1998) also reported a decrease in soil EC under NT which might be due to more downward movement of salts along with water infiltration into deeper layers.

4.2. Effect on soil pH: Long-term Long-term adoption of the ZT resulting-results in acidification of the surface soil which further affects the supply and distribution of other

nutrients within the rhizosphere. Under ZT, a significant lowering of pH was observed at the upper soil 0-7.5 cm on silt loam soil (**Dick *et al.*, 1986 and Tarkalson *et al.*, 2006**). Soil acidity with ZT was observed due to the decomposition of organic residues at the surface with subsequent leaching of resultant organic acids into mineral soil (**Blevins *et al.*, 1977**). While **Rahman *et al.* (2008)** observed ~~no significant~~ no significant differences in soil pH among ~~no tillage~~ no tillage and conventional tillage practices.

4.3. Effect on soil organic matter: Both quality and quantity of the organic matter (OM) in a particular soil ~~is-are~~ indicators of its quality as ~~they~~ they affect almost all the physico-chemical properties. Generally, OM content of upper soil under ZT is higher, than for tilled soil. The OM quantity will generally improve with conservation tillage, but remain fairly constant, or perhaps decrease further, with intensive tillage (**Frye *et al.*, 1985**). Improvement in soil OM status in the upper 0.2 m to 0.4 m soil depth under ZT was reported by **Freitas *et al.*, (1999)** and **Six *et al.*, (2002)**. The surface soil beneath the canopy had higher OM content in both tillage systems, particularly under ~~no tillage~~ no tillage. Contents of SOC ~~was-were~~ slightly greater under ~~no tillage~~ no tillage than under conventional tillage. ~~No tillage~~ No tillage ~~can increase~~ can increase soil C stock by an average of 4-7 ~~per cent~~ per cent%. The enhancement of SOC and SOM contents in the soil under ~~no tillage~~ no tillage is often accompanied by the enhancement of the cation exchange capacity (**Moussa-Machraoui *et al.*, 2010**). The SOC was significantly higher when stubble was left on the surface. Under conservation tillage the organic carbon increased by 11 ~~per cent~~ per cent (**Bricchi *et al.*, 2004**) and 14-17 ~~per cent~~ per cent (**Hazarika *et al.*, 2009**), while a reverse trend was observed in the lower depths.

5. Effect on biological properties of soil

5.1. Soil microbial populations: ZT conditions increased micro and macro soil organisms resulting ~~in~~ in larger number of worm channels ~~and to their continuity~~ that attributed to the higher infiltration rate of soil (**Hopp and Slater, 1961**). Earthworm channels increased ~~conveyed~~, which ~~increase~~ improved soil porosity, and provided for rapid water entry into a soil (**Doran *et al.*, 1980**). In comparison to that of conventional tilled plots, ~~observed~~ ZT plots had greater earthworm activity (up to five times), 35 and 57% ~~per cent~~ per cent higher aerobic counts and facultative anaerobic counts, ~~under ZT conditions~~. The population of denitrifying bacteria was almost half in tilled plots in comparison ted to ~~than~~ that of the ZT plots.

5.2. Soil respiration: CO₂ flux as impacted by agricultural management practices ~~need~~ needs to be delineated (**Reicosky *et al.*, 1997**). Tillage ~~opens the soil, thus~~ improves the soil respiration and ~~increased~~ increases the emission of the CO₂ (**Reicosky and Archer *et al.*, 2007**). Nowadays, a rapid increase of CO₂ in the environment is one of the main issues because

of reported global warming consequences (Wood *et al.*, 1990). However, soil management practices need to be refined to reduce soil respiration and organic matter decomposition without decreasing crop yield and ZT might be the suitable answer. But However, scientists are of different opinions as some reported similar soil CO₂ emission rates from ZT and CT (Elder and Lal, 2008), while others (Oortset *et al.*, 2007) observed large CO₂ emissions under zero tillage in comparison to the CT. Thus, a bridge between the two tillage systems might be an answer.

6. Effect on nodulation and root growth:

A long-term chickpea experiment showed that in comparison to a conventional tillage system, the quantity and dry weight of root nodules/plant under a zero tillage system were 33.3 and 21.1% greater, respectively (Lopez-Bellido *et al.*, 2011). According to a study conducted in Spain, during conventional tillage (0.34 mm/cm²), chickpea root length in the 0-15 cm layer was 38 and 27% larger than under zero tillage (0.18 mm/cm²) (Munoz-Romero *et al.*, 2012). The study carried out in Jabalpur (India) revealed that ZTs chickpea produced after transplanted rice had a considerably greater nodule dry weight (71.8 mg/plant) than both ZTs direct-seeded rice (55.2 g/plant) and direct-seeded rice (59.2 mg/plant) (Mishra *et al.*, 2012).

7. Effect on crop production economics and energy efficiency:

The data on the economics of the rabi chickpea crop emphasized that conventional tillage requires a higher cost of cultivation (16799.40 ha⁻¹) as compared to a minimum (16790.70 ha⁻¹) and zero tillage (16277.50 ha⁻¹) (Ahmad *et al.*, 2019). While they realized maximum gross return (Rs. 60599.78 ha⁻¹) and net returns (Rs. 43800.38 ha⁻¹) under conventional tillage. The higher returns under above conventional tillage might be due to higher seed yield coupled with lower cost of weed management treatments. Similar findings were also reported by (Singh *et al.*, 1991; Porwal, 2000 and Patel *et al.*, 2006).

Mishra *et al.* (2012) reported that the total cost of cultivation in TPR-CT chickpea system (₹ 31410/ha) was higher than in other systems due to higher costs involved in field preparation and transplanting operations in rice. The highest net returns (₹33,600/ ha) and benefit: cost ratio (2.33) was accrued with ZT-DSR, ZT- chickpea cropping system and the lowest with DSR-CT chickpea system (₹ 20,222/ha and 1.67, respectively). The PBR-CT chickpea system required maximum energy (₹ 20 867 MJ/ha) closely followed by TPR-CT chickpea (20,594 MJ/ha) due to higher energy required for puddling in rice and tillage operations in chickpea. The energy productivity (yield per unit energy consumed) was maximum (0.401 kg/MJ) in ZTDSR-ZT chickpea, followed by TPR-CT chickpea (0.327 kg/MJ). The output energy was maximum in TPR-CT chickpea (99,049 MJ/ha), followed by ZT DSR-ZT chickpea (96,344

Commented [KA15]: Pls explain.... Whether you mean Transplanted

Commented [KA16]: Pls chk units

Commented [KA17]: Pls chk units

MJ/ha) due to higher system productivity. The energy output:input ratio (5.90) was the highest in ZT-DSR-ZT—chickpea, followed by TPR-CT-chickpea (4.81). The lowest energy productivity (0.253 kg/MJ), output energy (72 765 MJ/ha) and output: input ratio (3.71) was observed from DSR-CT chickpea system.

According to **Roy and Singh (2012)**, on the sandy loam soils of Ludhiana, Punjab, the net monetary returns of chickpea were higher under ZT with Pantnagar seed drill (₹2650/ha) and ZT with paddy stubbles (₹2100/ha) than under reduced tillage-RT (₹1950/ha). [Similar A similar view was also reported by \(Kumari et al., 2019\) by conducting an experimentexperimenting](#) on chickpea at Bhagalpur.

8. Effect of zero tillage on weed dynamics in chickpea:

Weed is also another serious problem in rice based cropping system. Infestation of weeds suppresses the crop especially during initial growth period. Weeds can reduce the chickpea yield by the tune of 10-50% loss of crop yield depends upon the intensity of weed flora and management practices (**Yaduraju and Mishra., 2002**). A significant obstacle to achieving potential chickpea yield is weeds. The weed species infesting the chickpea fields vary from one location to another depending on the agro-climatic conditions, prevailing cropping systems, tillage practices and weed management strategies adopted.

The experimental field of chickpea was dominated with broad-leaved weeds (95.4%), viz. Clover (*Medicago hispida* Gaertn) (50 %), common vetch (*Vicia sativa* L.) (22.6 %), common lambsquarters (*Chenopodium album* L.) (7.3 %), and others including field bindweed (*Convolvulus arvensis* L.), *V. hirsuta* L., sweet clover (*Melilotus indica* All.) and *M. alba* L. (15.5 %). The grassy weeds, viz littleseed canary grass (*Phalaris minor* Retz.) (3.1%) and wild oats (*Avena sterilis*) (1.5%) were of minor importance. **(Mishra et al., 2011)** stated that Zero till chickpea significantly increased the population of *V. sativa* but reduced the problem of *C. album* as compared to conventional tillage. The population of *M. hispida*, *A. ludoviciana*, *P. minor*, and total weeds did not vary significantly due to change in tillage systems. Under zero tillage significantly reduced the dry matter of *M. hispida* and *C. album* ~~album was observed~~ by 20.3 and 58% as compared to conventional tillage, however, the dry matter of *A. ludoviciana* and *V. sativa* increased.

In a study at Kanpur, **(Kumar et al., 2022)** observed that, post-rainy seasons zero tillage with and without residues attributed higher weed diversity indices (Shannon and Simpson) compared with conventional tillage in seedbank. Importantly, **CT-ZT + R with R-C-Mb (interaction)** reduced 24% total viable seed density at 0–15 cm depth than CT-CT with R-W.

Commented [KA18]: Pls chk

Commented [KA19]: Pls explain

Zero tillage in post-rainy seasons after puddled transplanted rice and intensive ~~pulse based~~pulse-based cropping (rice-chickpea-mungbean) can minimize viable weed seeds in soil vis-a-vis above-ground weed density over time than conventional tillage and rice-wheat system. The reduced weed density could reduce soil fertility degradation, enhance crop ~~system~~ productivity, restore soil health and ~~provides~~provide opportunity for sustainable cropping intensification in rice ecologies of the Indo-Gangetic plains.

In a study conducted at IGFRI, Jhansi, zero tillage in the chickpea cropping sequence considerably reduced the weed density by 14.5-19.5% when compared to reduced and conventional tillage conditions (Dixit *et al.*, 2015).

Conclusion

Based on the research work carried out by various workers as reviewed, it may be concluded that:

1. Zero tillage significantly reduces weed density and biomass, mitigating weed competition and enhancing chickpea growth and development.
2. Weed dynamics under zero tillage exhibit a favorable trend, facilitating improved resource utilization by chickpea crops and reducing the need for intensive weed management.
3. Chickpea yields demonstrate marked improvement under zero tillage which is attributed to decreased weed interference and enhanced soil health.
4. The adoption of zero tillage offers a sustainable approach to weed management, promoting soil conservation and minimizing environmental impact.
5. Integrating zero tillage into rice-chickpea cropping systems holds promise for optimizing agricultural productivity while fostering long-term sustainability.

By reviewing the research work done by several workers it may be concluded that the efficacy and benefits of zero till chickpea in enhancing chickpea productivity and efficient weed management under rice-chickpea cropping systems may offer ~~,-thereby offering-~~ a pathway towards sustainable agricultural practices and improved farm profitability.

References -

Adarsh, S., Giffy, T. and Jacob, J. (2019). Role of pulses in cropping systems: A review. *Agricultural Reviews* 40(3): 185–191.

Commented [KA20]: Not as per journal style

AICRP-Sorghum. (2017). Sorghum agronomy. Indian Institute of Millets Research, Hyderabad.

Amgain, L. P., Sharma, A. R., Das, T. K., & Behera, U. K. (2013). Effect of residue management on productivity and economics of pearl millet (*Pennisetum glaucum*)-based cropping system under zero-till condition. *Indian Journal of Agronomy*, 58(3), 298–302.

Dixit, A. K., Kumar, S., Rai, A. K., & Kumar, T. K. (2015). System productivity, profitability, nutrient uptake and soil health under tillage, nutrient and weed management in rainfed chickpea (*Cicer arietinum*)–fodder sorghum (*Sorghum bicolor*) cropping system. *Indian Journal of Agronomy*, 60(2), 205–211.

Hemmat, A., & Eskandari, I. (2004). Tillage system effects upon productivity of a dryland winter wheat–chickpea rotation in the northwest region of Iran. *Soil and Tillage Research*, 78(1), 69–81.

Kumar, B. R., and Angadi, S. S. (2016). Effect of tillage, mulching and weed management practices on the performance and economics of chickpea. *Legume Research-An International Journal*, 39(5), 786–791.

Kumari, S., Kumar, S., Chaurasiya, A., Dutta, S. K., Pradhan, A., & Behera, S. K. (2019). Conservation agriculture based tillage options for rabi crops cultivation in eastern India: Effects on Productivity and Profitability. *International Journal of Current Microbiology and Applied Sciences*, 8(7), 1678–1689.

Mishra, J. S., Singh, V. P., Chandra, B., & Subrahmanyam, D. (2012). Crop establishment, tillage and weed management techniques on weed dynamics and productivity of rice (*Oryza sativa*)-chickpea (*Cicer arietinum*) cropping system. *Indian Journal of Agricultural Sciences*, 82(1), 15–20.

Piggin, C., Haddad, A., Khalil, Y., Loss, S., & Pala, M. (2015). Effects of tillage and time of sowing on bread wheat, chickpea, barley and lentil grown in rotation in rainfed systems in Syria. *Field Crops Research*, 173, 57–67.

Rathod, P. S., Patil, D. H., & Dodamani, B. M. (2017). Integrated weed management in chickpea (*Cicer arietinum* L.) under rainfed conditions of Karnataka, India. *Legume Research-An International Journal*, 40(3), 580–585.

Roy, R., & Singh, A. (2012). Crop establishment techniques for improving soil properties and productivity of chickpea (*Cicer arietinum* L.). *Journal of Food Legumes*, 25(3), 237–239.

Singh, D., Pazhanisamy, S., Kumar, S., Kumar, A., & Reddy, S. L. (2020). Bio-efficacy of different herbicides in broad spectrum weed management for chickpea. *International Journal of Current Microbiology and Applied Sciences*, 9(3), 2313–2317.

Singh, N. P., & Praharaj C. S. (2020). Scenario of pulses production in India, Abstracts of international conference on pulses as the climate smart crops: challenges and opportunities, Bhopal, India, 3–9.

Munoz-Romero, V., López-Bellido, L., & López-Bellido, R. J. (2012). The effects of the tillage system on chickpea root growth. *Field Crops Research*, 128, 76–81.

Ahmad, A., Chowdhury, T., & Kumar, A. (2019). Effect of Tillage and Weed Control Techniques on Energy Dynamics and Profitability of Chickpea (*Cicer arietinum* L.)-Rice Cropping Sequence in Irrigated Ecosystem of CG Plains. *Curr. J. App. Sci. and Tec*, 38(6), 1-8.

Ali M. (1992). Genotypic compatibility and spatial arrangement in chickpea (*Cicer arietinum* L.) and Indian mustard (*Brassica juncea*) in intercropping in north-eastern plains. *Indian Journal of Agricultural Sciences* 62(4) :249-53.

Arya, R.L., Kumar, L., Singh, K.K. and Kushwaha, B.L., (2005). Effect of fertilizers and tillage management in rice (*Oryzasativa*)–chickpea (*Cicer arietinum*) cropping system under varying irrigation schedules. *Indian Journal of Agronomy*, 50(4), pp.256-259.

López-Bellido, L., Benítez-Vega, J., García, P., Redondo, R. and López-Bellido, R.J., 2011. Tillage system effect on nitrogen rhizodeposited by faba bean and chickpea. *Field crops research*, 120(1), pp.189-195.

Banjara, T. R., Pali, G. P., Tigga, B. K., Kumar, S., &Shori, A. (2017). Effect of Different Tillage Practices on Growth, Yield and Economics of Chickpea (*Cicer arietinum* L.) under Rainfed Condition of Chhattisgarh. *International Journal of Current Microbiology and Applied Sciences*, 6(2), 1464-1470. <https://doi.org/10.20546/ijcmas.2017.602.164>.

Barley, K. P. (1961). The abundance of earthworms in agricultural land and their possible significance in agriculture. *Advances in agronomy*, 13, 249-268.

Bhatt, R and S.S. Kukal. (2015e). Soil physical environment as affected by double ZT in rice wheat cropping system of North-West India. *Asian J. Soil Sci.* 10 (1): 166-172.

Bhatt, R. (2015). Soil water dynamics and water productivity of rice-wheat system under different establishment methods. Dissertation submitted to the Punjab Agricultural University, Ludhiana.

Bhatt, R. (2017). Zero tillage impacts on soil environment and properties. *Journal of Environmental and Agricultural Sciences*, 10, 01-19.

Bhuler, D.D. (1995). Influence of tillage systems on weed population dynamics and management in corn and soybean production in central USA. *Crop science*.133 (35):1247-1257.

Blevins, R. L., Thomas, G. W., & Cornelius, P. L. (1977). Influence of No-tillage and Nitrogen Fertilization on Certain Soil Properties after 5 Years of Continuous Corn 1. *Agronomy journal*, 69(3), 383-386.

Blevins, R.L., G.W. Thomas and P.L. Cornelius (1977). Influence of no-tillage and nitrogen fertilization on certain soil properties after five years of continuous corn. *Agron. J.* 69:383-386.

- Boguzas, Vaclovas, Ausra Marcinkeviciene, and Rita Pupaliene (2010). "Weed response to soil tillage, catch crops and farmyard manure in sustainable and organic agriculture." 43-50.
- Bricchi E, Formia F, Esposito G, Riberi L, Aquino H (2004). The effect of topography, tillage and stubble grazing on soil structure and organic carbon levels. *Span. J. Agric. Res.* 2: 409-18.
- Bricchi, E., Formia, F., Espósito, G., Riberi, L., & Aquino, H. (2004). The effect of topography, tillage and stubble grazing on soil structure and organic carbon levels. *Spanish Journal of agricultural research*, 2(3), 409-418.
- Chatterjee, A and Lal, R. (2009). On farm assessment of tillage impact on soil carbon and associated soil quality parameters. *Soil Till. Res.* 104: 270-77.
- Chatterjee, A., & Lal, R. (2009). On farm assessment of tillage impact on soil carbon and associated soil quality parameters. *Soil and Tillage Research*, 104(2), 270-277.
- Daraghme, O. A., Jensen, J. R., & Petersen, C. T. (2009). Soil structure stability under conventional and reduced tillage in a sandy loam. *Geoderma*, 150(1-2), 64-71.
- Dick, W. A., Van Doren Jr, D. M., Triplett Jr, G. B., & Henry, J. E. (1986). Influence of long-term tillage and rotation combinations on crop yields and selected soil parameters
- Dick, W.A., D.M. Van Doren, G.B. Triplett and J.E Henry (1986). Influence of long term tillage and rotation combinations on crop yields and selected soil parameters. II. Results obtained for a TypicFragiudalf soil. *OhilAgric Res Dev Cent* 1181: 1-34.
- Dinnes, D. L., Karlen, D. L., Jaynes, D. B., Kaspar, T. C., Hatfield, J. L., Colvin, T. S., &Cambardella, C. A. (2002). Nitrogen management strategies to reduce nitrate leaching in tile drained Midwestern soils. *Agronomy journal*, 94(1), 153-171.
- Doran, J. W. (1980). Soil microbial and biochemical changes associated with reduced tillage. *Soil Science Society of America Journal*, 44(4), 765-771.
- Edwards, J. H., Thurlow, D. L., & Eason, J. T. (1988). Influence of tillage and crop rotation on yields of corn, soybean, and wheat. *Agronomy Journal*, 80(1), 76-80.
- Edwards, J.H., D.L. Thurlow and J.T. Eason. (1988). Influence of tillage and crop rotation on yield of corn, soybean, and wheat. *Agron. J.* 80: 76-80.
- Elder, J. W., & Lal, R. (2008). Tillage effects on gaseous emissions from an intensively farmed organic soil in North Central Ohio. *Soil and Tillage Research*, 98(1), 45-55.
- FAO. (2012). Advancement and impact of conservation agriculture/no-till technology adoption in Kazakhstan. FAO Information Note. Retrieved from <http://www.fao.org/ag/ca/doc/Importance Zero Tillage Northern Kazakhstan.pdf>
- Francis, G. S., & Knight, T. L. (1993). Long-term effects of conventional and no-tillage on selected soil properties and crop yields in Canterbury, New Zealand. *Soil and Tillage Research*, 26(3), 193-210.

- Franzluebbers, A.J and M.A. Arshad. (1996b). Soil organic matter pools with conventional and ZT in a cold, semiarid climate. *Soil till. Res.* 39: 1–11.
- Freitas, P.L., P. Blancaneaux, E. Gavinelli, M.C. Larré-larrouy and C. Feller. (1999). Nature and level of organic stock in clayey Oxisols under different land use and management systems. *Pesq Agro pec Bras.* 35:157-170.
- Frye, W. W., Smith, W. G., & Williams, R. J. (1985). Economics of winter cover crops as a source of nitrogen for no-till corn. *Journal of Soil and Water Conservation*, 40(2), 246-249.
- Frye, W.W., O.L. Burnett and G.J. Buntley. (1985). Restoration of crop productivity on eroded or degraded soils. In *Soil Erosion and Crop Productivity*, eds. R.F. Follet and B.A. Stewart. Madison, WI: *American Society of Agronomy Journal*. Pp: 335- 356.
- Ghosh, P.K., A. Das, R. Saha, E. Kharkrang, A.K. Tripathi, G.C. Munda and S.C. Ngachan. (2010). Conservation agriculture towards achieving food security in North East India. *Curr. Sci.* 99(7): 915-921.
- Gangwar, K.S. and Singh, H.R., (2010). Effect of rice (*Oryza sativa*) crop establishment technique on succeeding crops. *Indian Journal of Agricultural Sciences*, 80(1), pp.24-28.
- Ghuman, B.S and H.S. Sur. (2001). Tillage and residue management effects on soil properties and yields Eastern Gangetic Plains of South Asia: Yield trends and economic profitability. *Field Crops Res.* 164: 199-210.
- Green, V. S., Cavigelli, M. A., Dao, T. H., & Flanagan, D. C. (2005). Soil physical properties and aggregate-associated C, N, and P distributions in organic and conventional cropping systems. *Soil Science*, 170(10), 822-831.
- Hazarika S, Parkinson R, Bol R, Dixon L, Russel P, Donovan S, Allen D (2009). Effect of tillage system and straw management on organic matter dynamics. *Agron. Sust. Dev.* 29: 525-33.
- Hobbs, P. R., Sayre, K., & Gupta, R. (2008). The role of conservation agriculture in sustainable agriculture. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 363(1491), 543-555.
- Jat, M. L., R. K. Gupta, O. Erenstein, and R. Ortiz. (2006). Diversifying the intensive cereal cropping systems of the Indo - Ganges through horticulture. *ChronicaHorticulturae* 46 (3), 27–31.
- Karlen D.L., Wollenhaupt N.C., Erbach D.C., Berry E.C., Swan J.B., Eash N.S., Jordahl J.L. (1994). Long-term effects on soil quality. *Soil and Tillage Research*, 32: 313–327.
- Kumar Narendra, Nath C.P., Hazra K.K., Praharaj, C.S. Singh S.S., and Singh N.P. (2022). Long-term impact of zero-till residue management in post-rainy seasons after puddled rice and cropping intensification on weed seedbank, above-ground weed flora and crop productivity. *Ecological Engineering* 176:106540.

- Lal, R. *et al.* (1989). Conservation tillage for sustainable agriculture: tropics vs. temperate environments. *Advances in Agronomy*, 42, 85–197.
- Mannering, J.V., R. Griffith and B. Richey. (1975). Tillage for moisture conservation. Ameri. Soc. of Agriculture and Engineering. *St Joseph MI, Paper No. 75-2523*.
- Martino, D. L., & Shaykewich, C. F. (1994). Root penetration profiles of wheat and barley as affected by soil penetration resistance in field conditions. *Canadian Journal of Soil Science*, 74(2), 193-200.
- McConkey, B.G., B.C. Liang, A. Campbell, D. Curtin, A. Moulin, S.A. Brandt and G.P. Lafond. (2003). Crop rotation and tillage impact on carbon sequestration in Canadian prairie soils. *Soil Till. Res.* 74:81–90.
- Miller, P., McKay, K., Jenks, B., Riesselman, J., Neill, K., Buschena, D. and Bussan, A. J. (2002). Growing chickpea in the northern Great Plains. MontGuide fact sheet #200204/Agriculture, Montana St Univ Ext Serv, Bozeman, MT, USA [Online] Available: <http://www.montana.edu/wwwwpb/pubs/mt200204.html> [2005 Dec. 15]
- Mishra, J.S. and Singh, V.P., (2011). Effect of tillage and weed control on weed dynamics, crop productivity and energy-use efficiency in rice (*Oryzasativa*)-based cropping systems in Vertisols. *Indian Journal of Agricultural Sciences*, 81(2), p.129.
- Mishra, J.S., Singh, V.P., Bhanu, C. and Subrahmanyam, D., (2012). Crop establishment, tillage and weed management techniques on weed dynamics and productivity of rice (*Oryzasativa*)—Chickpea (*Cicer arietinum*) cropping system. *Indian Journal of Agricultural Sciences*, 82(1), p.15.
- Moussa-Machraoui SB, Errouissi F, Ben-Hammouda M, ouira S (2010). Comparative effects of conventional and no-tillage management on some soil properties under Mediterranean semi-arid conditions in north-western Tunisia. *Soil Till Res* 106: 247-53.
- Oorts, K., Merckx, R., Gréhan, E., Labreuche, J., & Nicolardot, B. (2007). Determinants of annual fluxes of CO₂ and N₂O in long-term no-tillage and conventional tillage systems in northern France. *Soil and Tillage Research*, 95(1-2), 133-148.
- Patni, N. K., Masse, L., & Jui, P. Y. (1998). *Groundwater quality under conventional and no tillage: I. Nitrate, electrical conductivity, and pH* (Vol. 27, No. 4, pp. 869-877). *American Society of Agronomy, Crop Science Society of America, and Soil Science Society of America*.
- Poonia, T. C., and M. S. Pithia (2013). "Pre-and post-emergence herbicides for weed management in chickpea." 223-225.
- Quddus, M. A., Naser, H. M., Siddiky, M. A., Ali, M. R., Mondol, A. T. M. A. I., Islam, M. A., & Gazipur, B. (2020). Impact of zero tillage and tillage practice in chickpea production. *Journal of Agricultural Science*, 12(4), 106-118.

Rahman MH, Okubo A, Sugiyama S, Mayland HF (2008). Physical, chemical and microbiological properties of an Andisol as related to land use and tillage practice. *Soil Till.*

Reicosky, D. C., & Archer, D. W. (2007). Moldboardplow tillage depth and short-term carbon dioxide release. *Soil and Tillage Research*, 94(1), 109-121.

Reicosky, D. C., Dugas, W. A., & Torbert, H. A. (1997). Tillage-induced soil carbon dioxide loss from different cropping systems. *Soil and Tillage Research*, 41(1-2), 105-118. Res. 101:10-19.

Six, J., Conant, R. T., Paul, E. A., & Paustian, K. (2002). Stabilization mechanisms of soil organic matter: implications for C-saturation of soils. *Plant and soil*, 241, 155-176.

Smart, J.R and Bradford, J.M(1999) Conservation tillage corn production for a semi-arid, subtropical environment. *Agronomy Journal*. 91:116-21

TarkalsonDD, Gary WH, Kassman KG (2006). Long-term effects of tillage on soil chemical properties and grain yields of a dryland winter wheat-sorghum/ corn-fallow rotation in the great plains. *Agron. J.* 98: 26-33.

Wood, D. M. (1990). *Soil behaviour and critical state soil mechanics*. Cambridge university press.

Yaduraju N T and Mishra J S. (2002). Weeds management in chickpea: Challenges and Opportunities. *Agricultural Situation in India*, pp 423–30. October (2002).

Singh, A., Phogat, V.K., Dahiya, R. and Batra, S.D., (2014). Impact of long-term zero till wheat on soil physical properties and wheat productivity under rice–wheat cropping system. *Soil and Tillage Research*, 140, pp.98-105.

Monneveux, P., Quill rou, E., S nchez, C. and Lopez-Cesati, J., (2006). Effect of zero tillage and residues conservation on continuous maize cropping in a subtropical environment (Mexico). *Plant and Soil*, 279, pp.95-105.

Nazeer, S., Malik, A.U., Nazir, G. and Ahmad, J., 2012. Effectiveness of tillage systems and farm manure levels on rice productivity.