
On Sums of Squares Involving Integer Sequence:

$$\sum^n w^2 + \frac{n}{3}d^2 = 3\left(\frac{nd^2}{3} + \sum^{\frac{n}{3}} w^2\right)$$

**Authors
Contribution**

This work was carried out by the all authors

**Original Research
Article**

Abstract

Let w_r be a given integer sequence in arithmetic progression with a common difference d . The study of diophantine equations, which are polynomial equations seeking integer solutions, has been a very interesting journey in the field of number theory. Historically, these equations have attracted the attention of many mathematicians due to their intrinsic challenges and their significance in understanding the properties of integers. In this current study, we examine a diophantine equation relating the sum of squared integers from specific sequences to a variable d . In particular, the diophantine equation $\sum_{r=1}^n w_r^2 + \frac{n}{3}d^2 = 3\left(\frac{nd^2}{3} + \sum_{r=1}^{\frac{n}{3}} w_{3r-1}^2\right)$ is introduced and partially characterized. The objective is to determine the conditions under which integer solutions for (w_r, d) exist within this diophantine equation. The methodology of solving this problem entails, decomposing polynomials, factorizing polynomials, and exploring the solution set of the given equation.

*hlao@kyu.ac.ke

Keywords: Diophantine Equation; Sums of Squares, Integer Sequence
 2010 Mathematics Subject Classification: 53C25; 83C05; 57N16

1 Introduction

Diophantine equations, tracing their roots back to the era of the ancient Greek mathematician Diophantus, continue to be a captivating challenge within number theory. These equations, seeking integer solutions, hold significant importance due to their real-life applications. Despite the extensive exploration of various Diophantine equations, including renowned challenges like Fermat's Last Theorem, Ramanujan Nagell equation, and Lebesgue Nagell equation, as well as studies focusing on polynomials of degree less than 5, the specific examination of the diophantine equation $\sum_{r=1}^n w_r^2 + \frac{n}{3}d^2 = 3(\frac{nd^2}{3} + \sum_{r=1}^{\frac{n}{3}} w_{3r-1}^2)$ remains largely uncharted. Recent research has delved into the intricacies of polynomials with degrees less than 5, as referenced in [1, 2,3,10,11,12]. For a comprehensive understanding of studies related to Fermat's Last Theorem and Ramanujan Nagell equations, readers are encouraged to explore [4,5,6,7,8,9,13,14,15,16]. Within the existing body of work, the literature concerning the diophantine equation $\sum_{r=1}^n w_r^2 + \frac{n}{3}d^2 = 3(\frac{nd^2}{3} + \sum_{r=1}^{\frac{n}{3}} w_{3r-1}^2)$ remains largely unexplored. This study aims to contribute to this knowledge gap by introducing and developing the formula $\sum_{r=1}^n w_r^2 + \frac{n}{3}d^2 = 3(\frac{nd^2}{3} + \sum_{r=1}^{\frac{n}{3}} w_{3r-1}^2)$, seeking to enhance our comprehension of this specific diophantine equation within the broader landscape of mathematical exploration.

2 Main Results

In the following sections, we begin by articulating our observations as a conjecture, and subsequently, we proceed to obtain solutions for particular instances of the aforementioned diophantine equation.

Conjecture 2.1. *For any integer n divisible by 3, the diophantine equation*

$$\sum_{r=1}^n w_r^2 + \frac{n}{3}d^2 = 3(\frac{nd^2}{3} + \sum_{r=1}^{\frac{n}{3}} w_{3r-1}^2) \cdots (1)$$

admits solutions in integers if $w_n - w_{n-1} = w_{n-1} - w_{n-2} = \cdots = w_2 - w_1 = d$

In the subsequent sections, the focus of this investigation revolves around identifying the values of the variables $(n, w_1, w_2, \dots, w_n, d)$ that fulfill the conditions of equation (1). Consequently, distinct cases have been established.

Theorem 2.2. *Consider equation (1) satisfying the condition $(n, w_1, w_2, w_3, d) = (3, w_1, w_2, w_3, d)$ Then, the diophantine equation*

$$w_1^2 + w_2^2 + w_3^2 + d^2 = 3(d^2 + w_2^2)$$

has solution in integers if $w_3 - w_2 = w_2 - w_1 = d$.

Proof. Assume that $w_2 = w_1 + d, w_3 = w_1 + 2d, w_3 = w_1 + 2d$ and Consider the equation $w_1^2 + w_2^2 + w_3^2 + d^2 = 3(d^2 + w_2^2) \cdots (2.1)$. The, left hand side of equation (2.1) expressed as

$$w_1^2 + w_2^2 + w_3^2 + d^2 = w_1^2 + (w_1 + d)^2 + (w_1 + 2d)^2 + d^2$$

simplifies to

$$3w_1^2 + 6w_1d + 6d^2 = 3(w_1^2 + 2w_1d + 2d^2) \cdots (2.1.1).$$

Decomposing equation (2.1.1) into thrice sums of two squares, we obtain,

$$\begin{aligned} 3w_1^2 + 6w_1d + 6d^2 &= 3(w_1^2 + 2w_1d + 2d^2) \\ &= 3((w_1^2 + 2w_1d + d^2) + d^2) = 3(d^2 + (w_1 + d)^2) = 3(d^2 + w_2^2). \end{aligned}$$

This complete the proof. □

Theorem 2.3. Consider equation (1) satisfying the condition $(n, w_1, w_2, \dots, w_6, 2d) = (6, w_1, w_2, \dots, w_6, 2d)$. Then, the diophantine equation

$$w_1^2 + w_2^2 + w_3^2 + w_4^2 + w_5^2 + w_6^2 + 2d^2 = 3(2d^2 + w_2^2 + w_5^2)$$

has solution in integers if $w_6 - w_5 = w_5 - w_4 = w_4 - w_3 = w_3 - w_2 = w_2 - w_1 = d$.

Proof. Let $w_2 = w_1 + d, w_3 = w_1 + 2d, w_4 = w_1 + 3d, w_5 = w_1 + 4d, w_6 = w_1 + 5d$ and Consider the equation $w_1^2 + w_2^2 + w_3^2 + w_4^2 + w_5^2 + w_6^2 + 2d^2 = 3(2d^2 + w_2^2 + w_5^2) \dots (2.2)$. The, left hand side written as

$$w_1^2 + (w_1 + d)^2 + (w_1 + 2d)^2 + (w_1 + 3d)^2 + (w_1 + 4d)^2 + (w_1 + 5d)^2 + (w_1 + 6d)^2 + 2d^2$$

reduces to

$$6w_1^2 + 30w_1d + 57d^2 = 3(w_1^2 + 10w_1d + 19d^2) = 3(2d^2 + w_1^2 + 10w_1d + 17d^2) \dots (2.2.1).$$

Breaking equation (2.2.1) into thrice sums of sums of four squares, we get,

$$\begin{aligned} 3(2d^2 + w_1^2 + 10w_1d + 17d^2) &= 3(2d^2 + (w_1^2 + 2w_1d + d^2) + (w_1^2 + 8w_1d + 16d^2)) \\ &= 3(2d^2 + (w_1 + d)^2 + (w_1 + 4d)^2) = 3(2d^2 + w_2^2 + w_5^2) \end{aligned}$$

This concludes the proof. □

Theorem 2.4. Consider equation (1) satisfying the condition

$$(n, w_1, w_2, w_3, w_4, w_5, w_6, w_7, w_8, w_9, 3d) = (9, w_1, w_2, w_3, w_4, w_5, w_6, w_7, w_8, w_9, 3d).$$

Then, the diophantine equation

$$w_1^2 + w_2^2 + w_3^2 + w_4^2 + w_5^2 + w_6^2 + w_7^2 + w_8^2 + w_9^2 + 3d^2 = 3(3d^2 + w_2^2 + w_5^2 + w_8^2)$$

has solution in integers if $w_9 - w_8 = w_8 - w_7 = w_7 - w_6 = w_6 - w_5 = w_5 - w_4 = w_4 - w_3 = w_3 - w_2 = w_2 - w_1 = d$.

Proof. Consider the equation $w_1^2 + w_2^2 + w_3^2 + w_4^2 + w_5^2 + w_6^2 + w_7^2 + w_8^2 + w_9^2 + 3d^2 = 3(3d^2 + w_2^2 + w_5^2 + w_8^2)$. Assume that $w_2 = w_1 + d, w_3 = w_1 + 2d, w_4 = w_1 + 3d, w_5 = w_1 + 4d, w_6 = w_1 + 5d, w_7 = w_1 + 6d, w_8 = w_1 + 7d, w_9 = w_1 + 8d$. The, left hand side expressed as,

$$w_1^2 + (w_1 + d)^2 + (w_1 + 2d)^2 + (w_1 + 3d)^2 + (w_1 + 4d)^2 + (w_1 + 5d)^2 + (w_1 + 6d)^2 + (w_1 + 7d)^2 + (w_1 + 8d)^2 + 3d^2$$

simplifies to

$$9w_1^2 + 72w_1d + 207d^2 = 3(3w_1^2 + 24w_1d + 69d^2) \dots (2.3).$$

Decomposing equation (2.3) into triple sums of squares, we obtain,

$$\begin{aligned} &= 3(3d^2 + (w_1^2 + 2w_1d + d^2) + (w_1^2 + 8w_1d + 16d^2) + (w_1^2 + 14w_1d + 49d^2)) \\ &= 3(3d^2 + (w_1 + d)^2 + (w_1 + 2d)^2) + (w_1 + 4d)^2 + (w_1 + 7d)^2 = 3(3d^2 + w_2^2 + w_5^2 + w_8^2) \end{aligned}$$

This complete the proof. □

Theorem 2.5. Consider equation (1) satisfying the condition $(n, w_1, w_2, \dots, w_{12}, 4d) = (12, w_1, w_2, \dots, w_{12}, 4d)$. Then, the diophantine equation

$$w_1^2 + w_2^2 + w_3^2 + w_4^2 + w_5^2 + w_6^2 + w_7^2 + w_8^2 + w_9^2 + w_{10}^2 + w_{11}^2 + w_{12}^2 + 4d^2 = 3(4d^2 + w_2^2 + w_5^2 + w_8^2 + w_{11}^2)$$

has solution in integers if $w_{12} - w_{11} = w_{11} - w_{10} = w_{10} - w_9 = w_9 - w_8 = w_8 - w_7 = w_7 - w_6 = w_6 - w_5 = w_5 - w_4 = w_4 - w_3 = w_3 - w_2 = w_2 - w_1 = d$.

Proof. Consider the equation,

$$w_1^2 + w_2^2 + w_3^2 + w_4^2 + w_5^2 + w_6^2 + w_7^2 + w_8^2 + w_9^2 + w_{10}^2 + w_{11}^2 + w_{12}^2 + 4d^2 = 3(4d^2 + w_2^2 + w_5^2 + w_8^2 + w_{11}^2)$$

and Suppose that $w_2 = w_1 + d, w_3 = w_1 + 2d, w_4 = w_1 + 3d, w_5 = w_1 + 4d, w_6 = w_1 + 5d, w_7 = w_1 + 6d, w_8 = w_1 + 7d, w_9 = w_1 + 8d, w_{10} = w_1 + 9d, w_{11} = w_1 + 10d, w_{12} = w_1 + 11d$. The left hand side expressed as

$$w_1^2 + (w_1 + d)^2 + (w_1 + 2d)^2 + (w_1 + 3d)^2 + (w_1 + 4d)^2 + (w_1 + 5d)^2 + (w_1 + 6d)^2 + (w_1 + 7d)^2 + (w_1 + 8d)^2 + (w_1 + 10d)^2 + (w_1 + 11d)^2 + 4d^2.$$

simplifies to

$$12w_1^2 + 132w_1d + 510d^2 = 3(4w_1^2 + 44w_1d + 170d^2) \dots (2.4).$$

Splitting equation (2.4) into thrice sums of squares, we obtain,

$$\begin{aligned} &3(4d^2 + (w_1^2 + 2w_1d + d^2) + (w_1^2 + 8w_1d + 16d^2) + (w_1^2 + 14w_1d + 49d^2) + (w_1^2 + 18w_1d + 81d^2) + \\ &(w_1^2 + 20w_1d + 100d^2) + (w_1^2 + 22w_1d + 121d^2)) \\ &= 3(4d^2 + (w_1 + d)^2 + (w_1 + 4d)^2 + (w_1 + 7d)^2 + (w_1 + 10d)^2) \\ &= 3(4d^2 + w_2^2 + w_5^2 + w_8^2 + w_{11}^2). \end{aligned}$$

This complete the proof. □

3 Conclusion

In summary, the solution of the diophantine equation $\sum_{r=1}^n w_r^2 + \frac{n}{3}d^2 = 3(\frac{nd^2}{3} + \sum_{r=1}^{\frac{n}{3}} w_{3r-1}^2)$ under the specified conditions of a common difference d between consecutive terms $w_n, w_{n-1}, \dots, w_2, w_1$ where $w_n - w_{n-1} = w_{n-1} - w_{n-2} = \dots = w_2 - w_1 = d$ has been achieved for some cases. This solution provides valuable insights into the relation among the sequence terms, enhancing our understanding of the inherent patterns and structures within the equation. For future investigations, it is recommended to explore extensions of this diophantine equation by proving conjecture (1).

Acknowledgements

The author would like to thank the anonymous reviewers for carefully reading the article and for their helpful comments.

References

- [1] Amir .F., Pooya .M., Rahim. F., (2012). A Simple Method to Solve Quartic Equations. *Australian Journal of Basic and Applied Sciences*, 6(6): 331-336, ISSN ,1991-8178.
- [2] Bombieri. E., Bourgain. J., (2015) A problem on sums of two squares, *Internatinal Mathematics Research* , (11):3343-3407.

-
- [3] Cavallo, A. (2019). Galois groups of symmetric sextic trinomials. arXiv:1902.00965v1 [math.GR]. <https://arxiv.org/abs/1902.00965>.
- [4] David A., (2016) A partition-theoretic proof of Fermat's two squares theorem. *Discrete Mathematics* ,339:4:1410–1411, DOI:10.1016/j.disc.2015.12.002.
- [5] Giorgos P. Kouropoulos., (2021). A combined methodology for approximate estimation of the roots of the general sextic polynomial equation. *Research Square* , DOI: <https://doi.org/10.21203/rs.3.rs-882192/v2>.
- [6] Lao H., (2022).Some Formulae For Integer Sums of Two Squares.*Journal of Advances in Mathematics and Computer Science* , 37(4): 53-57, Article no.JAMCS.87824,ISSN: 2456-9968.DOI: 10.9734/JAMCS/2022/v37i430448.
- [7] Lao H., (2024).Radical Solution of Some Higher Degree Equation Via Radicals.*Journal of Advances in Mathematics and Computer Science* ,volume 39(3):20-28,Article no.JAMCS.113540|ISSN: 2456-9968, DOI: 10.9734/JAMCS/2024/v39i31872
- [8] Lao H.,Zachary K., Kinyanjui J. (2023).Some Generalized Formula For Sums of Cube.*Journal of Advances in Mathematics and Computer Science* , 37(4): 53-57, Article no.JAMCS.87824,ISSN: 2456-9968, DOI: 10.9734/JAMCS/2023/v38i81789.
- [9] Lao. H., Maurice. O, Michael . O., (2023).On The Sum of Three Square Formula,*Science Mundi* ,Vol 3(Iss.1), pp.111-120, Science Mundi ISSN:2788-5844 <http://sciencemundi.net>. DOI: <https://doi.org/10.51867/scimundi.3.1.11>
- [10] Kimtai B., Lao, H., (2023). *On Generalized Sum of Six, Seven and Nine Cube*. Mundi,Vol 3(1),pp.135-142, Science Mundi ISSN:2788-5844 <http://sciencemundi.net> DOI: <https://doi.org/10.51867/scimundi.3.1.14>
- [11] Mochimaru, Y. (2005). Solution of Sextic Equations. *International Journal of Pure and Applied Mathematics*, 23(4), 575-583. <https://ijpam.eu/contents/2005-23-4/9/9>.
- [12] Najman .F., (2010). On The Diophantine Equation $x^4 \pm y^4 = iz^2$ in Gaussian Integers. *Amer. Math. Monthly*, 117(7), 637-641.
- [13] Najman .F., (2011).Torsion of elliptic curves over quadratic cyclotomic fields *Math. J. Okayama Univ.* , 53, 75-82.
- [14] Par Y. , (2016).Waring-Golbach problem. Two squares and Higher Powers. *Journal de Theorie des Nombres* ,791-810.
- [15] Ruffini, P. (1799). Teoria Generale delle Equazioni, in cui si dimostra impossibile la soluzione algebrica delle equazioni generali di grado superiore al quarto [General Theory of equations, in which the algebraic solution of general equations of degree higher than four is proven impossible]. Book on Demand Ltd. ISBN: 978-5519056762
- [16] Tignol, J. P. (2001). Galois' Theory of Algebraic Equations. World Scientific, Louvain, Belgium. DOI: 10.1142/4628 .