

Formulation and Evaluation of Calendula and Lavender Oil Based Topical Antifungal Gel

ABSTRACT

Aim: *Calendula officinalis* and *Lavendula angustifolia* are traditional medicinal plants that have antifungal activity. A combination of these two plants has not been known for its activity against fungus. The present study deals with the formulation and evaluation of herbal topical Antifungal gel containing essential oils of both plants.

Study design: Proto type research design

Place and Duration of Study: Smt. R. D. Gardi Pharmacy College, Nyara- Rajkot. January 2024 to April 2024.

Methodology: As 90% of skin infections are caused by *Candida albicans*, the antifungal activity of topical gel against this species was assessed. In this investigation, we created an antifungal herbal gel by combining carbopol 940 and other excipients with the essential oils of *Calendula officinalis* and *Lavendula angustifolia* in varying concentrations. For compatibility study, FTIR study would be done. Evaluation of Antifungal gel performed by using different tests likes, pH, Stability, Extrudability, Spreadability, Viscosity, Antifungal activity against *Candida albicans*, and *in-vitro* drug diffusion study.

Results: As a result, the formulation's physicochemical characteristics, *in vitro* antifungal efficacy, and stability analysis (stable even after 30 days) were assessed. Every herbal gel formulation with a pH of 5–7 had favorable outcomes for physicochemical measures. Out of all the formulations, batch no 4 exhibited superior release characteristics (98 %) and zone of inhibition in comparison to the other formulated batches. The drug release profile of batch 4 showed great results.

Conclusion: This is the first study on the scientific evaluation of *Calendula officinalis* and *Lavendula angustifolia* essential oil as a gel for antifungal activity. Thus this study reveals both good antifungals and *in-vitro* drug release; their essential oil may be formulated as Topical Gel Antifungal with satisfactory physicochemical parameters.

Keywords: *Calendula officinalis*, *Lavendula angustifolia*, *Candida albicans*, Essential Oil, Topical Antifungal Gel.

1. INTRODUCTION OF ANTIFUNGAL TOPICAL GEL

"A gel is a semi-solid, three-dimensional matrix created by the permeability of a solvent into a network of intertwined polymer chains or by an interspersed system of colloidal particles".[1,2,3] "A gelator, also known as a gelling agent, is added to the solvent and active component mixture to create pharmaceutical gels".[4,5,6] "Small compounds with low molecular weight or polymers can be utilized as gelators in gel formulation".[7,8] "As a dispersion media, the solvent might be an aqueous, organic, inorganic, or a combination of solvents".[9,10,11,79,80,81] "Topical gels are applied to the skin to act as a contact or transport medium for active medications".[12,13,14,15] The gel's three-dimensional mesh entangles the active drug molecules. Gels differ from conventional dosage forms in a few

key ways, including stiffness[16], rheology, swelling, aging, and syneresis[17]. Gels can be categorized based on a range of factors, including their physical makeup, kind of colloidal phase, and type of solvent utilized.[18,19] “Due to its advantages over creams and ointments, topical gels are a topical medication delivery dosage form that is frequently employed in cosmetics and treatments for skin ailments. They are created by combining a gelator, solvent, active medication, and other excipients”. [20,21]

1.1 Introduction of calendula essential oil



Figure 1: *Calendula officinalis*

“*Calendula officinalis* is a flowering plant in the daisy family asteraceae”. [22,23] “Its florets are edible”. [24] “They are often used to add color to salads or added to dishes as a garnish instead of saffron”. [25] “The essential oil of calendula is high in triterpenoid content which heals dry skin, eczema, and hemorrhoids. *Calendula officinalis* is widely cultivated and can be grown easily in sunny locations in most kinds of soils”. [26]

1.1.1 Phytoconstituents

“The plant has flavonoids, triol triterpenes, xanthophylls, esquiterpene glycosides, saponins, and volatiles, terpenoids, flavonoids, triterpeneol esters, steroids, phenolic compounds, carotenes, triterpenoids, essential oils, quinones, fatty acids, minerals, saponins, carbohydrates, sterols, and tocopherols, among other classes of chemical compounds. *Calendula officinalis* flowers are orange-colored because it is rich in carotenoids. The oil is rich in volatile oils, sterols, beta carotene, rutin, lupeol, narcissin, and calendulin. *Calendula officinalis* contains flavonoids, including quercetin, which has strong wound-healing properties”. [27, 28]

1.1.2 Medicinal benefits of calendula oil

Calendula oil works wonders as a moisturizer for all skin types. The oil relieves the damaged region and does wonders for dry or split skin. Calendula essential oil helps to cure hemorrhoids, eczema, and dry skin. [28,29] It cures mouth infections and lowers body temperature. One can consume calendula oil internally. [30,31,32] On the outside, the oil is utilized in soaps, lip balms, lotions, and creams. [33,34] According to studies, the oil known as “the mother of the skin” is highly regarded by homeopathic herbalists. [35,36,37] It exhibits pharmacological activity like anti-inflammatory [38], antioxidant activity [39], cytotoxic and anti-tumor activity [40], wound-healing activity [41], antimicrobial activity [42], antibacterial activity [43], and antifungal activity [44].

1.1.3 Pharmacological activity

Calendula oil shows pharmacological activities like antiviral[30,31], antifungal[32,33], antioxidant[34,35], antiseptic[36], antitumor[37,38], anti-inflammatory[39,40], antioxidant[41,42], hepatoprotective[43], and antidiabetic activity.[44]

1.2 Introduction of Lavender essential oil



Figure 2: *Lavendula angustifolia*

The plant's scent is produced by shimmering oil glands embedded in the microscopic plant hairs, or trichomes, that cover the stems, leaves, and flowers of the plant. Taxonomic classification of *lavendula angustifolia* is: scientific name: lavandula, family: lamiaceae, subfamily: nepetoideae, kingdom: plantae, order: lamiales, tribe: ocimeae.

1.2.1 Medicinal benefits of lavender oil

Lavender may have antifungal effects, as shown by numerous research.[45] Studies suggest that the essential oil of lavender may help inhibit the growth of certain fungi, such as candida albicans. The oil can also be used to treat athlete's foot and ringworm.[46] The pharmacological actions of lavender angustifolia oil are well-known, and they include sedative, antidepressant, healing, antiseptic, antifungal, relaxing, and antiemetic qualities.[47]

1.2.2 Phytoconstituents

The main component of lavender essential oils is monoterpenes. The most common monoterpenes are linalool and linalyl acetate.[48,49,50] Volatile oils like linalool, limonene, perillyl alcohol[51,52], linalyl acetate[53,54], cis-smine[55,56], terpene[57], coumarin, tannin[59], caffeic acid[58,60], and camphor[61]. The main constituents that give antifungal activities are linalyl acetate and linalool.

1.2.3 Pharmacological activity

The plant shows pharmacological activities like antimicrobial[62], anti-inflammatory, antinociceptive properties, anxiolytic, antioxidant[63], and antifungal activity[64].

2. MATERIAL AND METHODS

2.1 API and Excipients

API (Active pharmaceutical ingredients): Calendula oil and Lavandula oil.

Excipients: Carbopol 940, Glycerine, Propylene glycol, DM DM Hydantoin, PEG-40 Hydrogenated Castor oil, Diethylene glycol monoethyl ether, Vitamin E, Triethanolamine [65-74]

2.2 Collection of Essential Oil

2.2.1 Collection of Calendula Oil:

Veda oil Calendula Essential Oil, Marigold Essential Oil | ACTIZEET, Calendula Oil Pure & Natural Carrier Oil, Deve Herbs Pure Calendula Oil, Ks Essentials *Calendula Officinalis* From above mentioned different branded calendula oil products we have selected the Ks Essentials *Calendula Officinalis* as it is good quality and low-cost product compared to other products. Ks Essentials 100% Pure Calendula Essential Oil Pure Natural & Undiluted for Skin Care & Hair. [72]

Based on certificate of analysis, it is proven that the selected essential oil product complies with the standard.

Table 1: Certificate of Analysis of Calendula Essential Oil

Certificate of Analysis		
Product: Calendula Oil		
Shelf Life: 2 Years		
Test	Standards	Results
Botanical Name	<i>Calendula Officinalis</i>	Complies
Appearance & Color	Clear yellow to orange liquid	Complies
Odor	Mild characteristic odor	Complies
Refractive index at 20°C	1.450 - 1.510	1.467
Acid Value	Max 4.0	1.04
Specific gravity at 25°C	0.905 - 0.923	0.920
Iodine Value	112 – 145	123.20
Peroxide Value	Max 20.0	3.36
Saponification Value	190-200	197
Unsaponifiable Matter	Max 1.50%	1.0%
Water	Max 0.1%	Complies
Solubility	Very slightly soluble in Alcohol and completely miscible With light petroleum	
Storage	In a well-fitted container, in a cool and dark place.	

2.2.2 Collection of Lavender Essential oil

Vagad's khadi Herbal Lavender Essential Oil is used for this preparation.

2.2.3 Collection of Excipients

Carbopol 940, glycerine, propylene glycol and triethanolamine were collected at Smt. R. D. Gardi Pharmacy College, Nyara-Rajkot. PEG-40 Hydrogenated Castor Oil and DM DM Hydantoin were bought online from purenso select official website. Diethylene Glycol, Monoethyl Ether were bought from CHEMDYES CORPORATION "Rasayan Ghar" Kotharia Naka Chowk, Near Soni Bazaar RAJKOT - 360001 (Gujarat).

3. FORMULATION AND EVALUATION OF ANTIFUNGAL GEL [73]

3.1 Gel Formulation Ingredients

Formulation of topical gels is determined by important factors such as appearance, odor, Spreadability, Extrudability, viscosity, pH, texture, microbial contamination potential and bioavailability. The components of the vehicle should serve to make the skin surface more penetrable to the drug. Characteristics of the gel such as consistency and viscosity are

affected by formulation design. Consistency and viscosity affect the adhesion and retention property of the gel, and are important in ensuring the gel is retained at the site of application and effective delivery of the drug. The ingredients in topical gel formulation can be broadly categorized into four types: gelator, solvent, drug, and excipients.

3.1.1 Gelator

Gelators thicken the gel solution while preserving its flexibility by acting as stabilizers and thickeners. Gelators provide a stable internal structure to the gel when they are distributed as a colloid through the solvent. Gelators are typically selected according to their compatibility with the solvent and the intended use of the gel. The rigidity of the gel is determined by the type of gelators that are utilized. Aloe vera gel, tragacanth, gelatin, collagen, guar gum, and other natural gelators are examples of semi-synthetic gelators. On the other hand, carbomers, polyvinyl alcohol, polyethylene, and its copolymers are examples of synthetic gelators.

3.1.2 Solvent

Solvents are usually chosen based on the applications of the gel. They can be hydrophilic, lipophilic, or organic. Individual solvents can be used alone or as a mixture.^[5] Some examples of solvents include purified water, glycerin, glycols, alcohols, sucrose, toluene, and mineral oils.

3.1.3 Drug

Topical delivery is often used for drugs that are easily degraded in the GI tract, or are highly susceptible to hepatic first pass effect. Even if the drug has to be administered for long periods or can induce adverse drug reactions in parts of the body other than the target location, it can still be formulated as a topical gel.

3.1.4 Excipients

Excipients are substances that are inert to the medication and are added to dosage forms to enhance their overall quality. Antioxidants, sweeteners, stabilizers, dispersion agents, penetration enhancers, buffers, and preservatives are a few examples. Excipients known as penetration enhancers have the capacity to raise skin permeability. Numerous excipient classes, including glycerin, sulfoxides and their analogues, pyrrolidines, fatty acids and ethanol, surfactants, etc., can be employed as penetration enhancers. The pH of gels with an aqueous or hydroalcoholic basis can be adjusted by adding buffers. Because of their antibacterial properties, preservatives are crucial, but they become much more crucial throughout the hydrogel production process. The oxidation of gel components is inhibited by the application of antioxidants. When choosing the antioxidant to be used, it is important to consider the nature of the solvent. Since the solvent of most gels are aqueous in nature, water-soluble antioxidants are more commonly used. Sweetening agents are only used in gels that are designed to be used in the oral cavity such as dental gels.

3.2 Gel Preparation Methods

The process of gel formation involves finding a balance between the concentrations of the gelator and the solvent. When adding a gelator to the solvent, the mixture remains in liquid state. As the concentration of the gelator increases to a certain critical concentration, gelation occurs through swelling to form the semi-solid gel. Further increasing the concentration of the gelator beyond the gelling point will increase gel viscosity.

The exact gelling point varies depending on the properties of the gelator and the solvent, such as structure uniformity, molecular weight of the polymer, and flexibility of the polymer chain. Generally, gels are prepared by first dissolving the soluble excipients in the solvent. The gel is allowed to settle for one to two days before the final consistency of the gel can be reached. The exact method of preparing gels depends on the properties of the formulation ingredients.

3.3 Advantages of Topical gels

Topical gels have a less oily texture than creams and ointments because they contain a larger percentage of water. Due to the evaporation of the solvent, these gels have a greater spreading capacity, a cooling effect, and a longer skin retention period.

Topical gels can stick well to the application site and are more stable than creams and ointments. On the application site, they create an occlusive layer that can serve as a barrier. Their distinct composition and structure make them harmless and easily washable. Because of their confined impact, they have little adverse effects. Topical gels are straightforward to use and convenient. Topical gels also have a non-invasive topical method of action. The release profile of the gel can be modified by altering the properties of the gelator, allowing for continuous drug delivery. Topical gels are also eco-friendly, biocompatible, and biodegradable. The topical dosage form allows stable and continuous drug delivery to the site of application while having a faster drug release than ointments and creams. All these can increase the drug's bioavailability in the body. Better application properties and stability in comparison to cream and ointments.

3.4 Method for Formulation of Gel [73]

3.4.1 Dispersion Method

Gelling agent was dispersed in water with stirring at 1200 rpm for 30 min. Drug was dissolved in non-aqueous solvent with preservative, this solution was added in above gel with continuous stirring.

Phase A

Take water in a beaker, add Carbopol 940 & dissolve completely. Add glycerine on continue stirring then add PG. Mix well. Add DM DM Hydantoin on continue stirring.

Phase B

In another beaker, weigh PEG40 hydrogenated castor oil & Add Diethylene glycol monoethyl ether. Mix well until it becomes transparent. Add lavender oil, calendula oil & vitamin E one by one. Mix well until clear. Add phase B to phase A. Mix well. Check pH & adjust with Triethanolamine. The transparent gel is formed.

3.5 Evaluation of Antifungal Gel [74-78]

3.5.1 FTIR (Fourier Transform Infrared Spectroscopy)

Interactions between an API and its excipients impact the API's stability, chemical structure, and bioavailability. As FTIR spectroscopy detects vibrational shifts that may indicate possible intermolecular interactions between dosage components, it is helpful not only for studying the behavior of solid-state APIs and their excipients but also as a compatibility screening tool.

3.5.2 FTIR Analysis

FTIR study was performed and FTIR spectra were recorded on FTIR spectrophotometer with IRspirit_SHIMADZU-IR-Instrument1 at Public Testing Laboratory, GTU, Gandhinagar, Gujarat, India and Instrumental Facility Section Department of Pharmaceutical Sciences Saurashtra University Rajkot-India by MS method. FTIR spectral data were recorded and functional group stretching and bending wavelength were matched with literature published data.

3.5.3 Physical appearance

The physical attributes (color, look, and feel), organoleptic parameters (phase separation, and liquefaction), pH, viscosity, Spreadability, and oil content were also observed at various intervals for 30 days.

Appearance - Colour is important for patient compliance. The prepared gels were inspected visually for clarity, colour, and presence of any particles.

Homogeneity - All developed gels were tested for homogeneity by visual inspection after the gels had been set in the container. They were tested for their appearance and the presence of any aggregates.

pH

pH 1.0 g gel was accurately weighed and dispersed in 100 ml purified water. The pH of the dispersion was measured using a digital pH meter, which was calibrated before use with a standard buffer solution at 4.0, 7.0, and 9.0. The measurements of pH were done in triplicate and average values were calculated.

Viscosity

Brookfield DV-III ULTRA PROGRAMMABLE RHEOLOGICAL RHEOMETER and DV-II+ Pro viscometer and was used for the determination of viscosity. Gel samples were placed at room temperature for 30 min. Then, they were poured into an apparatus container. Number 64 spindle was attached then viscosity was determined at 25°C and 100–250 rpm. The results were reported on average after triplicate experiments.

Spreadability

One of the criteria for a topical formulation to meet the ideal qualities is that it should possess good Spreadability. It is the term expressed to denote the extent of the area to which formulation readily spreads on application to the skin or affected part. The therapeutic efficacy of a formulation also depends upon its spreading value. To determine the Spreadability of formulation, 0.5 g of gel was placed within a circle of 1 cm diameter pre-marked on a glass plate of 20 × 20 cm, over which a second glass plate was placed. A weight of 500 g was allowed to rest on the upper glass plate for 5 min. The increase in the diameter due to gel spreading was noted.

Extrudability

To determine Extrudability a closed collapsible tube containing formulation was pressed firmly at the crimped end. When the cap was removed, the formulation extruded until the pressure dissipated. The weight in grams required to extrude a 0.5 cm ribbon of the formulation in 10 Seconds was determined. The average extrusion pressure in g was reported.

Stability

The stability study was assessed by storing the formulation at different storage conditions and at room temperature (25-28°C).

3.5.4 Antifungal Activity

For a variety of causes, the quantity of antifungal susceptibility testing (AFST) carried out has increased recently. In recent decades, there has been a rise in the number of patients with risk factors for invasive fungal infection (IFI), such as deep immunosuppression, prolonged use of broad-spectrum antibiotics, and implanted medical devices.

This has also led to an increase in the incidence of IFIs. Clinical microbiology labs frequently use AFST as a technique to help choose the best antifungal medication. By figuring out the medication concentration needed to inhibit an organism to a certain degree—referred to as the minimum inhibitory concentration, by definition, gives an *in vitro* assessment of susceptibility and resistance. Testing for antifungal susceptibility may be useful in ways other than choosing an antifungal medication for a certain patient.

Procedure: The antifungal activity of prepared formulation was tested at Neugen Unipath-Health Checkup / Pathology lab/ Diagnostics center in Rajkot against *candida albicans* by agar diffusion method.

3.5.5 In vitro Drug Diffusion Study

The Franz diffusion cell was mainly used to assess the stability and permeability of formulations, including transdermal and topical. The in-vitro and ex-vivo drug release from topical treatments such as creams, ointments, liposome formulations, and gels can be accurately measured using the simple diffusion cell assay. It provided very important viewpoints on the interactions between the skin, medication, and formulation.

Furthermore, it is utilized for quality control and toxicity testing.

Procedure: *In vitro* drug release was determined using a Franz diffusion cell and synthetic membrane. 1 g of the test sample was dispersed uniformly on the membrane surface; finally, it was fixed on the cell. cell receiver phase contained phosphate buffer, pH 6.8. The temperature of 37°C was controlled by a pumped water bath circulating between 2 shells encompassing the chamber.

Franz diffusion cell was placed at receiver phase space by a magnetic stirrer to obtain sink conditions. This set was also put on a magnetic mixer then the cell mouth was covered by parafilm to avoid evaporation from the donor phase. A volume of 1 ml samples was taken at specified time intervals. After each sampling, the aliquots were replaced by fresh phosphate buffer, pH 6.8 subsequently to gain the same volume of receiver phase during the experiment. The test was repeated three times for each sample, and the absorbance was measured in a UV Spectrophotometer.

4. RESULTS AND DISCUSSION

4.1 FTIR Study

Based on FTIR spectral data of essential oils as well as excipients, it is confirmed that the characteristic peaks of essential oils are preserved in the gel formulations, showing the absence of any type of interaction among formulation constituents.




Figure 3: FTIR analysis

4.2 Formulations of Different Batches

Table 2: Formulas of Different Batches of Antifungal Gel

Ingredient	Batch 1	Batch 2	Batch 3	Batch 4	Batch 5	Batch 6	Batch 7
Distilled water	q.s	q.s	q.s	q.s	q.s	q.s	q.s
Carbopol 940	0.5 %	0.6%	1%	1%	1.5%	1.5%	2%
Glycerin	3%	1.6 %	2%	2%	2%	3%	3%
Propylene glycol	5%	1.6%	3%	5%	4%	5%	3%
DM DM Hydantoin	0.5%	0.5%	0.5%	0.5%	0.5%	0.5%	0.5%
PEG40 hydrogenated castor oil	4%	5%	3%	5%	5%	5%	4%
Diethylene glycol monoethyl ether	4%	5%	2%	4%	4%	5%	4%
Calendula essential oil	0.5%	1%	1%	1%	1%	1%	1%
Lavender essential oil	0.5%	0.5%	1%	1%	1%	1%	1%
Vitamin E	0.5%	0.5%	0.5%	0.5%	0.5%	0.5%	0.5%
Triethanolamine	q.s	q.s	q.s	q.s	q.s	q.s	q.s

By using all the above Formulas Antifungal Gel was prepared in different batches shown in Figure 4.




Figure 4: different batches of formulations

4.3 pH and Viscosity of All Formulations

Batch no.	pH	Viscosity (cP)
-----------	----	----------------

Table 3: pH & formulations

1	5.5	2983
2	5.7	1082
3	6.8	1297
4	6.2	1987
5	5.9	2787
6	6	1095
7	6.9	1276

Viscosity of all

The pH of all formulations was found to be in the range of 5-7 (Table 3), which was near to skin pH. Batch no. 1,2,5 and 6 showed a pH near to range of skin pH 5.

4.4 Spreadability of All Formulations

Batch 1, 2, and 4 showed better Spreadability than other batches.

4.5 Extrudability of All Formulations

Extrudability study shows batches 1,2 and 4 have good Extrudability.

4.6 Stability of All Formulations

Stability study was done at 0, 15th, and 30th days by examining all formulations of any visible and above-performed evaluation parameters regarding changes. All the formulations were found stable as none of them showed any changes in parameters. Batch no. 4 was found to be more stable than other formulations.

4.7 Antifungal Activity Against *Candida Albicans*

Antifungal activity against *candida albicans* was performed by using the agar diffusion method of batches 1,2 and 4. Batch 2 showed slight activity at different concentrations as shown in the figures. Batch 4 was found to be a better antifungal topical gel as results shown in the figures.



Figure 5: Antifungal activity of Batch 4

Table 4 Zone of inhibition in *Candida albicans* species

Batch no	Concentration (μ l)	Zone of inhibition
1	100	0
2	200-400	2.5 mm
4	100-200	5 mm

4.8 *In vitro* Drug Diffusion Study

Table 5: Drug Release of Batch 4

Time (hours)	Drug Release %
0	0
1	30
2	45
3	57
4	78
5	89
6	98

Drug release of batch 4 formulation was carried out by using a Franz cell diffusion study for 6 hours and results were determined by using a UV spectrophotometer at 413 nm wavelength.

5. CONCLUSION

The risk of fungal infections to human health is still there and getting worse. This innovative herbal topical antifungal gel, which contains essential oils of *calendula officinalis* and *lavender angustifolia*, is biocompatible and provides an alternative form of treatment for fungal infections. The gel is intended to reduce the adverse effects of standard antifungal drugs that are commercially accessible, including drug resistance and toxicity. To evaluate the patients for improvements in clinical signs and symptoms, more clinical research must be done. In summary, all evaluation parameters showed positive results for the topical antifungal gel based on calendula and lavender oil, which was prepared with carbopol 940 and other excipients. This suggests that the gel is efficient against pathogens. Out of all the formulations, batch number four demonstrated excellent results and was subsequently selected for a study on drug release, revealing drug release within six hours. To improve patient consistency, topical medications will be utilized more often in the future.

COMPETING INTERESTS

Authors have declared that they have no known competing financial interests OR non-financial interests OR personal relationships that could have appeared to influence the work reported in this paper.

Disclaimer (Artificial intelligence)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

9. REFERENCES

1. Salm S, et al. Current state of research on the clinical benefits of herbal medicines for non-life-threatening ailments. *Frontiers in Pharmacology*. 2023 Sep 28;14. Available from: <https://doi.org/10.3389/fphar.2023.1234701>
2. Wang H, et al. Advancing herbal medicine: enhancing product quality and safety through robust quality control practices. *Frontiers in Pharmacology*. 2023 Sep 25;14. Available from: <https://doi.org/10.3389/fphar.2023.1265178>
3. Seladi-Schulman J PhD. Everything you need to know about fungal skin infections. *Healthline*. 2023. Available from: <https://www.healthline.com/health/fungal-skin-infection>
4. Fungal infections of the skin. *WebMD*. 2024. Available from: <https://www.webmd.com/skin-problems-and-treatments/fungal-infections-skin>
5. Professional CCM. Fungal infections (Mycosis). *Cleveland Clinic*. Available from: <https://my.clevelandclinic.org/health/diseases/24401-fungal-infections-mycosis>
6. Wikipedia contributors. Topical gels. *Wikipedia*. 2024. Available from: https://en.wikipedia.org/w/index.php?title=Topical_gels&oldid=1209671347
7. Kasiram, K., Sakharkar, P., & Patil, A. Antifungal Activity Of *Calendula Officinalis*. *Indian Journal of Pharmaceutical Sciences*, 2000;62:464.
8. D'Auria FD, et al. Antifungal activity of *Lavandula angustifolia* essential oil against *Candida albicans* yeast and mycelial form. *Med Mycol*. 2005 Aug;43(5):391-6. doi: 10.1080/13693780400004810. PMID: 16178366.
9. Gazim ZC, et al. Antifungal activity of the essential oil from *Calendula officinalis* L. (Asteraceae) growing in Brazil. *PubMed*. 2008 Jan 1; Available from: <https://pubmed.ncbi.nlm.nih.gov/24031180>
10. Mehraliyeva, S., et al. Development of novel antibacterial gel using clove and calendula extracts with colloidal silver nanoparticles. *Eurasian Chemical Communications*, 2021; 3(3): 170-179. doi: 10.22034/ecc.2021.269869.1131
11. Ak G, et al. Chemical composition and biological activities of essential oils from *Calendula officinalis* L. flowers and leaves. *Flavour and Fragrance Journal*. 2021 Apr 30;36(5):554–563. Available from: <https://doi.org/10.1002/ffj.3661>
12. Xiong X, et al. Antifungal mechanisms of lavender essential oil in the inhibition of rot disease caused by *Monilinia fructicola* in postharvest flat peaches. *Canadian Journal of Microbiology*. 2021 Oct 1;67(10):724–736. Available from: <https://doi.org/10.1139/cjm-2020-0484>
13. Patil K, et al. A review of *Calendula Officinalis* Magic in Science. *Journal of Clinical and Diagnostic Research*. 2022 Jan 1; Available from: <https://doi.org/10.7860/jcdr/2022/52195.16024>
14. Mijatovic S, et al. Antifungal Activity of *Lavandula angustifolia* Essential Oil against *Candida albicans*: Time-Kill Study on Pediatric Sputum Isolates. *Molecules/Molecules Online/Molecules Annual*. 2022 Sep 24;27(19):6300. Available from: <https://doi.org/10.3390/molecules27196300>
15. Adaszyńska-Skwirzyńska M, Dzięcioł M, Szczerbińska D. *Lavandula angustifolia* Essential Oils as Effective Enhancers of Fluconazole Antifungal Activity against *Candida albicans*. *Molecules/Molecules Online/Molecules Annual*. 2023 Jan 25;28(3):1176. Available from: <https://doi.org/10.3390/molecules28031176>
16. Shahane K, et al. An Updated Review on the Multifaceted Therapeutic Potential of *Calendula officinalis* L. *Pharmaceuticals*. 2023 Apr 18;16(4):611. Available from: <https://doi.org/10.3390/ph16040611>
17. Maj, Wiktoria & Pertile, Giorgia & Rozalska, Sylwia & Frac, M. Assessing the Effect of *Calendula Officinalis* on *Neosartorya* spp. Mycelium Viability. 2023.
18. Paolino B, et al. *Lavandula angustifolia* mill. for a suitable non-invasive treatment against fungal colonization on organic-media cultural heritage. *Heritage Science*. 2024 Feb 14;12(1). Available from: <https://doi.org/10.1186/s40494-024-01166-9>
19. Kasar PM, Kale K, Phadtare DG. Formulation and Evaluation Of Topical Antifungal

- Gel Containing Itraconazole. International Journal of Current Pharmaceutical Research. 2018 Jul 16;10(4):71. Available from: <https://doi.org/10.22159/ijcpr.2018v10i4.28470>
20. A U, Kumari C, et al. Development and Evaluation Of Polyherbal Gel For Antifungal Activity. International Journal of Current Pharmaceutical Research. 2018 Sep 15;10(5):40. Available from: <https://doi.org/10.22159/ijcpr.2018v10i5.29694>
 21. Jayasankar P, et al. Formulation and Development of a Herbal Antifungal Gel Containing Origanum vulgare and Syzygium aromaticum Essential Oils Against Oral Candida albicans. Curēus. 2024 Feb 17; Available from: <https://doi.org/10.7759/cureus.54348>
 22. Wikipedia contributors. Calendula officinalis. Wikipedia. 2024. Available from: https://en.wikipedia.org/w/index.php?title=Calendula_officinalis&oldid=1215002221
 23. Kaviena Baskaran. Pharmacological Activities of Calendula Officinalis. International Journal of Science and Research (IJSR) ISSN: 2319-7064 Index Copernicus Value (2015): 78.96. Available from; https://en.m.wikipedia.org/wiki/Calendula_officinalis.
 24. "Calendula officinalis". Germplasm Resources Information Network. Agricultural Research Service, United States Department of Agriculture.
 25. Calendula officinalis (Calendula, Calendula (Pot Marigold), Common Marigold, Garden Marigold, Gold bloom, Holligold, Marigold, Mary Bud, Pot Marigold, Ruddles) | North Carolina Extension Gardener Plant Toolbox. Available from: <https://plants.ces.ncsu.edu/plants/calendula-officinalis>
 26. Ramos A., et al. Genotoxicity of an extract of Calendula officinalis L. J. Ethnopharmacol. 1988;61:49–55.
 27. Shahane K, et al. An Updated Review on the Multifaceted Therapeutic Potential of Calendula officinalis L. Pharmaceuticals. 2023 Apr 18;16(4):611. Available from: <https://doi.org/10.3390/ph16040611>
 28. Calendula oil - Calendula Officinalis Uses & Benefits. Available from: <https://www.essentialoil.in/calendula-oil.html>
 29. Fonseca YM, et al. Protective effect of Calendula officinalis extract against UVB-induced oxidative stress in skin: Evaluation of reduced glutathione levels and matrix metalloproteinase secretion. Journal of Ethnopharmacology 2010 Feb 1;127(3):596–601. Available from: <https://doi.org/10.1016/j.jep.2009.12.019>
 30. Hamburger M, et al. Preparative purification of the major anti-inflammatory triterpenoid esters from Marigold (Calendula officinalis). Fitoterapia. 2003 Jun 1;74(4):328–338. Available from: [https://doi.org/10.1016/s0367-326x\(03\)00051-0](https://doi.org/10.1016/s0367-326x(03)00051-0)
 31. Babae N, et al. Antioxidant capacity of calendula officinalis flowers extract and prevention of radiation induced oropharyngeal mucositis in patients with head and neck cancers: a randomized controlled clinical study. Daru. 2013 Mar 7;21(1). Available from: <https://doi.org/10.1186/2008-2231-21-18>
 32. Patil K, et al. A review of Calendula Officinalis Magic in Science. Journal of Clinical and Diagnostic Research. 2022 Jan 1; Available from: <https://doi.org/10.7860/jcdr/2022/52195.16024>
 33. Pandey P, Despande B. Antioxidant activity in the leaves and petals of Calendula officinalis Linn. Asian Pacific Journal of Health Sciences. 2022 Apr 1;9(2):130–132. Available from: <https://doi.org/10.21276/apjhs.2022.9.2.26>
 34. Elias R, et al. Antimutagenic activity of some saponins isolated from Calendula officinalis L., C. arvensis L. and Hedera helix L. Mutagenesis. 1990 Jan 1;5(4):327–332. Available from: <https://doi.org/10.1093/mutage/5.4.327>
 35. Cruceriu D, Balacescu O, Rakosy E. Calendula officinalis: Potential Roles in Cancer Treatment and Palliative Care. Integrative Cancer Therapies. 2018 Oct 5;17(4):1068–1078. Available from: <https://doi.org/10.1177/1534735418803766>
 36. Pommier P, et al. Phase III randomized trial of Calendula officinalis compared with trolamine for the prevention of acute dermatitis during irradiation for breast cancer. Journal of Clinical Oncology. 2004 Apr 15;22(8):1447–1453. Available from: <https://doi.org/10.1200/jco.2004.07.063>

37. Singh M, Bagewadi A. Comparison of effectiveness of *Calendula officinalis* extract gel with lycopene gel for treatment of tobacco-induced homogeneous leukoplakia: A randomized clinical trial. *International Journal of Pharmaceutical Investigation*. 2017 Jan 1;7(2):88. Available from: https://doi.org/10.4103/jphi.jphi_19_17
38. Deka NB, et al. Mechanism of Action of Wound Healing Activity of *Calendula officinalis*: A Comprehensive Review. *Pharmaceutical and Biosciences Journal*. 2021 Jan 3;28–44. Available from: <https://doi.org/10.20510/ukjpb/9/i1/1609684673>
39. Gunasekaran S, Nayagam AAJ, Natarajan R. Wound healing potentials of herbal ointment containing *Calendula officinalis* Linn. on the alteration of immunological markers and biochemical parameters in excision wounded animals. *Clinical Phytoscience*. 2020 Nov 23;6(1). Available from: <https://doi.org/10.1186/s40816-020-00215-7>
40. Górniak I, Bartoszewski R, Króliczewski J. Comprehensive review of antimicrobial activities of plant flavonoids. *Phytochemistry Reviews*. 2018 Oct 6;18(1):241–272. Available from: <https://doi.org/10.1007/s11101-018-9591-z>
41. Manandhar S, Luitel S, Dahal RK. In Vitro Antimicrobial Activity of Some Medicinal Plants against Human Pathogenic Bacteria. *Journal of Tropical Medicine*. 2019 Apr 2;2019:1–5. Available from: <https://doi.org/10.1155/2019/1895340>
42. Vinola SelvanathanMJ, et al. Comparative evaluation of *Calendula officinalis* and 2% chlorhexidine against *Enterococcus faecalis* and *Candida albicans*. *Journal of Interdisciplinary Dentistry*. 2021 Jan 1;11(3):119. Available from: https://doi.org/10.4103/jid.jid_28_21
43. Mali S.N., et al. *Essential Oils: Applications and Trends in Food Science and Technology*. Springer; Cham, Switzerland: 2022. *Molecular Modeling Approaches to Investigate Essential Oils (Volatile compounds) Interacting with Molecular Targets*; pp. 417–442.
44. History & Uses of Lavender - Lavender | Lavender Sense browse. Available from: <http://www.lavendersense.com/index.php/Lavender/Index>
45. De Melo Alves Silva LC, et al. Use of *Lavandula angustifolia* essential oil as a complementary therapy in adult health care: A scoping review. *Heliyon*. 2023 May 1;9(5):e15446. Available from: <https://doi.org/10.1016/j.heliyon.2023.e15446>
46. Woronuk G., et al. Biosynthesis and therapeutic properties of *Lavandula* essential oil constituents. *Planta Med*. 2011;77:7–15. doi: 10.1055/s-0030-1250136.
47. Demissie Z.A., et al. Cloning, functional characterization and genomic organization of 1, 8-cineole synthases from *Lavandula*. *Plant Mol. Biol*. 2012;79:393–411. doi: 10.1007/s11103-012-9920-3.
48. Lane A., et al. A genomics resource for investigating regulation of essential oil production in *Lavandula angustifolia*. *Planta*. 2010;231:835–845. doi: 10.1007/s00425-009-1090-4.
49. Batiha G.E.-S., et al. A Review of the Bioactive Components and Pharmacological Properties of *Lavandula* Species. *Naunyn-Schmiedeberg's Arch. Pharmacol*. 2023;396:877–900. doi: 10.1007/s00210-023-02392-x.
50. Wells R., et al. *Lavandula* Essential Oils: A Current Review of Applications in Medicinal, Food, and Cosmetic Industries of Lavender. *Nat. Prod. Commun*. 2018;13:1934578X1801301038. doi: 10.1177/1934578X1801301038.
51. Ghanimi R., et al. Antioxidant and Antibacterial Activities of Essential Oils from Three Moroccan Species (*Lavandula Mairei* Humbert, *Lavandula dentata* L. and, *Lavandula stoechas* L.) *J. Pharm. Res. Int*. 2021;33:64–71. doi: 10.9734/jpri/2021/v33i45B32779.
52. Dobros N., Zawada K., Paradowska K. Phytochemical Profile and Antioxidant Activity of *Lavandula Angustifolia* and *Lavandula x Intermedia* Cultivars Extracted with Different Methods. *Antioxidants*. 2022;11:711. doi: 10.3390/antiox11040711.
53. Cavanagh H.M.A., Wilkinson J.M. Biological Activities of Lavender Essential Oil. *Phytother. Res*. 2002;16:301–308. doi: 10.1002/ptr.1103.
54. Gismondi A., et al. The Antimicrobial Activity of *Lavandula Angustifolia* Mill.

- Essential Oil against *Staphylococcus* Species in a Hospital Environment. *J. Herb. Med.* 2021;26:100426. doi: 10.1016/j.hermed.2021.100426.
55. Sayout A., et al. Antibacterial Activity and Chemical Composition of Essential Oil from *Lavandula Tenuisecta* Coss.Ex Ball. an Endemic Species from Morocco. *Eur. J. Integr. Med.* 2020;33:101017. doi: 10.1016/j.eujim.2019.101017
 56. Zuzarte M., et al. The Anti-Inflammatory Response of *Lavandula Luisieri* and *Lavandula Pedunculata* Essential Oils. *Plants.* 2022;11:370. doi: 10.3390/plants11030370.
 57. Maurya S., et al. Interspecies Comparative Features of Trichomes in *Ocimum* Reveal Insights for Biosynthesis of Specialized Essential Oil Metabolites. *Protoplasma.* 2019;256:893–907. doi: 10.1007/s00709-018-01338-y.
 58. Prusinowska R., et al. Hydrolates from Lavender (*Lavandula angustifolia*)—Their Chemical Composition as Well as Aromatic, Antimicrobial and Antioxidant Properties. *Nat. Prod. Res.* 2016;30:386–393. doi: 10.1080/14786419.2015.1016939.
 59. Dobros N, Zawada KD, Paradowska K. Phytochemical Profiling, Antioxidant and Anti-Inflammatory Activity of Plants Belonging to the *Lavandula* Genus. *Molecules.* 2022 Dec 28;28(1):256. doi: 10.3390/molecules28010256. PMID: 36615453; PMCID: PMC9821988.
 60. Kajjari S, et al. The Effects of Lavender Essential Oil and its Clinical Implications in Dentistry: A Review. *Int J Clin Pediatr Dent.* 2022 May-Jun;15(3):385-388. doi: 10.5005/jp-journals-10005-2378. PMID: 35991803; PMCID: PMC9357533.
 61. Boligon A.A., Machado M.M., Athayde M.L. Technical evaluation of antioxidant activity. *Med Chem.* 2014;4:517–522. doi: 10.4172/2161-0444.1000188.
 62. Cavanagh HMA, Wilkinson JM. Biological activities of lavender essential oil, *Phytother Res*, 2002, vol. 16 p. 301-308.
 63. Safitri FI, Nawangsari D, Febrina D. Overview: Application of Carbopol 940 in gel. *Proceedings of the International Conference on Health and Medical Sciences (AHMS 2020).* 2021 Jan 1; Available from: <https://doi.org/10.2991/ahsr.k.210127.018>
 64. Nur Izyan Wan Azelee, et al. "Glycerol In Food, Cosmetics And Pharmaceutical Industries: Basics And New Applications" *International Journal of Scientific & Technology Research.* 2019;8(12). ISSN 2277-8616
 65. Tolu Ajiboye, What Is Propylene Glycol? 2023; <https://www.verywellhealth.com/propylene-glycol-5076050#:~:text=Propylene%20glycol%20is%20an%20emollient,well%20the%20products%20are%20absorbed.DMDM> HYDANTOIN <https://cosmetics.specialchem.com/inci-ingredients/dmdm-hydantoin>
 66. PEG40 HYDROGENATED CASTOR OIL. <https://cosmetics.specialchem.com/inci-ingredients/peg-40-hydrogenated-castor-oil>
 67. Osborne DW. Diethylene glycol monoethyl ether: an emerging solvent in topical dermatology products. *J Cosmet Dermatol.* 2011 Dec;10(4):324-9. doi: 10.1111/j.1473-2165.2011.00590.x. PMID: 22151944.
 68. ETHOXYDIGLYCOL <https://cosmetics.specialchem.com/inci-ingredients/ethoxydiglycol>
 69. Keen MA, Hassan I. Vitamin E in dermatology. *Indian Dermatol Online J.* 2016 Jul-Aug;7(4):311-5. doi: 10.4103/2229-5178.185494. PMID: 27559512; PMCID: PMC4976416.
 70. ECHEMI 2020-02-06 "The role of triethanolamine in cosmetics the effect of Triethanolamine on skin" <https://www.echemi.com/cms/85497.html#:~:text=Triethanolamine%20is%20the%20most%20commonly,effect%20of%20thickening%20and%20moisturizing>
 71. BROOKE SHUNATONA "Triethanolamine for Skin: What You Need to Know" 2022; <https://www.byrdie.com/triethanolamine-for-skin-4777052>
 72. Kanth Saanvi Essentials, New Delhi <https://m.indiamart.com/proddetail/calendula->

oil-20122626297.html

73. Available from: <https://www.slideshare.net/FarazaJaved/gels-33285405>
 74. Adriana I. Segall Preformulation: The use of FTIR in compatibility studies *Journal of Innovations in Applied Pharmaceutical Science (JIAPS)* e-ISSN: 2455-5177 Vol. 4(3), 01-06, Jul-Sep, 2019. www.jiapsonline.com
 75. Berkow EL, Lockhart SR, Ostrosky-Zeichner L. Antifungal Susceptibility Testing: Current Approaches. *Clin Microbiol Rev.* 2020 Apr 29;33(3):e00069-19. doi: 10.1128/CMR.00069-19. PMID: 32349998; PMCID: PMC7194854.
 76. Kumar, et al. A Franz diffusion cell and its implication in skin permeation studies. *Journal of Dispersion Science and Technology*, 45(5), 943–956. 2024; <https://doi.org/10.1080/01932691.2023.2188923>
 77. Jayasankar P, et al. Formulation and Development of a Herbal Antifungal Gel Containing *Origanum vulgare* and *Syzygium aromaticum* Essential Oils Against Oral *Candida albicans*. *Cureus.* 2024 Feb 17;16(2):e54348. doi: 10.7759/cureus.54348. PMID: 38500909; PMCID: PMC10945991.
 78. Harish NM, et al. Formulation and Evaluation of in situ Gels Containing Clotrimazole for Oral Candidiasis. *Indian J Pharm Sci.* 2009 Jul;71(4):421-7. doi: 10.4103/0250-474X.57291. PMID: 20502548; PMCID: PMC2865814.
-
79. PAULUS D, LUCHESI LA, BUSO C, FRATA MT, OLIVEIRA PJB DE. CHEMICAL COMPOSITION, ANTIMICROBIAL AND ANTIOXIDANT ACTIVITIES OF ESSENTIAL OILS OF FOUR SPECIES OF THE LAMIACEAE FAMILY. *EURO. J. MED. PLANTS.* [INTERNET]. 2020 JUL. 22 [CITED 2024 JUN. 7];31(10):129-40. AVAILABLE FROM: [HTTPS://JOURNALEJMP.COM/INDEX.PHP/EJMP/ARTICLE/VIEW/918](https://JOURNALEJMP.COM/INDEX.PHP/EJMP/ARTICLE/VIEW/918)
-
80. DRA LA, AGHRAZ A, BOUALY B, OUBAASSINE S, BARAKATE M, MARKOUK M, LARHSINI M. CHEMICAL CHARACTERIZATION AND IN VITRO ANTIMICROBIAL ACTIVITY OF *CARALLUMA EUROPAEA* ESSENTIAL OIL AND ITS SYNERGISTIC POTENTIAL WITH CONVENTIONAL ANTIBIOTICS. *J. ADV. MED. PHARM. SCI.* [INTERNET]. 2019 FEB. 20 [CITED 2024 JUN. 7];19(4):1-11. AVAILABLE FROM: [HTTPS://JOURNALJAMPS.COM/INDEX.PHP/JAMPS/ARTICLE/VIEW/378](https://JOURNALJAMPS.COM/INDEX.PHP/JAMPS/ARTICLE/VIEW/378)
-
81. CASSELLA S, CASSELLA JP, SMITH I. SYNERGISTIC ANTIFUNGAL ACTIVITY OF TEA TREE (*MELALEUCA ALTERNIFOLIA*) AND LAVENDER (*LAVANDULA ANGUSTIFOLIA*) ESSENTIAL OILS AGAINST DERMATOPHYTE INFECTION. *INTERNATIONAL JOURNAL OF AROMATHERAPY.* 2002 MAR 1;12(1):2-15.