

# Detection of Metallo $\beta$ -Lactamase in *Acinetobacter* species using Phenotypic Methods with their Antibiotic Resistance Profile in Tertiary Care Hospital

## Abstract

**Background:** *Acinetobacter species* are commonly found in nature, in water, and in soil, and also, and they have been isolated from humans and animals. In the hospital setting, *Acinetobacter species* has emerged as a major opportunistic pathogen, being able to colonize and develop infections in patients. In *Acinetobacter species*,  $\beta$ -lactamases and outer membrane alterations are likely to function together to impart resistance. Understanding the epidemiology, and resistance mechanism, and for infection management and preventing a potential global health crisis, techniques for identifying Carbapenem-Resistant-*Acinetobacter species* are essential. **Aim:** To detect Metallo  $\beta$ -Lactamase (MBL) in *Acinetobacter species* with their antimicrobial susceptibility pattern. **Material and Methods:** 100 *Acinetobacter species* isolates from various clinical samples such as blood, cerebrospinal fluid, sputum, bronchoalveolar lavage, pus, urine, wound swab, and body fluids were included in the study. Phenotypic tests such as, Disc Potentiation test & Modified Hodge test were used to detect the presence of Metallo  $\beta$ -lactamase in imipenem & meropenem-resistant isolates. **Results:** The highest prevalence of *Acinetobacter* was found in blood (37%), followed by sputum (31%) and others [pus, urine, ascitic fluid, cerebrospinal fluid (CSF), bronchoalveolar Lavage (BAL), and pleural fluid]. A maximum number of *Acinetobacter* were isolated from MICU followed by medicine and surgery. The Modified Hodge test yielded the largest percentage of MBL-positive *Acinetobacter* isolates (67.26%) when compared to the Disc Potentiation test method (50.45%). **Conclusion:** Early detection of Metallo  $\beta$ -lactamase production is essential for planning appropriate treatment according to the resistance mechanisms of the multi-drug resistant strains, as well as for efficient infection control procedures to prevent the spread of infection. **Keywords:** *Acinetobacter species*; Metallo  $\beta$ -lactamase; Antibiotic resistance; Disc Potentiation test (DPT); Modified Hodge test (MHT)

**Introduction:** *Acinetobacter species* are non-fastidious, strict aerobe, gram-negative coccibacilli, oxidase-negative, catalase-positive, non-fermentative, and non-motile [1,2,3]. *Acinetobacter species* are commonly found in nature, in water, and in soil and also, and they

have been isolated from animals. In humans, they are found to reside on the mucous membrane, skin, and also sometimes the membrane of the intestinal tract <sup>[3,4]</sup>. They are usually plumpy, and small, measuring 1.0-1.5  $\mu\text{m}$  by 1.5-2.5  $\mu\text{m}$  in diameter[21-24]. More than 50 species in the *Acinetobacter* genus, the *Acinetobacter baumannii* complex (*Acinetobacter nosocomialis*, *Acinetobacter pittii*, and *Acinetobacter baumannii*) are the most clinically important ones. But sometimes they develop into more coccoid form, usually present in pairs or long chains of variable duration <sup>[5]</sup>. It has led to many hospital outbreaks in the past years due to its ability to survive in the hospital environment and stay long on surfaces. In recent years, it has been labelled as a “red alert” human pathogen, expressing problems among the medical fraternity due to its wide range of antibiotic resistance <sup>[6]</sup>.

In the hospital setting, *Acinetobacter baumannii* has emerged as a major opportunistic pathogen, being able to colonize and develop infections in patients with ventilator-associated pneumonia, secondary meningitis, urinary tract infection, septicemia, and other conditions in the intensive care unit (ICU). It also causes infections in other immune-compromised individuals, including burn patients <sup>[7]</sup>.

Infections with *Acinetobacter baumannii*, particularly Carbapenem-Resistant *Acinetobacter baumannii* (CRAB), because of their high case rates, mortality, and morbidity, are of public health concern worldwide. According to the World Health Organization (WHO), carbapenem-resistant *Acinetobacter baumannii* has high rates globally in the target list of antibiotic-resistant bacteria, as a vital priority pathogen to drive drug research and development <sup>[8]</sup>. Owing to the widespread use of  $\beta$ -lactam antimicrobials, bacterial resistance has increased and is now a significant threat to the continued use of antibiotic therapy <sup>[9]</sup>. The objective of this study is to enhance the clinical management of patients suffering from infections and detection of metallo  $\beta$ -lactamase which in turn will also provide us with relevant epidemiological details.

## **Materials and methods**

This cross-sectional prospective study was conducted in the department of microbiology, MGM Medical college & hospital, Navi Mumbai, India from July 2019 to January 2021.

**Sample size** 100 *Acinetobacter species* isolates from various clinical samples such as blood, cerebrospinal fluid, sputum, bronchoalveolar lavage, pus, urine, wound swab, and body fluids were included in the study. Samples received from age group below 18 years were excluded. Antibiotic Susceptibility testing of all *Acinetobacter species* isolated was carried out by the

Modified Kirby Bauer Disc diffusion technique by using Mueller Hinton Agar (MHA) plates according to clinical and laboratory standard institute (CLSI) guidelines 2020. <sup>[10]</sup>

### **Test for Metallo $\beta$ -lactamase detection**

#### **Disc Potentiation test (DPT) <sup>[11]</sup>**

The standard suspension of test isolate was inoculated on Mueller Hinton Agar (MHA) plate. A 0.5M solution of EDTA was prepared by dissolving 186.1gm of disodium EDTA 2H<sub>2</sub>O in 1000ml of distilled water and pH was adjusted to 8.0 by using NaOH. The mixture was sterilized by autoclaving. Two 10 $\mu$ g Meropenem discs and Imipenem discs were placed 20mm apart from center to center onto MHA plate. To one disc of Imipenem and Meropenem each, 5 $\mu$ l of 0.5M EDTA was added. The zone of inhibition of Meropenem and Meropenem/EDTA discs and Imipenem and Imipenem/EDTA discs were compared after 16-18 hours of incubation at 35°C. An increase in the zone of inhibition at least by 7 mm with Meropenem/EDTA discs and Imipenem/EDTA disc than Meropenem and Imipenem alone respectively was considered as MBL positive [as shown in Fig.1].

#### **Modified Hodge test (MHT) <sup>[11]</sup>**

A 0.5 McFarland dilution of the *Escherichia coli* ATCC 25922 in 5 ml of broth or saline was prepared. In the Mueller Hinton agar (MHA) plate, 1:10 dilution was streaked as a lawn. In the center of the test region, a 10 g Imipenem disc was placed, the test organism was streaked in a straight line from the disc edge to the plate edge, and it was then kept in a bacteriological incubator at 35°C for 24-48 hours. The appearance of the enhanced ATCC *E. coli* 25922 growth along the test organism, which showed a cloverleaf-like indentation, which indicated a positive test, was used to detect the production of carbapenemase [as shown in Fig.2].

### **Results**

A total 100 isolates of Acinetobacter species collected from various clinical samples of patients admitted and also out door patients (OPD) were included in the study. Clinical samples were from all the clinical areas of the hospital. Samples received from age group below 18 years were excluded from the study.

All isolates were non-duplicate and identified by conventional methods.

### **Demographic distribution**

Age and sex-wise distribution of 100 strains of *Acinetobacter* species in various clinical samples is shown in Table 1. The maximum number of *Acinetobacter* species were isolated from male patients (79%) in both age groups as compared to the female population (21%). The highest prevalence of *Acinetobacter* was found in Blood (37%), followed by Sputum (31%) and others (Pus, Urine, Ascetic Fluid, CSF, BAL and Pleural Fluid).

### **Antimicrobial Susceptibility Profile**

All the 100 isolates of *Acinetobacter* species were subjected to antimicrobial susceptibility testing (AST) by using first line antibiotics to identify the multi-drug resistant strains. Table 2 shows the resistance pattern of *Acinetobacter* species, revealing that 88% of *Acinetobacter* species were resistant to Augmentin (AMC), followed by Nitrofurantoin (82 %), Tobramycin (68 %), Cefotaxime, and Piperacillin (67 %).

62 isolates of *Acinetobacter* species were found to be resistant to three or more antibiotic groups and considered as multi-drug resistant *Acinetobacter* species, which were further subjected to AST by using second line antibiotics as shown in Table 3. In the second-line antibiotics, *Acinetobacter* species were found to be highly resistant to Ticarcillin/ Clavulanic acid (61, 98.38%), followed by Piperacillin/Tazobactam (59, 95.16%), Meropenem (57, 91.19%) and Imipenem (56, 90.32%).

### **Phenotypic detection of Metallo $\beta$ -lactamase (MBL) production in *Acinetobacter* species:**

The phenotypic detection of Metallo  $\beta$ -lactamase (MBL) was done on those isolates of *Acinetobacter* species which were resistant to Imipenem (56) and Meropenem (57). Disc Potentiation test (DPT) and Modified Hodge test (MHT) were used for the phenotypic detection of MBL. Table 4 shows the proportion of MBL in imipenem and meropenem resistant strains of *Acinetobacter* species. The highest percentage of MBL-positive *Acinetobacter* isolates were given by the M.H.T (67.26%) method as compared to D.P.T (50.45%) method.

### **Result summary**

The study analyzed 100 non-duplicate *Acinetobacter* isolates from clinical samples of patients aged 18 and above. Most isolates were from male patients (79%) and were most commonly

found in blood (37%) and sputum (31%) samples. Antimicrobial susceptibility testing revealed high resistance rates to first-line antibiotics, with 88% resistant to Augmentin and 82% to Nitrofurantoin. Additionally, 62 isolates were multi-drug resistant (MDR), showing significant resistance to second-line antibiotics such as Ticarcillin/Clavulanic acid (98.38%) and Piperacillin/Tazobactam (95.16%). Phenotypic detection of Metallo  $\beta$ -lactamase (MBL) among Imipenem and Meropenem-resistant strains indicated that the Modified Hodge Test (MHT) identified a higher proportion of MBL-positive isolates (Imipenem: 66.07%, Meropenem: 68.42%) compared to the Disc Potentiation Test (DPT) (both around 50%). This highlights a significant challenge in treating *Acinetobacter* infections due to the high prevalence of resistance to both first and second-line antibiotics and the presence of MBL producers.

## Discussion

During the study period of July 2019 to January 2021, 100 samples of *Acinetobacter species* was isolated from all the body fluids which were included in this study. Some of the studies and their findings are presented in Table 5.

Blood had the highest prevalence of *Acinetobacter species* accounting for 37%, followed by Sputum (31%), Pus (18%), Urine (9%), and Other body fluids (1%). Our study is comparable with the study conducted in 2010 in which the highest prevalence was also found in blood (47%)<sup>[12]</sup>. In controversy, in a study done by M. Hajjar et.al, the sputum sample had the highest prevalence (43%)<sup>[13]</sup>.

In our study, when tested the antimicrobial susceptibility of the *Acinetobacter species* it was revealed that the *Acinetobacter* showed 88% resistance to Augmentin (AMC), which was followed by Tobramycin (68%). On the other hand, the *Acinetobacter species* showed the highest sensitivity to Tetracycline (TE) with a percentage of 68%. In the second-line antibiotics, the highest degree of resistance was found to be 98.38% for Ticarcillin/ Clavulanic acid (TCC) and Piperacillin/ Tazobactam (PIT). On the other hand, the highest sensitivity was found against Polymyxin B (PB) with 96.77% followed by 93.5% to Colistin (CL), 87.09% to Levofloxacin (LE), and 66.12% to Tigecycline (TGC). Another study done by **Dr. S. Das, et.al**,<sup>[14]</sup> presented the Antimicrobial Susceptibility Pattern which showed 100% susceptibility to Colistin, followed by 75% to Aztreonam and 65% sensitivity to Imipenem and Meropenem. Along with this, it was also found that the lowest sensitivity was towards Ceftazidime (21.7%).

Our present study is comparable with the study done by **M. Hajjar, et.al** <sup>[13]</sup> in which according to Antibiotic Susceptibility Testing, 95% of *Acinetobacter species* were resistant to Cefoxitin, 87% were resistant to Ciprofloxacin, 86% resistant to Trimethoprim-sulfamethoxazole and Piperacillin- tazobactam, 83% were resistant to Cefotaxime, 81% to Ceftazidime and 80% to Gentamicin. Also, 78% of the isolates of *Acinetobacter species* were resistant to Imipenem and 84% to Meropenem. There was only 1 isolate that was resistant to Colistin. In another study done in the year 2018 by **A. Kaur, et.al**, <sup>[15]</sup>, most of the antibiotics examined demonstrated high levels of resistance in *Acinetobacter baumannii isolates*. *Acinetobacter baumannii strains* were found to be resistant to Ceftazidime in 96.6% of cases, Cefepime in 94.8% of cases, Imipenem in 60.3% of cases and Meropenem in 68.1% of cases. However, Polymyxin B has a susceptibility of 96.5% and Colistin has a susceptibility of 97.4%.

In the present study, two age groups were considered to analyze their distribution. In this study, it was found that the species of *Acinetobacter* were widely distributed in the age group of 1850 (52%) as compared to 51 and above (48%). The other study done in the year 2020 by **S.K. Yadav, et.al**, <sup>[16]</sup>, included various age groups in their study. In their study, the highest distribution of *Acinetobacter species* was in the age group of 16-32 years (23.6%) followed by  $\geq 65$  years (22.4%),  $\leq 15$  years (19.8%), 33-48 years (18.7%) and 49-64 years (15.5%). The other study done in the year 2018 by **V. Rebic, et.al**, <sup>[17]</sup>, studied the distribution of *Acinetobacter species* in three age groups. The ratio of isolates was higher in the over 60 years age group ( $p= 0.763$ ). In the present study, it was discovered that the isolates of *Acinetobacter species* were more in males (79%) as compared to females (21%). *Acinetobacter* infections were more frequent in males (54.20 %) as compared to females (45.80 %). In the present study, the growth of *Acinetobacter* isolates in blood was more in the age group of 18-50 with males (12) and females (10), which was followed by sputum in which the highest number of isolates were found from the age group of 51 and above with males (16) and females (4). In the pus sample the maximum number of *Acinetobacter* was isolated from the age group of 18-50 in which males were 13 and female was 1. In the urine sample, the highest isolates were found in the age group of 51 and above with males (5) and female (1).


The present study was conducted on 100 isolates of *Acinetobacter species*, to detect the prevalence of MBL. The production of MBL was carried out on those isolates that were resistant to Meropenem (57) and Imipenem (56). MBL was detected with the help of 2 methods- Disc Potentiation Test (DPT) and Modified Hodge Test (MHT). 56 Imipenem resistant *Acinetobacter* isolates showed 50% positivity by DPT and 66.07% by MHT. Whereas,

out of 57 Meropenem-resistant *Acinetobacter species* isolates, 50.87% were positive by DPT and 68.42% by MHT. In a study done by **Zahra Moulana, et.al** <sup>[18]</sup>, in the year 2020, the use of DDST was one of the more effective techniques for the detection of Ambler class B MBL production with a high rate of positivity, according to the results of several studies. The results from the MHT showed that 84% of *A.baumannii* isolates were carbapenemase producers. Besides that, their study showed the lowest positivity rate (30%) by the DDST method. Our study is comparable with the study done in the year 2019 by **Mittal N, et.al.** <sup>[19]</sup> in which 292 (76.8%) isolates were likely MBL producers with only 72 (19%) confirmed as MBL producers.

**Figure 1:** Showing Disc Potentiation Test (DPT) positive for Metallo  $\beta$ -Lactamase (MBL)



**Figure 2:** Showing Modified Hodge Test for Metallo  $\beta$ -Lactamase (MBL) production



**Table 1:** Age-wise and Sex-wise distribution of *Acinetobacter* species in all the clinical samples

Samples	18-50 Age		51 & above		Isolates of <i>Acinetobacter</i>
	Male	Female	Male	Female	
Blood	12	10	12	3	37
Sputum	9	2	16	4	31
Pus	13	1	4	0	18
Urine	3	0	5	1	9
Ascitic Fluid	0	0	2	0	2
Cerebrospinal Fluid (CSF)	1	0	0	0	1
Pleural Fluid	0	0	1	0	1
Broncho alveolar Lavage (BAL)	1	0	0	0	1

**Table 2:** Antibiotics (first line antibiotics)

<b>Antibiotic</b>	<b>Concentration (mcg)</b>	<b>Sensitive</b>	<b>Intermediate</b>	<b>Resistant</b>
Amoxyclav (AMC)	20/ 10 mcg	10	2	88
Nitrofurantoin (NIT)	300 mcg	10	8	82
Tobramycin (TOB)	10 mcg	31	1	68
Cefotaxime (CTX)	30 mcg	30	3	67
Piperacillin (PI)	100 mcg	24	9	67
Amikacin (AK)	30 mcg	34	0	66
Ceftazidime (CAZ)	30 mcg	36	0	64
Gentamicin (GEN)	10 mcg	30	7	63
Trimethoprim- sulfamethoxazole (COT)	23.75/ 1.25 mcg	30	9	61
Ciprofloxacin (CIP)	5 mcg	34	10	56
Levofloxacin (LE)	5 mcg	64	0	36
Tetracycline (TE)	30 mcg	73	3	24

**Table 3:** Antibiotics (second line antibiotics)

<b>Antibiotics</b>	<b>Concentration (mcg)</b>	<b>Sensitive</b>	<b>Intermediate</b>	<b>Resistant</b>
Ticarcillin/ Clavulanic acid (TCC)	75/10 mcg	0	1	61
Piperacillin/ Tazobactam (PIT)	100/ 10 mcg	3	0	59
Meropenem (MRP)	10 mcg	3	2	57
Aztreonam (AT)	30 mcg	2	3	57
Imipenem (IPM)	10 mcg	5	1	56
Cefepime (CPM)	30 mcg	4	2	56

Tigecycline (TGC)	15 mcg	41	11	10
Levofloxacin (LE)	5 mcg	54	2	6
Colistin (CL)	25 mcg	58	0	4
Polymyxin B (PB)	300 units	60	0	2

**Table 4:** Metallo  $\beta$ -lactamase Production by Different Methods among the Imipenem and Meropenem Resistant Acinetobacter spp.

Antibiotics	Disc Potentiation Test (DPT)		Modified Hodge Test (MHT)	
	POSITIVE	NEGATIVE	POSITIVE	NEGATIVE
Imipenem (n=56)	28	28	37	19
Meropenem (n=57)	29	28	39	18

**Table 5:** Showing various studies and their sample-wise distribution

Sample	Present Study	S.K. Yadav, et.al 2020 [16]	Dr. S Das, et.al 2018 [14]	M Hajjar, et.al 2018 [13]	A.Rezaei, et.al,2018 [20]	F Sahcher aghi, et al 2010 [12]
Blood	37%	6.2%	13.3%	5%	3%	47%
Sputum	31%	-	3%	43%	3%	10%
Pus	18%	27.3%	38.3%	21%	9%	12%
Urine	9%	6.8%	15%	10%	-	7%
Other Body Fluids	5%		-	19%	-	
Catheter Tips	-		-	2%	1.2%	

Medical Devices	-		20%	-	-	
-----------------	---	--	-----	---	---	--

## Conclusions

The maximum number of *Acinetobacter species* was isolated from the age group 18-50 years as compared to the age group of 50 & above. The present study showed that Males had predominance over Females (4:1) in both age groups in all the samples. According to the present study, the *Acinetobacter species* showed the highest percentage of resistance to Augmentin (AMC) (88%), which was followed by Nitrofurantoin (NIT) (82%). On the other hand, the *Acinetobacter species* showed the highest susceptibility to Tetracycline (TE) (73%). In the second line of antibiotics, the highest degree of resistance was found to be for both Ticarcillin/ Clavulanic acid (TCC) and Piperacillin/ Tazobactam (PIT) i.e., 98.38%. On the other hand, the highest susceptibility was found against Polymyxin B (PB) (96.77%) followed by Colistin (CL) (93.5%), Levofloxacin (LE) (87.09%), and Tigecycline (TGC) (66.12%). The highest percentage of MBL-positive *Acinetobacter* isolates were given by the M.H.T (67.26%) method as compared to D.P.T (50.45%) method. The potential spread of these multi-drug resistant strains may be stopped by the early detection of MBL-producing *Acinetobacter* species. To find MBL producers, active monitoring is required. In district health laboratories or other locations lacking access to molecular diagnostic tools, the phenotypic tests described in the current research can be used to quickly identify these types of resistance.

## Highlights

- To identify and characterize the proportion and distribution of MBL in Imipenem and Meropenem Resistant isolates within the clinical isolates of *Acinetobacter*.
- To analyse the phenotypic characteristics of MBL producing *Acinetobacter* isolates, including their antimicrobial resistance pattern.

## Limitations

The lack of PCR analysis for the verification of phenotypical techniques constitutes one of the study's limitations. It was a uni-center study so our findings cannot be generalized to the entire region.

### Ethical Approval and Consent:

The approval of the Institutional Ethics Committee (N-EC/2019/SC/07/97) IRB name (Dr. Ipsheeta Ray & Dr. Sarita Sahani) and the written consent was obtained from all the patients.

### Disclaimer (Artificial intelligence)

#### Option 1:

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

#### Option 2:

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc have been used during writing or editing of manuscripts. This explanation will include the name, version, model, and source of the generative AI technology and as well as all input prompts provided to the generative AI technology

Details of the AI usage are given below:

- 1.
- 2.
- 3.

### References

1. Almaghrabi M, Joseph M, Assiry M, Hamid M. Multidrug-Resistant *Acinetobacter baumannii*: An Emerging Health Threat in Aseer Region, Kingdom of Saudi Arabia. *Canadian Journal of Infectious Diseases and Medical Microbiology*. 2018;2018:1-4. <https://doi.org/10.1155/2018/9182747>
2. Mathai A, Oberoi A, Madhavan S, Kaur P. *Acinetobacter* infections in a tertiary level intensive care unit in northern India: Epidemiology, clinical profiles, and outcomes. *Journal of Infection and Public Health*. 2012;5(2):145-152. <https://doi.org/10.1016/j.jiph.2011.12.002>
3. Lazim, A.J. and Al-Jumaili, E.F.A. . (2021). Production, Purification and Characterization of  $\beta$ -Lactamase from Clinical Isolate of *Acinetobacter baumannii* A7. *Indian J. of Forensic Medicine & Toxicology*. 2021; 15 (3). 1893-1900.

4. Manchanda V, Sinha S, Singh N. Multidrug resistant Acinetobacter. Journal of Global Infectious Diseases. 2010;2(3):291. <https://doi.org/10.4103/0974-777X.68538>
5. Almasaudi S. Acinetobacter spp. as nosocomial pathogens: Epidemiology and resistance features. Saudi Journal of Biological Sciences. 2018;25(3):586-596. <https://doi.org/10.1016/j.sjbs.2016.02.009>
6. Peleg A, Seifert H, Paterson D. Acinetobacter baumannii: Emergence of a Successful Pathogen. Clinical Microbiology Reviews. 2008;21(3):538-582. <https://doi.org/10.1128/CMR.00058-07>
7. Gupta V. Coexistence of Extended Spectrum Beta-Lactamases, AmpC Beta-Lactamases and Metallo-Beta-Lactamases in Acinetobacter baumannii from burns patients: a report from a tertiary care center of India. Ann Burns Fire Disasters. 2013;26(4):189-192. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3978590/>
8. Ayobami O, Willrich N, Harder T, et.al, The incidence and prevalence of hospital-acquired (carbapenem-resistant) Acinetobacter baumannii in Europe, Eastern Mediterranean, and Africa: a systematic review and meta-analysis. Emerging Microbes & Infections. 2019;8(1):1747-1759. <https://doi.org/10.1080/22221751.2019.1698273>
9. Palzkill T. Metallo- $\beta$ -lactamase structure and function. Annals of the New York Academy of Sciences. 2012;1277(1):91-104. <https://doi.org/10.4103/0974-2727.129082>
10. CLSI, Performance Standards for Antimicrobial Disk Susceptibility Test. 13<sup>th</sup> ed. CLSI standard M02. Wayne, PA: Clinical and Laboratory Standards Institute, 2018:46-49.
11. Umadevi, Sivaraman, "Detection of extended spectrum beta lactamases, ampc beta lactamases and metallo betalactamases in clinical isolates of ceftazidime resistant Pseudomonas Aeruginosa." Brazilian journal of microbiology : [publication of the Brazilian Society for Microbiology] vol. 42,4 (2011): 1284-8.
12. Shahcheraghi F, Abbasalipour M, Akbari N, et.al, P24.09 Study of beta-lactamases in clinical Acinetobacter isolates from Tehran, Iran. Journal of Hospital Infection. 2010;76:S70. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3279807/>
13. Hajjar Soudeiha M, Dahdouh E, Daoud Z, et.al, Phenotypic and genotypic detection of  $\beta$ lactamases in Acinetobacter spp. isolates recovered from Lebanese patients over a 1-year period. Journal of Global Antimicrobial Resistance. 2018;12:107-112. <https://doi.org/10.1016/j.jgar.2017.09.016>
14. Das S, Basak S. Acinetobacter baumannii complex and its beta-lactamase production: are we moving towards pre-antibiotic era? Int J Health Sci Res.

15. Rebic V, Masic N, Teskeredzic S, et.al. The Importance of *Acinetobacter Species* in the Hospital Environment. *Medical Archives*. 2018;72(3):330. <https://doi.org/10.5455/medarh.2018.72.330-334>
16. Yadav S, Bhujel R, Hamal P, Mishra S, et.al., Burden of Multidrug-Resistant *Acinetobacter baumannii* Infection in Hospitalized Patients in a Tertiary Care Hospital of Nepal. *Infection and Drug Resistance*. 2020;13:725-732 <https://doi.org/10.2147/IDR.S239514>
17. Kaur A, Singh S. Prevalence of Extended Spectrum Betalactamase (ESBL) and Metallobetalactamase (MBL) Producing *Pseudomonas aeruginosa* and *Acinetobacter baumannii* Isolated from Various Clinical Samples. *Journal of Pathogens*. 2018;:1-7. <https://doi.org/10.1155/2018/6845985>
18. Moulana Z, Babazadeh A, Eslamdost Z, et.al, Phenotypic and genotypic detection of metallo-beta-lactamases in Carbapenem-resistant *Acinetobacter baumannii*. *Caspian Journal of Internal Medicine*. 2020;11(2):171-176. 10.22088/cjim.11.2.171. PMID: 32509245; PMCID: PMC7265521.
19. Mittal N, Chhina D, Gupta U. Incidence of ESBL, AmpC  $\beta$ -lactamases and Metallo  $\beta$ -lactamase producing *Acinetobacter baumannii* in intensive care units of a tertiary care hospital from north India. *International Journal of Research in Health Sciences*. 2019;7(1):9-13. <http://www.ijrhs.org/>
20. Rezaei A, Fazeli H, Halaji M, et.al. Prevalence of metallo-beta-lactamase producing *Acinetobacter baumannii* isolated from intensive care unit in tertiary care unit in tertiary care hospitals. *Department of Microbiology, School of Medicine, Isfahan University of Medical Sciences, Iran*. 2018;30:330-336 <https://doi.org/10.7416/ai.2018.2224>
21. Mohamed ME, Hassan AT, Ahmed SMK. Carbapenem-hydrolyzing Oxacillinase Genes in Clinical Isolates of *Acinetobacter baumannii*. *Microbiol. Res. J. Int.* [Internet]. 2017 Sep. 7 [cited 2024 Jun. 12];21(3):1-8. Available from: <https://journalmrji.com/index.php/MRJI/article/view/1324>
22. Lo G, Dieng A, Ba-Diallo A, Samb M, Tine A, Mbaye Lo Ndiaye S, Karam F, Diagne-Samb H, Ngom-Cisse S, Sow Ndoeye A, Diop-Ndiaye H, Toure-Kane C, Gaye-Diallo A, Mboup S, Saad Bouh Boye C, Camara M. Molecular Epidemiology of Carbapenem-resistant *Acinetobacter baumannii* Isolates in a Senegalese University Teaching Hospital. *J. Adv. Microbiol.* [Internet]. 2022 Mar. 23 [cited 2024 Jun. 12];22(3):73-82. Available from: <https://journaljamb.com/index.php/JAMB/article/view/632>

23. Jung J, Park W. Acinetobacter species as model microorganisms in environmental microbiology: current state and perspectives. Applied microbiology and biotechnology. 2015 Mar;99:2533-48.

24. Poirel L, Bonnin RA, Nordmann P. Genetic basis of antibiotic resistance in pathogenic Acinetobacter species. IUBMB life. 2011 Dec;63(12):1061-7.

UNDER PEER REVIEW