

Description and Comparative Analysis of Status of Soil Fertility in Suwana block of Bhilwara District of Rajasthan, India

Abstract

The present research took place from Suwana block in Bhilwara district of Rajasthan for analyze the fertility of the soils on cultivated land both close to and far from industrial areas was carried out at the SHUATS, Prayagraj. Department of Soil Science and Agricultural Chemistry formerly called Naini Agricultural Institute, (U.P.). During the year 2023-2024. The soil samples were collected at the single-depth ranges: 0–15 cm from 30 different villages of one block of Bhilwara district, a total of 90 samples collected and analyzed for their physical and chemical parameter by using SLT. The outcome demonstrated that the soil samples' pH was observed from neutral, slightly alkaline to strongly alkaline. The EC⁻¹ became secure variety for cultivated land. %OC range was found in low to medium range. Available nitrogen was low, phosphorous was moderately to low., potassium levels ranged from medium to high, Sulphur was range from low too high in range in soils of Suwana block. Ca and Mg had been enough in variety of research area.

Keywords: Physical, Parameters, Suwana, Bhilwara, Standard Laboratory technique

1. Introduction

"Soil" is a loose material consisting of worn rock and other things, including partly decomposed OM. It is sometimes referred to as the soul of infinite life. It serves as a storehouse for nutrients and is essential to the development of plants and other crops that keep the environment on Earth clean. It additionally acts as a supply and sinks for atmospheric gases (Ratan et al., 2011). Soil provides habitat to millions, fuel and feed to support the basic needs of humans and animals; as the population of humans and animals grows, there is a growing need for more food production. However, the soil's ability to produce is restricted, and intrinsic qualities determine the boundaries of production., agroecological setting, use and management. In order to create a land use system that works, it is necessary to systematically assess soil resources in terms of their quantity, distribution, qualities, behaviors, and potential for use. For augmenting agricultural manufacturing on sustainable basis (FAO, 1993). "Soil trying out makes whole nutrient manipulate a possibility; fertilizer experiments are being patterned to decide economically surest costs of vitamins utility excessive yields with low manufacturing expenses according to contemporary farming, unit area is essential. Farmers of today are in different mind to the failure is more certain and sooner unless they are obtaining reasonably high yields, improved drainage, many improved cultural practices, better varieties, and control of insects and disease have helped to set the stage for high yields" (Joshi et al., 2013). Soil is formed by the weathering of rocks, and it can take up to 1000 years to develop one cm of soil. The horizons of the soil are composed of both organic and mineral materials that are split into different depths and have different morphologies, compositions, physical and chemical characteristics, and biological characteristics from the parent material. While pedology studies soil production, classification, surveying, and geological distribution, edaphology studies soil in connection to higher plants.

2. Materials and Methods

Figure 1 depicts the researched area chosen from the Bhilwara district in the Suwana block. The location of it is 25°34'07"N latitude and 74°63'13"E longitude, respectively. The area under study is part of the Central Plateau and Hills Region (VIII) Agro Climatic Zone (NARP) Sub Humid Southern Plain Zone (RJ-06), Hot Semi-Arid Eco Region Agro-Climatic Zone (Planning Commission), and Sub Agro ecological Sub Region (ICAR) Northern Plain (And Central Highlands) Including Aravalli's. Because of Rajasthan's arid geography and little rainfall, Bhilwara district has dry weather


Table 1. Characterisation of soil test value for different nutrients

Nutrients	Evaluation of the soil test results		
	Low	Medium	High
OC (%)	<0.5	0.5-0.75	>0.75
Available N (kg ha ⁻¹)	<280	280-560	>560
Available P (kg ha ⁻¹)	<10	10-25	>25
Available K (kg ha ⁻¹)	<110	110-280	>280
Available S (kg ha ⁻¹)	<10	10-20	>20
	Inadequate		Adequate
Ca ²⁺ (Cmol(P+) kg ⁻¹)	<1.5		>1.5
Mg ²⁺ (Cmol(P+) kg ⁻¹)	< 1		> 1

Its humid environment, with average rainfall and a mean annual temperature of above 22 degrees Celsius 550 mm. Using khurpi, soil samples were randomly gathered to a depth of 015 cm from farmed land in several villages within the Suwana block. After the soil samples were well combined, 500 grams of composite soil samples were collected for examination.

Using certain standard protocols for primary and macronutrients, the processed soil samples were examined for various nutrients. Using a pH meter, conductivity meter, bulk density, particle density, porosity, and pycnometer, the soil's pH, EC, and water holding capacity were all assessed. A measuring cylinder was also used to record the data.

The OM content was extracted by Walkey and Black (1934) method, available N was estimated by alkaline KMnO₄ method, available phosphorus was extracted by 0.5 M Sodium bicarbonate solution pH 8.5. Available potassium was determined by 1N NH₄Oc method, with the help of flame photometer. Sulphur was estimation by CaCl₂ method. Ex. Ca and Mg were analyzed by complexometric titration method.

Fig. 1 Location of Suwana block in Bhilwara dist. of (Raj.)

3. Result and Discussion

3.1 Physical Properties of Soil

The Status of physical properties is shown in table 2. Out of the total 90 soil samples, pH in categories as 5 (5.55 %) soil samples were neutral, 67 (87.77 %) soil samples were alkaline and 10 (5.84 %) soil samples were strongly alkaline in condition. The electrical conductivity was varied from 0.28 to 3.1 dSm⁻¹ with an average value of 0.71 dSm⁻¹. Out of 90 soil samples, 38 (31.11 %) soil samples under in safe use, 55 soil samples (66.66 %) are normal range of use and 2 sample (2.22 %) is unsafe range of use so, studies present in soil EC, no harm in germinating of seed and normal growth of plant, as maximum number of samples. The BD and PD ranges from 1.13-1.49 and 2.12-2.76 Mgm⁻³ respectively with a mean of 1.33 and 2.53 Mgm⁻³. WHC of soil ranges from 31.34 to 47.24 %, with a mean of 37.68 %. Porosity of soil ranges from 37.25 to 53.7%, with a mean of 46.57%. The data on %OM content was ranges from 0.15 to 0.69 with a mean value of 0.38. Out of 90 soil samples to 69 soil samples (76.66 %) were found low, 21 soil samples (23.33%) were found medium in OM. Thus, majority of the soil samples of Suwana block are low and few samples were medium in %OC due to the arid area is responsible low in OMaccumulation and some areas were medium in OM percent due to use of farm yard manure each year and crop residues left after harvesting. The similar results were also reported by Aggarwal and Kumar (1994), Prakash (2001), Meena *et al.*, (2006) and Yadav and Meena (2009) and Dinesh Khadka *et al.*, (2016) in ARS, Belachapi, Dhanusha, Nepal.

3.2 Status of 1⁰ nutrients in soil

The status of nitrogen, phosphorous and potassium has been shown in table2. Available nitrogen content of collected soils was ranged from 110.52 to 312.89 kg ha⁻¹ with a mean value of 199.81 kg ha⁻¹. Out of 90 collected soil samples, 85 soil samples (94.44 %) were range low, and 5 (4.67%) samples were range in medium. Climatic condition has also a major role on availability of nitrogen, max. soil samples were in low range, it may be due to denitrification

by denitrifying bacteria, arid climatic condition and application of low amount of organic manure, in situ green manures and use of chemical fertilizers. The available P content in these collected soils samples was in range from 8.5 to 32.50 Kg ha⁻¹ with a mean value of 14.21 Kg ha⁻¹. Out of 90 soil samples collected, 28 (31.11%) soil samples were in range of low, 60 (66.66%) soil samples were in range of medium and 2 (2.22%) soil samples were high in Phosphorus content. These results are strongly supported by the findings of Kumar and Seth (1983) who reported that the soil was medium in available phosphorus.

The K value in these collected soil samples was ranged of 204.32 to 383.69 kg ha⁻¹ with a mean value of 283.13 kg ha⁻¹ K. Out of 90 collected soil samples, 49 soil samples (54.44 %) were medium in range and 41 soil samples (45.55 %) were in high range, this high range of potassium is due to presence of K containing primary and secondary minerals (muscovite, biotite and anorthite).

As per criterion laid down by Muhr *et al.*, (1965), “maximum soils samples 72.72% were found under high category (above 280 kg K₂O ha⁻¹) of available potassium”. “Hence, it is found that all soils of the study area were high in available potassium”. [25]

3.3 Status of 2⁰ nutrients in soil

The status of Sulphur, Calcium and Magnesium is present in table 2. The available S content in soils of Suwana block was ranged from 4.61–19.95 kg ha⁻¹ with an average value of 12.54 kg ha⁻¹. Out of total 90 soil samples, 18 soil samples (20.00 %) were range from of low, 72 soil samples (80.00 %) were range of medium in Suwana block. The similar results were found by Jat and Yadav (2006) in soils of Entisol of Jaipur District, (Raj).

The ex. calcium content of these soils was ranges from 2.24 to 18.50 Cmol (P+) kg⁻¹ with a mean of 7.59 Cmol(P+) kg⁻¹. Out of 90 collected soil samples, all the 90 soil samples (100.00 %) were found sufficient in available Ca²⁺ content. It may be due to use of Ca²⁺ rich fertilizer and application of gypsum on the soil of studied area. The similar trend was also recorded by Nayak *et al.*, (2006) in swell and shrink soils of Vertisol order in Vidarbha region.

“The ex. magnesium value in soils of Suwana block were in range of 0.31 to 7.11 Cmol (P+) Kg⁻¹ with a mean of 2.55 Cmol (P+) kg⁻¹. Out of 90 soil samples, all the 90 soil samples (100.00%) were found sufficient in available Mg²⁺”. [25] The similar results were observed by Prasad *et al.*, (2006) in swell-shrink orange cropped soils of Nagpur District.

Table 2: Nutrients status in soil

Features of the Soil	Range	Mean	SD	CV (%)
pH (Soil: water) (1:2.5)	7.2-8.6	8.20	0.40	3.86
EC (dSm ⁻¹)	0.25-2.2	0.78	0.30	42.20
B.D. (Mg m ⁻¹)	1.13-1.49	1.34	0.08	5.68
P.D (Mg m ⁻¹)	2.12-2.77	2.54	0.16	6.52
W.H.C. (%)	23.23-54.68	37.68	6.43	7.89
Porosity (%)	37.25-52.7	47.12	3.15	6.46
OC%	0.15-0.69	0.39	0.14	36.64

Features of the Soil Available N (kg ha ⁻¹)	Range 110.52-312.89	Mean 199.81	SD 57.75	CV (%) 28.90
Available P (kg ha ⁻¹)	8.5-32.50	14.22	3.87	27.25
Available K (kg ha ⁻¹)	204.32-383.69	283.13	45.78	16.17
Features of the Soil Available S (kg ha ⁻¹)	Range	Mean	SD	CV (%)
Available Ca (Cmol (P ⁺) (kg ⁻¹))	3.71-19.95	12.54	3.29	26.20
Available M (Cmol(P ⁺) (kg ⁻¹))	2.24-18.50	12.54	3.53	46.47
	0.31-7.11	0.46	3.53	51.06

Table 3. Comparison between cultivation soils near cloth industry area, steel plant, bricks industry, sewage water irrigation and area away from industry

S.No.	Parameter	Area near industries (cultivation area)				Area far away from industries (cultivation area)
		Steel Plant	Cloth industry	Sewage water irrigation	Brick industry	
1	pH	7.81	7.63	7.65	8.25	8.19
2	EC (dSm ⁻¹)	0.62	1.17	1.07	0.64	0.65
3	BD (Mg m ⁻³)	1.42	1.31	1.32	1.35	1.33
4	PD (Mg m ⁻³)	2.63	2.37	2.42	2.56	2.55
5	WHC (%)	38.16	32.01	42.59	38.7	40.95
6	OC (%)	0.43	0.39	0.61	0.36	0.35
7	N (Kg ha ⁻¹)	192.33	192.33	240.42	165.09	200.63
8	P (Kg ha ⁻¹)	12.12	13.84	14.81	12.99	14.49
9	K (Kg ha ⁻¹)	237.63	271.93	306.31	282.29	285.06
10	S (Kg ha ⁻¹)	14.30	13.83	11.46	13.54	12.14
11	Ca (Cmol(p ⁺)Kg ⁻¹)	5.14	6.19	13.09	9.37	7.59

	1)					
12	Mg (Cmol(p+)Kg⁻¹) 1)	2.90	2.29	3.24	2.44	2.49

Conclusion

The Suwana block of the Bhilwara district's soil is classified as neutral to alkaline with a slightly saline reaction; the soluble salt level is within a safe limit and has no effect on crop germination.

In the soils of the examined area, %OC is in the low to medium range.

The research areas' soils have low levels of N, P, and S, medium levels of S, and medium to high levels of K. There was also enough magnesium and calcium found in the Suwana block's soils. With the exception of WHC, which is higher in sewage water irrigated areas than in nonindustrial areas, all areas display comparable B.D. and P.D. values

Disclaimer (Artificial intelligence)

Option 1:

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

Option 2:

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc have been used during writing or editing of manuscripts. This explanation will include the name, version, model, and source of the generative AI technology and as well as all input prompts provided to the generative AI technology

Details of the AI usage are given below:

- 1.
- 2.
- 3.

References

1. AberaDonisLikisa, Kefyalew Assefa Gejea 2016. Assessment of Physicochemical Properties of Soils as Influenced by Different Land Uses in Bedele Area in Ilubabor Zone, Southwestern Ethiopia. *International Journal of Agriculture Innovations and Research* 5(4): 2319-1473
2. Bhuyan, S.I. et al. 2013. Soil nutrients status in prominent agroecosystems of East Siang district, Arunachal Pradesh. *International Journal of Environmental Sciences*. 3(5).
3. Chaurasia et al. (2013). Study of macro nutrient and physical status of soil in a part of Varuna River in Varanasi, India. *International Journal of Environmental Sciences*. 4(4).
4. Dwivedi, S.K., Sharma, V.K. and Bhardwaj, V. 2005. Status of available nutrients in soils of cold arid region of Ladakh. *Journal of the Indian Society of Soil Science*, 53: 421– 423.
5. Jayaprakash, R., Shetty, V. Punitha B. C. and Shilpashree, V. M. 2012. Status of macro nutrients in non-traditional arecanut growing soils. *Indian Journal of Fundamental and Applied Life Sciences*, 2(1): 59 -62.
6. Lal R, Singh BR. Effects of Soil Degradation on Crop Productivity in East Africa. *Journal of sustainable Agriculture*. 1998; 13(1):15-36.

7. Klute A, Dirksen C. Hydraulic conductivity of saturated soils. In method of soil Analysis, ed. A. Klute. American Society of Agronomy and Soil Science Society of America, Madison United States. 1986, 694-700.
8. Mishra, D., & Singh, Y. V. (2019). Micronutrients status and their relationship with soil physico-chemical properties of Naugarh block, Chandauli district, Uttar Pradesh, India. *Journal of Pharmacognosy and Phytochemistry*, 8(4), 528-531.
9. Akbari, K.N., Karan, F. and Panday, H.R. 1993. Distribution of available P, K and organic carbon in Mewar Region. *Annals of Arid Zone*, 32: 251-252.
10. Black, G.R., 1965. Particle density and bulk density. In C.A. Black (ed). *Method of soil analysis part I*. Am. Soc. Agron. Madison WI. Pp. 371–390.
11. Ghosh, A.B., and Hussan, R. 1976. Available potassium status of Indian soil. *Bull. Indian Soc. Soil Sci.*, 12: 1-18.
12. Hasan, R., 1996. Phosphorus status of soils in India. *Better Crop International*, 10(2): 4-5.
13. Jat, J.R., and B.L. Yadav, Different forms of sulphur and their relationship with properties of Entisols of Jaipur district (Rajasthan) under mustard cultivation. *J. Indian Soc. Soil Sci.*, 2006, 54(2): 208-212.
14. Jatav, G.K., Mishra VN 2012. Evaluation of soil fertility status of available N, P & K in Inceptisol of Baloda block of Janigar district of Chhattisgarh. *J ProgAgric*3: 28—32.
15. Meena, H.B., Sharma, R.P. and Rawat, U.S. 2006. Status of macro and micronutrients in some soils of Tonk district of Rajasthan. *J. Indian Soc. Soil Sci.* 54(4): 508- 512.
16. Muhr, G. R., Datta, N. P., SankaraSubramoney, N., Dever, F., Lecy, V. K. and Donahue, R. R. 1965. Soil testing in India. USAID. Mission to India, New Delhi.
17. Nayak, A.K., Chinchmalatpure, R. Rao, G. Gururaja and A.K. Verma, Swell-shrink potential of vertisols in relation to clay content and exchangeable sodium under different Ionic Environment. *J. Indian Soc. Soil Sci.*, 2006, 54(1): 1-5.
18. Olsen, S.R., Cole CV, Watanabe FS, 1954. Estimation of available phosphorus in soil by extraction with sodium by carbonate. *Circular US Dep of Agric: 1400 Independence Ave SW Washington DC*. Pp. 939.
19. Singh, M.K., Raina, S.K., Sharma, P. and Kumar, B. 2011. Status of available major and micronutrients in arid soils of Churu district of western Rajasthan. *Journal of the Indian Society of Soil Science*, 59(2): 188-192.
20. Soni, Rajbala and Bhaskar, R. 2012. Impact of steel industry waste on Physico-Chemical property of soil, *International Journal of Environmental Sciences*, 2(3).
21. Walkley, A., and Black, I. A. 1934. An examination of Degtjareff method for determining soil organic matter and a proposed modification of the chromic acid titration method. *Soil Science*, 37: 29-37.
22. Vyas, V.S., 1996, Diversification in Agricultural: Concept, Rational and Approaches, *Indian Journals of Agricultural Economics*, 51(3), pp. 636-643.
23. Verma, V.K., Patel, L.B., Toor, G.S. and Sharma, P.K. (2005) Spatial Distribution of Macronutrients in Soils of Arid Tract of Punjab, India. *International Journal of Agriculture & Biology*, 7(1): 370- 372.
24. Tiwari, R.C. 2016, *Geo Figurey of India*, Prawalika Publication Allahabad. 6, pp.59-62.
25. O.P. Gurjar, Ramawatar Meena and Achinkumar and Surya Kant. Soil Fertility Status of Mandal Block in Bhilwara District of Rajasthan, India. *Int.J.Curr.Microbiol.App.Sci* (2017) 6(9): 1154-1158