

EFFECT OF MID-SEASON MOISTURE STRESS ON FATTY ACID COMPOSITION OF GROUNDNUT GENOTYPES

Commented [HT1]: Add scientific name after groundnut for modify the title.

ABSTRACT

A field experiment was conducted at RARS, Tirupati, Acharya N.G. Ranga Agricultural University during *rabi*, 2022 and 2023 to study the effect of mid-season moisture stress on fatty acid composition of groundnut genotypes. The experiment was carried out in randomized block design with factorial concept with two main treatments *i.e.*, control and mid-season moisture stress (40-80 DAS) and 12 sub treatments *i.e.*, two check varieties and ten contrasting groundnut genotypes sown in three replications. The study revealed that among the groundnut genotypes tested for their tolerance to mid-season moisture stress, the genotypes, TCGS 2326, TCGS 1792, K9 and TCGS 2333 were found to withstand moisture stress and maintained higher fatty acid composition *i.e.*, palmitic acid, stearic acid, oleic acid and lower linoleic acid indicating tolerance to moisture stress during both the years.

Commented [HT2]: Full form ?

Commented [HT3]: Also write (RBD)

Key words: Mid season moisture stress, Fatty acid composition, Groundnut

Commented [HT4]: Compare with ? Lowest was found in what ? Also at higher how many percent and add at par.

Commented [HT5]: Add more keywords.

1. INTRODUCTION

Groundnut is an important oilseed crop in the world and cultivated mainly for its edible oil, protein and fatty acids. Groundnut kernel contains high protein content (22-30 %), edible oil (44-56 %) and fatty acids among which oleic acid, linoleic acid and palmitic acid constitutes for 90 % of the total fatty acid composition in groundnut (Chakraborty *et al.*, 2013). Oleic acid and linoleic acid ratio (O/L) determine the durability and quality of groundnut. High-oleic groundnut has longer shelf-life and better flavor quality than low-oleic groundnut. Groundnut kernel also consists of minor fatty acids such as stearic acid, arachidic acid, eicosenoic acid, behenic acid, lignoceric acid and gadoleic acid constitutes for 1 and 3% of the total fatty acid composition (Abadya *et al.*, 2021).

Commented [HT6]: Spelling

In India, as 70 % of groundnut is grown as *rainfed* crop, it is subjected to intermittent soil moisture stress where yield and quality of groundnut is reduced. To improve the nutritional quality of groundnut under rainfed condition, it is essential to study the effect of soil moisture stress on composition of fatty acids (Chaiyadee *et al.*, 2013).

There are only limited reports on effect of drought on quality of groundnut kernels. Therefore, the objective of this research was to evaluate the effect of mid-season moisture stress on fatty acid composition of groundnut genotypes with different levels of drought resistance.

2. MATERIAL AND METHODS

The experiment was carried out during *rabi*, 2022 and 2023 at RARS, Tirupati, Acharya N.G Ranga Agricultural University. The experimental site is geographically located at 13.5° N latitude and 79.5° E longitude and at an altitude of 182.9 m above the mean sea level (MSL) in the Southern Agro-Climatic zone of Andhra Pradesh, India. The experiment was laid out in randomized block design with factorial concept with two main treatments and twelve sub treatments in three replications. Main treatments include control and mid-season moisture stress (40-80 DAS) and sub treatments *i.e.*, two check varieties *i.e.*, K6 (drought susceptible) and K9 (drought tolerant) and ten contrasting groundnut genotypes procured from Dept. of Genetics and Plant Breeding, RARS, Tirupati. Control treatment was maintained by providing irrigation while, mid-season moisture stress treatment was imposed by growing plants in rainout shelters by withholding irrigation from 40-80 DAS. On the completion of stress period, irrigation was provided to both control and rainout shelters. Observations of fatty acids such as palmitic acid, stearic acid, oleic acid and linoleic acid were recorded at harvest during both the years.

2.1 Estimation of fatty acids

Four fatty acids *i.e.*, palmitic acid, stearic acid, oleic acid and linoleic acid were estimated after harvest in both control and stress samples using near infrared reflectance spectroscopy *i.e.*, NIRS (FOSS DZ 2500, Denmark).

Importance:

Near infrared reflectance spectroscopy (NIRS) is a more rapid non-destructive technique for screening of large population of seed for analysis of desirable changes in the fatty acid

Commented [HT7]: Full form ??

Commented [HT8]: District ??????????

composition.

Commented [HT9]: References ???

Methodology:

A sample of 100 g of well matured dried kernels were used for fatty acids estimation in near infrared spectroscopy *i.e.*, NIRS (FOSS DZ 2500, Denmark) using ISI scan Nova software. Seed sample was loaded in a round cup which was filled up sufficiently to allow good absorption of the incident light and percentage of fatty acids was recorded.

2.2 STATISTICAL ANALYSIS

The data were analyzed statistically by following analysis of variance technique for randomized block design with factorial concept outlined by Panse and Sukhatme (1985). The statistical hypothesis of equalities of treatment means was tested by F- test at 1 to 5% per cent level of significance.

3. RESULTS AND DISCUSSION

3.1 Palmitic acid (%)

The mean data pertaining to palmitic acid composition was recorded at harvest in kernel of groundnut genotypes grown under mid-moisture stress conditions during *rabi*, 2022 and 2023 were presented in the Table 1. During *rabi*, 2022 significant differences were observed among moisture stress treatments with highest palmitic acid under irrigated conditions (control) (11.16 %) compared to moisture stress (10.61 %) with 5.18 % increase in palmitic acid over stress. Our results are in contrary with Dwivedi *et al.* (1996) who reported non-significant differences among water deficit treatments for palmitic acid. During *rabi*, 2023 non-significant differences were observed among moisture stress treatments. Results are in accordance with the reports of Dwivedi *et al.* (1996). During *rabi*, 2022 among the genotypes tested, TCGS 2326 recorded significantly highest palmitic acid (12.67 %) which was on par with K9 (11.96 %), TCGS 1792 (11.87 %) and TCGS 2333 (11.83 %). Whereas, lowest palmitic acid was recorded by TCGS 2024 (9.51 %) which was on par with TCGS 1785 (9.62 %), K6 (9.84 %) and TCGS 1784 (10.03 %). During *rabi*, 2023 significant differences were observed among the genotypes with highest palmitic acid composition in TCGS 2326 (12.99 %) which was statistically at par with TCGS 1792 (12.56 %), K9 (12.11 %), TCGS 2019 (11.75 %) and TCGS 2333 (11.71 %). While lowest palmitic acid was recorded in TCGS 2024 (10.00 %) which was on par with TCGS 1785 (10.10 %), K6 (10.15 %), TCGS 1784 (10.19 %), TCGS 1707 (10.58 %) and TCGS 2020 (10.65 %). Similar genotypic variability for

palmitic acid was reported by Hashim *et al.* (1993). No significant interactions were found among moisture stress treatments and genotypes with regard to palmitic acid during *rabi*, 2022 and 2023. Similar results of non-significant interactions were reported by Abadya *et al.* (2021).

3.2 Stearic acid

The mean data regarding stearic acid of groundnut genotypes influenced by mid-season moisture stress was presented in the Table 1. During *rabi*, 2022 and 2023 no significant differences were observed among the moisture stress treatments with respect to stearic acid. The obtained results are in contrary with the findings of Abadya *et al.* (2021) who found significant differences among moisture stress treatments. During *rabi*, 2022 significantly highest stearic acid was recorded in TCGS 1792 (3.19 %) which was on par with TCGS 2326 (3.14 %), TCGS 2333 (3.04 %), K9 (3.01 %), TCGS 2019 (2.94 %) and TCGS 1789 (2.90 %). Significantly lowest stearic acid composition was recorded by TCGS 2024 (2.47 %) which was on par with TCGS 1784 (2.52 %), TCGS 1785 (2.61 %), K6 (2.62 %) and TCGS 2020 (2.78 %). During *rabi*, 2023 significantly highest stearic acid was recorded by TCGS 2326 (3.34 %) which was on par with TCGS 1792 (3.16 %), K9 (3.14 %), TCGS 2333 (3.11 %), TCGS 2019 and TCGS 1789 (3.04 %). Whereas, lower stearic acid was recorded by TCGS 1785 (2.47 %) which was on par with TCGS 2024 (2.56 %) and K6 (2.73 %). Similar results of genotypic differences for stearic acid were reported by Dwivedi *et al.* (1996). During *rabi*, 2022 and 2023 non-significant interactions were observed between treatments and genotypes for stearic acid. Similar non-significant interactions were reported by Hashim *et al.* (1993).

Table 1. Effect of mid-season moisture stress on quality parameters of groundnut genotypes

Genotypes	Palmitic acid (%)						Stearic acid					
	2021-22			2022-23			2021-22			2022-23		
	T ₀	T ₁	Mean	T ₀	T ₁	Mean	T ₀	T ₁	Mean	T ₀	T ₁	Mean
TCGS 1707	10.77	10.01	10.39	10.76	10.40	10.58	2.86	2.75	2.81	2.75	2.92	2.83
TCGS 1784	10.20	9.85	10.03	10.26	10.12	10.19	2.63	2.40	2.52	2.80	2.90	2.85
TCGS 1785	10.16	9.08	9.62	10.15	10.05	10.10	2.51	2.71	2.61	2.56	2.38	2.47
TCGS 1789	11.27	11.15	11.21	11.27	11.33	11.30	2.99	2.82	2.90	3.00	3.08	3.04
TCGS 1792	11.75	12.00	11.87	12.90	12.22	12.56	3.12	3.25	3.19	3.18	3.14	3.16

Commented [HT10]: Add figure/ graph of this table

TCGS 2019	11.53	11.00	11.27	11.74	11.75	11.75	2.95	2.93	2.94	3.08	3.01	3.04
TCGS 2020	10.48	10.34	10.41	10.47	10.82	10.65	2.76	2.80	2.78	2.93	2.93	2.93
TCGS 2024	10.02	9.00	9.51	10.01	10.00	10.00	2.54	2.40	2.47	2.40	2.72	2.56
TCGS 2326	12.90	12.43	12.67	12.90	13.09	12.99	3.10	3.18	3.14	3.25	3.42	3.34
TCGS 2333	12.40	11.26	11.83	11.53	11.88	11.71	3.08	3.00	3.04	3.17	3.06	3.11
K6	10.27	9.42	9.84	10.20	10.10	10.15	2.68	2.56	2.62	2.71	2.75	2.73
K9	12.18	11.75	11.96	12.17	12.05	12.11	3.10	2.91	3.01	3.13	3.15	3.14
Mean	11.16	10.61		11.20	11.15		2.86	2.81		2.91	2.95	
	T	G	TXG	T	G	TXG	T	G	TXG	T	G	TXG
SEm±	0.12	0.3	0.43	0.18	0.44	0.63	0.04	0.11	0.15	0.05	0.12	0.17
CD (P=0.05)	0.353	0.865	NS	NS	1.267	NS	NS	0.306	NS	NS	0.336	NS
CV (%)	6.84			9.75			9.28			9.86		

T₀: Control, T₁: Mid-season moisture stress (40-80 DAS)

3.3 Oleic acid

Data pertaining to oleic acid composition in groundnut genotypes influenced by mid-season moisture stress were presented in Table 2. During *rabi*, 2022 and 2023 no significant differences were observed among treatments with regard to oleic acid composition. Our results were in contrary with those obtained by Chakraborty *et al.* (2013). During *rabi*, 2022 among the genotypes, TCGS 1792 significantly recorded highest composition of oleic acid (53.02 %) which was on par with TCGS 2333 (52.63 %), K9 (50.19 %) and TCGS 2326 (49.88 %). Lowest oleic acid composition was observed in TCGS 1785 (40.05 %) which was on par with TCGS 2024 (40.97 %), TCGS 1784 (41.80 %), K6 (43.33 %) and TCGS 1707 (44.08 %). During *rabi*, 2023 significantly highest oleic acid was recorded by TCGS 1792 (52.67 %) which was on par with TCGS 2333 (50.22 %), TCGS 2326 (49.25 %) and K9 (48.56 %). Whereas, lowest oleic acid was recorded by TCGS 1785 (40.01 %) which was statistically on par with TCGS 2024 (41.80 %), K6 (43.10 %), TCGS 1784 (44.15 %) and TCGS 1707 (44.67 %). Our results also concur with the published reports of Misra and Nautiyal (2005). Interaction between treatments and genotypes was non-significant for oleic acid during *rabi*, 2022 and 2023. Similar, non-significant interactions was reported by Abadya *et al.* (2021).

3.4 Linoleic acid

Mean data pertaining to linoleic acid of groundnut genotypes grown under mid-season moisture stress was presented in Table 2. During *rabi*, 2022 and 2023 moisture stress treatment recorded significantly highest linoleic acid (28.11 % and 27.43 %) compared to irrigated condition

(control) (26.06 % and 25.89 %). Moisture stress treatment recorded 7.29 % and 5.61 % increase in linoleic acid compared to control condition. Similar results of significant differences were observed under drought stress by Hashim *et al.* (1993). During *rabi*, 2022 among the genotypes tested, highest linoleic acid was recorded by TCGS 2024 (37.89 %) followed by K6 (30.35 %) and TCGS 1785 (29.92 %). Whereas, lowest linoleic acid composition was recorded by TCGS 1792 (20.17 %) which was statistically on par with TCGS 2333 (22.23 %), followed by and K9 (23.80 %). During *rabi*, 2023 highest linoleic acid composition was recorded by K6 (37.12 %) followed by TCGS 2024 (30.50 %). While lowest linoleic acid composition was recorded in TCGS 2326 (20.13 %) which was at par with TCGS 2333 (20.26) and TCGS 1792 (21.39 %). Similar results were reported by Misra and Nautiyal (2005). During *rabi*, 2022 under control condition, highest linoleic acid was recorded by TCGS 2024 (35.37 %) followed by TCGS 1785 (29.30 %) and lowest linoleic acid was recorded by TCGS 2326 (19.82 %) which was on par with TCGS 1792 (20.73 %) and K9 (21.89 %) and significantly highest linoleic acid was observed in TCGS 2024 (40.41 %) followed by K6 (32.65 %) and lowest linoleic acid was recorded by TCGS 1792 (19.62 %) which was statistically on par with TCGS 2333 (20.63 %). Our results also concur with the published reports by Hashim *et al.* (1993). During *rabi*, 2023 non-significant interactions were found among treatments and genotypes. Obtained results are in contrary with the reports of Hashim *et al.* (1993).

Table 2. Effect of mid-season moisture stress on quality parameters of groundnut genotypes

Genotypes	Oleic acid (%)						Linoleic acid (%)					
	2021-22			2022-23			2021-22			2022-23		
	T ₀	T ₁	Mean	T ₀	T ₁	Mean	T ₀	T ₁	Mean	T ₀	T ₁	Mean
TCGS 1707	45.65	42.52	44.08	46.11	43.22	44.67	27.30	27.09	27.19	27.33	26.25	26.79
TCGS 1784	38.68	44.92	41.80	44.09	44.21	44.15	26.88	28.88	27.88	27.52	30.68	29.10
TCGS 1785	39.25	40.85	40.05	39.14	40.88	40.01	29.30	30.54	29.92	29.05	31.00	30.03
TCGS 1789	47.25	48.00	47.63	46.29	47.21	46.75	25.89	26.35	26.12	26.24	27.21	26.73
TCGS 1792	52.75	53.29	53.02	52.40	52.95	52.67	20.73	19.62	20.17	21.99	20.79	21.39
TCGS 2019	46.04	46.50	46.27	46.12	46.01	46.06	26.56	26.40	26.48	26.00	26.12	26.06
TCGS 2020	45.37	45.00	45.19	45.01	45.09	45.05	27.13	30.36	28.75	27.02	27.00	27.01
TCGS 2024	40.09	41.85	40.97	43.59	40.00	41.80	35.37	40.41	37.89	28.00	33.00	30.50
TCGS 2326	48.47	51.28	49.88	47.66	50.85	49.25	19.82	28.66	24.24	20.52	19.75	20.13

Commented [HT11]: Add figure/ graph of this table

TCGS 2333	53.25	52.00	52.63	51.23	49.21	50.22	23.84	20.63	22.23	19.90	20.63	20.26
K6	43.64	43.02	43.33	44.36	41.85	43.10	28.06	32.65	30.35	33.25	41.00	37.12
K9	50.18	50.21	50.19	47.55	49.57	48.56	21.89	25.70	23.80	23.84	25.69	24.77
Mean	45.88	46.62		46.13	45.92		26.06	28.11		25.89	27.43	
	T	G	TXG	T	G	TXG	T	G	TXG	T	G	TXG
SEm±	0.66	1.62	2.29	0.68	1.67	2.36	0.39	0.94	1.34	0.44	1.08	1.53
CD (P=0.05)	NS	4.609	NS	NS	4.478	NS	1.097	2.688	3.801	1.26	3.086	NS
CV (%)	8.57			8.88			8.54			9.96		

T₀: Control, T₁: Mid-season moisture stress (40-80 DAS)

4. CONCLUSION

Variation in fatty acid composition of groundnut genotypes was assessed over 2 years in two check varieties and ten groundnut genotypes grown under mid-season moisture stress condition (40-80 DAS). The genotypes TCGS 2326, TCGS 1792, K9 and TCGS 2333 recorded higher palmitic acid, stearic acid, oleic acid and lower linoleic acid indicating its tolerance to mid-season moisture stress by maintaining better composition of fatty acids. Whereas, genotypes TCGS 2024, TCGS 1785, TCGS 1784 and TCGS 1707 recorded poor kernel quality performance in terms of fatty acids composition indicating its susceptibility to drought stress. The tolerant genotypes identified in the present study can be further evaluated for other kernel quality traits along with physiological and yield traits under moisture stress conditions for confirming their tolerance.

Commented [HT12]: Add author contribution, conflict of interest and acknowledgement

REFERENCES

1. Abadya S, Shimelis H, Pasupuleti J, Mashilo J, Chaudhari S, Manohar, SS. Assessment of the genetic diversity of groundnut (*Arachis hypogaea* L.) genotypes for kernel yield, oil and fodder quantity and quality under drought conditions. *Crop Science*. 2021;1-18.
2. Chaiyadee S, Jogloy S, Songsri P, Singkham N, Vorasoot N, Sawatsitang P, Holbrook CC, Patanothai A. Soil moisture affects fatty acids and oil quality parameters in peanut. *International Journal of Plant Production*. 2013;7(1): 82-95.
3. Chakraborty K, Bishi SK, Singh AL, Kalariya KA, Kumar L. Moisture deficit stress affects yield and quality in groundnut seeds. *Indian journal of plant physiology*. 2013;18: 136-141.

Commented [HT13]: Add more references are compulsory for quality of paper. Add more than 20 references.

4. Dwivedi SL, Nigam, SN, Rao RCN, Singh U, Rao KVS. Effect of drought on oil, fatty acids and protein contents of groundnut (*Arachis hypogaea L.*) seeds. *Field Crops Research*. 1996;125-133.
5. Hashim IB, Koehlerv PE, Eitenmiller RR, Kvien CK. Fatty acid composition and tocopherol content of drought stressed florunner peanuts. *Peanut Science*. 1993;20: 21-24.
6. Misra JB, Nautiyal PC. Influence of imposition of soil-moisture deficit stress on some quality components of groundnut, *Arachis, hypogaea, L.* kernel. *Journal of Oilseeds Research*. 2005; 22(1): 119-124.
7. Panse VG, Sukhatme PV. Statistical methods for agricultural workers, ICAR, New Delhi; 1985.

Commented [HT14]: Font of journal and color of typing is different which indicate the copy and paste. Don't do like that.

UNDER PEER REVIEW