

Review Article

Exosomes: A potential biomarker and therapeutic target for the treatment of diseases

Abstract

Nearly all types of cells release exosomes (30-120 nm, nanosized vesicles), both in diseased and healthy physiological situations. Initially, it was believed that they acted just as “garbage bags” allowing cells to expel undesired components. Exosome composition and function have been the subject of extensive research since 2007. They are expelled from the majority of cell types and are present in the majority of bodily fluids, such as blood, urine, saliva, breast milk, semen, ascitic fluid, and cell culture media. The study has shown that, the exosomal miRNA and lncRNA content in diseased patients and healthy persons varies. Scientists have successfully discovered in several studies on disease cells that they work as biomarkers, having particular proteins connected to sick pathology. This is due to their role in expressing RNAs, DNAs, and proteins from cell to cell and their presence in most bodily fluids. Exosome-based diagnostic procedures are now being used for the early diagnosis of cancer, diabetes, neurological illnesses, and other conditions. Exosomes have the ability to traverse the blood-brain barrier and may be utilised to deliver therapeutic agents such as proteins, small compounds, viral gene therapy, RNA treatments, and CRISPR gene editing.

Keywords: Garbage bags, miRNA, lncRNA, ascitic fluid, neurodegenerative, biomarker

1) Introduction

In recent years, scientific research has been uncovering the remarkable potential of exosomes – tiny vesicles secreted by cells – in various fields of medicine and biology. One area where exosomes are showing tremendous promise is animal science. These minuscule packages of biological information have been identified as key players in intercellular communication, immune response modulation, and tissue regeneration. As our understanding of exosomes deepens, their applications in animal science continue to expand, holding the promise of revolutionizing diagnostics, therapies, and overall animal health. The majority of eukaryotic cells create exosomes, membrane-bound extracellular vesicles (EVs), in the endosomal compartments .⁴⁰ They are phospholipid bilayer-enclosed vesicles and cannot proliferate like a cell. Their main function is to regulate the communications between cells of both normal and diseased states and are required for cell growth and homeostasis in multicellular organisms. These are ubiquitous in all species and they are found in various biofluids (i.e., blood, urine, cerebrospinal fluid, saliva, breast milk, semen, and ascitic fluid). EVs transport cargo proteins, lipids, nucleic acids or RNA from donor cells to receiver cells via biofluids as well as information⁴⁶. EVs assist in the removal of cellular waste.²⁷ Extracellular vesicles (EVs) have recently significantly contributed to the study and management of several disorders. In light of the fact that they carry several biomarkers, EVs are important for diagnostic purposes in many diseases, like cancer, infectious and neurodegenerative diseases, cardiovascular , diseases and pregnancy.³² A recent study also shows that EVs are capable of acting as naturally occurring drug delivery vehicles.




Fig 1: Exosomes as disease carrier & Drug delivery

2) History and Classification:

Rose, Johnstone, and group in 1983^{17, 28} and Stahl and group in 1983¹² made the initial discovery of exosomes in the growing mammalian reticulocyte (immature red blood cell). Rose, Johnstone, and the group later gave the word "exosomes" to these particles. Extracellular vesicles (EVs) include exosomes as a subclass.

Depending on the biogenesis and size of the EVs, exosomes (30–120 nm)³⁴, microvesicles (120–1000 nm), and apoptotic bodies (1–5 μm) might all be categorised.

3) The Normal physiological function of Exosomes:

Exosomes have shown immense potential in various areas of regenerative medicine. One of the most exciting applications is in tissue regeneration. Exosomes derived from stem cells have been found to promote tissue repair and regeneration in various animal models. These exosomes can enhance cell proliferation, reduce inflammation, and stimulate the formation of new blood vessels, all crucial factors in tissue regeneration.

In addition to tissue regeneration, exosomes also hold promise in wound healing.³⁰ Studies have shown that exosomes derived from platelets, known as platelet-derived exosomes, can accelerate the healing process and improve the quality of healed tissue. These exosomes contain growth factors and other signaling molecules that promote cell migration and proliferation, leading to faster wound closure and improved tissue remodeling.

By the majority of different body cell types exosomes are released through biological liquids such as synovial fluid, breast milk, blood, urine, saliva, and semen, demonstrating that they are crucial for intercellular interaction and inducing physiological reactions. Exosomes are crucial molecules with effector properties that control the body's typical normal physiological processes, including maintenance of stem cell, healing and repairing of tissues, and immune monitoring (activating immune response or suppressing inflammatory factors) and blood coagulation by providing the surface for coagulation factors, neovascularization, repairing of damaged tissues, wound healing and angiogenesis.³⁰ Exosomes are also called a 'garbage bag' due to involving in unnecessary proteins excretion out of cells, during the life cycle of cells. Exosomes are currently understood to have a role in many biological functions, including the maturation of erythrocytes. Exosomes produced by antigen-presenting cells (APCs) can also activate MHC class I and II molecules on the cell surface, thereby helping in the activation of CD8⁺ and CD4⁺ T cells and the induction of certain immunological responses.⁴¹ Exosomes released by platelets have a role in the inflammatory response by transporting prostaglandins¹³. Exosomes either activate or inhibit regulatory T cell function, reduce NK and CD8⁺ cell activity, or stimulate monocyte, B, and NK cell activity, thus in early pregnancy embryo can be protected from the attack of the host immune system, exosomes also help in embryo implantation. Exosomes are able to transport nucleic acids in addition to proteins and lipids between recipient cells⁴³. Mast cell-secreted exosomes contain small RNA and mRNA, which are then transferred to specific recipient cells and translated into the recipient cell. In general, the actions of exosomes promote target-specific cell-to-cell communication throughout the body.

4) Role in diseased conditions in humans and animals:

Exosomes have been linked to multiple common disorders in addition to normal biological functions. Exosomes' macromolecular constituents are important for cellular processes and pathological conditions like cancer, neurological disorders, angiogenesis, inflammation, and immunological responses¹⁶. In cancer, tumour-excreted exosomes help to grow tumours more rapidly and help in spreading. They inhibit the CD4 T cell and inhibit the action of NK cells hence inhibiting its cytotoxic activity thus, suppressing the immune response. Those oncogenic proteins and nucleic acids, such as TSG 101, viral oncogene proteins and viral materials, and c-Met oncoprotein, are transferred by exosomes² derived from cancer cells to create a pro-tumorial microenvironment. These molecules modulate the activity of recipient cells and play key roles in tumorigenesis, cell proliferation, progression, metastasis, and drug resistance. Exosomes were discovered to be responsible for transporting misfolded proteins from unhealthy neurons to nearby cells in neurodegenerative diseases such as Prion disease, Alzheimer's disease, Parkinson's disease, Creutzfeldt-Jakob disease in humans, bovine spongiform encephalopathy in cattle, and scrapie in sheep⁴⁴. This spreads the disease from cell to cell. Cells infected with HIV-1 produce exosomes, which are nonviral nanovesicles. The latter are intracellularly formed 50–100 nm lipid bilayer vesicles that result from endosome membrane invagination.

i. Role in cancer spread:

Exosomes could contribute in several ways to the development of cancer. They can alter both the local tumour environment and the overall environment to promote the development, spread, and early stages of metastasis in cancer cells^{14, 15}. Tumour cells can share ontogenically active proteins via exosomes inside the tumour microenvironment.

ii. Role in Human Breast cancer:

RAB22A, a membrane-bound protein that functions as a molecular switch to integrate intracellular signalling and membrane trafficking events, was discovered to be expressed by hypoxia-inducible factors, which, in turn, increased the release of microvesicles during studies of human breast carcinoma cell lines (MCF-7, MDA-MB-231, and MDA-MB-435) in low oxygen environments. Because of this, RAB22A may be a viable new therapeutic target for stopping the invasion and metastasis of cancer cells.³³

iii. Role in Neurodegenerative diseases:

The formation of amyloid plaques in the brain is a hallmark of Alzheimer's disease (AD)⁵. There is evidence that these amyloid β (Aβ) molecules travel to other neuronal cells in the patient's brain through exosomes. This conclusion has been further supported by the discovery that these molecules are physically linked to exosomes. In addition, it has been demonstrated that the exosomal marker Alix is concentrated in the brains of AD patients as

compared to the control (healthy) individuals, in whom Alix is almost nonexistent¹. Another neurological pathogenesis associated with exosomes is Parkinson's disease. A rise in α -synuclein accumulation has been linked to the disease's development. The accumulation lag time is shortened when neuroblastoma exosomes are present. Exosomes offer a favourable environment for the accumulation of α -synuclein, which is stimulated by the exosomes' lipids¹¹.




Fig 2: Sources and release of Exosomes and their function

5) Therapeutic applications or uses of Exosomes:

Extensive study has been done to explore the potential of exosomes as diagnostic and therapeutic agents since they can be extracted from practically any cell, are engaged in cell-to-cell communication, and play a role in both healthy and pathological processes.

a) Disease diagnostics (Biomarkers):

Exosomes contain and express infectious RNA and proteins, enabling the diagnosis of infectious disorders as well as the detection of active and latent types of intracellular infection. They are used to identify tumours in individuals with prostate cancer and breast cancer. For instance, swine PRRS might be detected using PRRSV-infected pigs that have the traditional tetraspanin exosomal markers (CD63 and CD81).²⁴ In dairy cows, exosomes contain unique proteins FLOT1 and TSG101 suggesting that the animal is suffering from uterine disease or metabolic disorder⁴. Staphylococcus aureus infected bovine mastitis-derived exosomes showed 5 mi RNAs (miR-2339, miR-213p, miR-23a, miR-365-3p and miR-92a)⁶. Horses with anaemia may have a marker for regeneration due to TfR1 expression in serum exosomes.³¹ The lncRNA colorectal neoplasia differentially expressed-h (CRNDE-h) found in the serum exosomes of CRC patients may be employed as a possible biomarker.²² MiR-92b-3p and miR-17-5p may also be used as potential circulating biomarkers for the early detection of swine pregnancy⁴⁸. Patients with heart failure (HF) have been shown to have elevated levels of the miRNAs miR-22, miR-320a, miR-423-5p, and miR-92b in their blood and serum exosomes¹⁰. These miRNAs can be employed as specific biomarkers for the diagnosis and prognosis of systolic HF. A-synuclein (Alzheimer's disease) has also been identified as a biomarker of neurodegeneration in cerebrospinal fluid, in addition to A-, T-, and P-Tau (Parkinson's disease).¹⁸

b) Biomarker of metabolic disorder in cows:

Dairy cows are most susceptible to metabolic diseases during the lactation to pregnancy period. Circulating exosomes may act as biomarkers to detect at-diseased cows to improve the productivity and health of the cows. Circulating exosomes are produced by a variety of organs and tissues that affect the behaviour of target tissues or cells. Exosomes in circulation provide a sensitive liquid biopsy of the physiological state. In dairy cows, exosomes contain unique proteins FLOT1 and TSG101 suggests that the animal is suffering from uterine disease or metabolic disorder³.

6) Drug delivery vehicle (Therapeutic agents)

Exosomes are not limited to promoting tissue regeneration and wound healing. They also have the potential to treat a wide range of diseases. For example, in neurodegenerative diseases such as Parkinson's and Alzheimer's, exosomes can deliver therapeutic molecules directly to affected brain cells, potentially slowing down disease progression or even reversing damage. Similarly, in heart failure, exosomes derived from cardiac cells can promote cardiac repair and regeneration, offering new hope for patients with this debilitating condition.

By releasing growth factors, proteins, miRNA, mRNA, non-coding RNA, and lipids, exosomes have been demonstrated to have the therapeutic potential to stimulate tissue regeneration. For instance, exosomes from stem cells and endothelial progenitor cells have been shown to repair heart tissue and neovascularization in a myocardial infarction and kidney injury model. RNAs were transported through exosomes reported in 2007⁴³. Lung-trapped MSCs release TSG-6, an anti-inflammatory protein that is helpful for myocardial infarction and is transported via exosomes.²⁰ Exosomes generated from MSCs encourage cell division and prevent skin cells from dying. Many anticancer drugs may be used via the exosomal carrier. Due to biocompatibility, they're easily absorbed and deliver drugs successfully. Many anticancer drugs, anti-inflammatory drugs, and anti-neurodegenerative disease drugs are successfully inserted inside the exosomes or bound with exosomes. Infections including toxoplasmosis, diphtheria, TB, and atypical severe acute respiratory syndrome are among the diseases for which exosomes are promising candidates for vaccines.⁴⁷

i. Treatment of Inflammatory Brain Disorders:

The development of CNS treatments has been complicated by the blood-brain barrier, which prevents the practical application of potentially potent medicinal medicines in the treatment of several brain cell disorders where inflammation plays a causal role. Anti-inflammatory drugs like curcumin or JSI124 are successfully carried to the brain via exosomes (intranasally) without producing noticeable adverse effects⁴⁹. They reported that intranasal administration of Exo-cur (exosome encapsulated curcumin) resulted in a significant decrease in the number of microglial cells and that Exo-JSI124 (exosome encapsulated cucurbitacin I) enhanced tumour apoptosis and a parallel decrease in disease development in all models.

ii. Gene therapy

Gene therapy can be performed using exosomes. Exosomes' transport capability can be employed to create anticancer treatments. The immune-stimulatory effects of anticancer medicines or RNAs may be reduced, and their hydrophilic properties may make it easier for them to pass cell membranes during gene therapy.³⁵

iii. Anti-inflammatory activity:

According to Sun et al. (2010), exosomal curcumin can lessen the production of IL-6 and TNF- in LPS-induced (i/p) mice.³⁶ 16 hours after the LPS injection, the IL-6 and TNF- concentrations in the sera were assessed using an ELISA assay. Exosomal curcumin (@4mg/kg) and LPS (@18.5mg/kg, Sigma-Aldrich) were administered intraperitoneally to C57BL/6j mice.

iv. Agents used as exosomal content for drug delivery: There are mainly 3 types:

- a) **Small molecules:** eg Doxorubicin (DOX)⁴², Paclitaxel¹⁹ chemotherapy medications that are frequently employed to treat a variety of malignancies. But prolonged use of DOX can cause cytotoxicity. Paclitaxel⁴⁵: a chemotherapeutic drug used to treat ovarian cancer as well as lung cancer, breast cancer, and pancreatic cancer.
- b) **Large Molecules:** eg. Curcumin, Antioxidant Curcumin: Studies are showing that the anti-inflammatory, antioxidant, endothelial, neuroprotective, and anticancer effects of curcumin-primed and curcumin-encapsulated exosomes are mediated by several intercellular and regulatory signalling systems^{26,36}. Rhodamine-12⁴⁵, Withaferin²⁵ use to control tumour growth.
- c) **Nucleic Acids:** si-RNA is utilised in genetic treatment to disrupt target genes⁸. In order to regulate post-transcriptional gene expression, mi-RNA⁹ binds to complementary regions on targeted mRNA. Exosomes are employed as therapeutic delivery systems for miRNA since they contain miRNA naturally. CRISPR/Cas9 systems have been utilised by researchers in recent years to cure a variety of genetic disorders, including cancer, by correcting, eliminating, or silencing specific genetic abnormalities linked to the disease.²¹

7) Classification of agents loaded within exosomes:




Fig. 3 Exosomes as drug delivery system

i) Types of methods perform to collect and purify exosomes

Table 1: Methods of Exosomes Collection and Purification:

Methods	Mechanisms	Specificity
---------	------------	-------------

Differential centrifugation	centrifugal force-based	Exosome isolation from bodily fluids is a common practice. ²³
Size exclusion chromatography	based on the force of gravity and material porosity	With this technique, particular beads are used for a particular size, and centrifugal force-sensitive vesicles are used to isolate them. This approach allows the simultaneous running of several biological samples ⁷ .
Filtration	based on the porosity and membrane materials	Separate exosomes from soluble chemicals and tiny particles very easily ⁷ .
Polymer-based precipitation	based on precipitation and polymer materials	Precipitation benefits include the use of neutral pH and minimal effect on separate exosomes ³⁷ .
Immunological separation	on the basis of antibody-receptor interaction	techniques for characterising and quantifying a protein implicated with specific exosome subtypes. ³⁸

Among all the methods differential-centrifugation is the most commonly used and traditional method.²³ Below table for exosome collection based on the centrifugation method:

Table 2: Four consequent centrifugation steps for exosome isolation³⁹:

Sl no	Centrifugation steps	Duration	Temperature	Purpose
1.	300g	10 min	4°C	Remove cells,
2.	2000g	10 min	4°C	Dead cells,
3.	10,000g	30 min	4°C	Cellular debris
4.	100,000g	1 hr 10 min	4°C	Collect exosome fraction

A repeated 100,000 g centrifugation of the re-suspended pellet is frequently employed to separate the exosome preparation from the lower mobility fractions, usually from free proteins.

- ii) **Exosome purification:** Both for disease detection and disease therapy purpose exosomes are collected in many different methods and purifying them is the essential stage for further clinical application and basic research. Based on the size, shape, density, and surface proteins of exosomes, many exosome separation techniques have been developed, including differential centrifugation, size-exclusion chromatography, filtration, polymer-based precipitation, and immunological separation techniques. To further the study of exosomes and their use, it is necessary to further develop these techniques, which have both benefits and drawbacks.

iii) **Exosome drug loading:**

Therapeutic agents must efficiently be loaded into the exosomes, for effective use as a drug delivery system of exosomes. Exosomes contain therapeutic substances that can either be actively loaded or passively loaded/encapsulated.

1. **Passive loading:** These techniques are actually simple and basic. Both drug incubation with sample exosomes and drug incubation with donor cells might result in passive loading.²⁹ By passive loading, Munagala and his colleagues have been able to encapsulate chemotherapeutic and chemopreventive drugs including paclitaxel, doxorubicin, and withaferin in exosomes derived from bovine milk.²⁵

2. **Active loading:** Actively disrupt of exosome membrane so the compound can diffuse inside. Mainly used for loading large molecules. Active drug loading can be performed by:

- i) **Sonication:** In order to enable the drug to penetrate into exosomes without damaging membrane-bound proteins, the exosomes' membrane integrity is impaired.²⁹
- ii) **Extrusion:** It is done by a syringe-based lipid extruder, disrupts the membrane and enhances drug diffusion.
- iii) **Electroporation:** In this technique Cells are subjected to an electrical field to improve the cell membrane's permeability, enabling the introduction of compounds, medications, or DNA within the cell.
- iv) **Drug conjugation techniques:** mostly applied in pharma industries.

8) Future Expectations

The future of exosome research in animal science is filled with promise. Researchers are working on refining techniques for isolating and characterizing exosomes, which will enhance our understanding of their biological functions and potential applications. As the technology advances, exosome-based therapies may become mainstream in veterinary medicine, offering safer and more effective treatment options for animals.

9) Conclusion:

Exosomes are unlocking new dimensions in our understanding of animal biology and health. Their ability to facilitate intercellular communication, modulate immune responses, and carry therapeutic payloads holds great promise for the field of animal science. Exosomes have great biocompatibility, making them interesting as therapeutic carriers. However, there are risks and issues, such as immunosuppression, and the development of tumours. Recent research demonstrates that one of the most popular medication delivery techniques uses both liposomes and polymeric nanoparticles, and they're causing toxicity, immunogenicity, less biocompatibility and long-distance travel issues. Because exosomes are able to carry several kinds of cargo molecules, including bioactive proteins, lipids, and nucleic acids, they have become an essential medication delivery method, and specific drugs for a variety of disorders, including solid tumours, bone regeneration, cardiac diseases, Parkinson's disease, and others. As research progresses and technology evolves, exosomes may very well reshape the landscape of diagnostics, treatments, and overall animal well-being. The journey into this microscopic realm promises exciting discoveries that will benefit animals and humans alike.

10) References

1. Aguzzi A, Rajendran L. The transcellular spread of cytosolic amyloids, prions, and prionoids. *Neuron*. 2009;64(6):783-90.
2. Ailawadi S, Wang X, Gu H, Fan GC. Pathologic function and therapeutic potential of exosomes in cardiovascular disease. *Biochimica et Biophysica Acta (BBA)-Molecular Basis of Disease*. 2015;1852(1):1-1.
3. Almughlhiq FB, Koh YQ, Peiris HN, Vaswani K, Holland O, Meier S, Roche JR, Burke CR, Crookenden MA, Arachchige BJ, Reed S. Circulating exosomes may identify biomarkers for cows at risk for metabolic dysfunction. *Scientific Reports*. 2019;9(1):13879.
4. Almughlhiq FB, Koh YQ, Peiris HN, Vaswani K, McDougall S, Graham EM, Burke CR, Arachchige BJ, Reed S, Mitchell MD. Proteomic content of circulating exosomes in dairy cows with or without uterine infection. *Theriogenology*. 2018;114:173-9.
5. Bellingham SA, Guo BB, Coleman BM, Hill AF. Exosomes: vehicles for the transfer of toxic proteins associated with neurodegenerative diseases?. *Frontiers in physiology*. 2012;3:124.
6. Cai M, He H, Jia X, Chen S, Wang J, Shi Y, Liu B, Xiao W, Lai S. Genome-wide microRNA profiling of bovine milk-derived exosomes infected with *Staphylococcus aureus*. *Cell Stress and Chaperones*. 2018;23:663-72.
7. Cheruvanky A, Zhou H, Pisitkun T, Kopp JB, Knepper MA, Yuen PS, Star RA. Rapid isolation of urinary exosomal biomarkers using a nanomembrane ultrafiltration concentrator. *American Journal of Physiology-Renal Physiology*. 2007;292(5):F1657-61.
8. Faruqu FN, Xu L, Al-Jamal KT. Preparation of exosomes for siRNA delivery to cancer cells. *JoVE (Journal of Visualized Experiments)*. 2018; 5(142):e58814.
9. Gong C, Tian J, Wang Z, Gao Y, Wu X, Ding X, Qiang L, Li G, Han Z, Yuan Y, Gao S. Functional exosome-mediated co-delivery of doxorubicin and hydrophobically modified microRNA 159 for triple-negative breast cancer therapy. *Journal of nanobiotechnology*. 2019;17(1):1-8.
10. Goren Y, Kushnir M, Zafrir B, Tabak S, Lewis BS, Amir O. Serum levels of microRNAs in patients with heart failure. *European journal of heart failure*. 2012;14(2):147-54.
11. Grey M, Dunning CJ, Gaspar R, Grey C, Brundin P, Sparr E, Linse S. Acceleration of α -synuclein aggregation by exosomes. *Journal of Biological Chemistry*. 2015;290(5):2969-82.
12. Harding C, Stahl P. Transferrin recycling in reticulocytes: pH and iron are important determinants of ligand binding and processing. *Biochemical and biophysical research communications*. 1983;113(2):650-8.
13. Heijnen HF, Schiel AE, Fijnheer R, Geuze HJ, Sixma JJ. Activated Platelets Release Two Types of Membrane Vesicles: Microvesicles by Surface Shedding and Exosomes Derived From Exocytosis of Multivesicular Bodies and α -Granules. *Blood, The Journal of the American Society of Hematology*. 1999;94(11):3791-9.
14. Hood JL, Pan H, Lanza GM, Wickline SA. Paracrine induction of endothelium by tumor exosomes. *Laboratory investigation*. 2009;89(11):1317-28.
15. Hood JL, San RS, Wickline SA. Exosomes released by melanoma cells prepare sentinel lymph nodes for tumor metastasis. *Cancer research*. 2011;71(11):3792-801.
16. Howitt J, Hill AF. Exosomes in the pathology of neurodegenerative diseases. *Journal of Biological Chemistry*. 2016;291(52):26589-97.

17. Johnstone RM, Adam M, Hammond JR, Orr L, Turbide C. Vesicle formation during reticulocyte maturation. Association of plasma membrane activities with released vesicles (exosomes). *Journal of Biological Chemistry*. 1987;262(19):9412-20.
18. Kang JH, Irwin DJ, Chen-Plotkin AS, Siderowf A, Caspell C, Coffey CS, Waligórska T, Taylor P, Pan S, Frasier M, Marek K. Association of cerebrospinal fluid β -amyloid 1-42, T-tau, P-tau181, and α -synuclein levels with clinical features of drug-naive patients with early Parkinson disease. *JAMA neurology*. 2013;70(10):1277-87.
19. Kim MS, Haney MJ, Zhao Y, Yuan D, Deygen I, Klyachko NL, Kabanov AV, Batrakova EV. Engineering macrophage-derived exosomes for targeted paclitaxel delivery to pulmonary metastases: in vitro and in vivo evaluations. *Nanomedicine: Nanotechnology, Biology and Medicine*. 2018;14(1):195-204.
20. Lee RH, Pulin AA, Seo MJ, Kota DJ, Ylostalo J, Larson BL, Semprun-Prieto L, Delafontaine P, Prockop DJ. Intravenous hMSCs improve myocardial infarction in mice because cells embolized in lung are activated to secrete the anti-inflammatory protein TSG-6. *Cell stem cell*. 2009;5(1):54-63.
21. Lin Y, Wu J, Gu W, Huang Y, Tong Z, Huang L, Tan J. Exosome-liposome hybrid nanoparticles deliver CRISPR/Cas9 system in MSCs. *Advanced Science*. 2018;5(4):1700611.
22. Liu T, Zhang X, Gao S, Jing F, Yang Y, Du L, Zheng G, Li P, Li C, Wang C. Exosomal long noncoding RNA CRNDE-h as a novel serum-based biomarker for diagnosis and prognosis of colorectal cancer. *Oncotarget*. 2016;7(51):85551.
23. Livshits MA, Khomyakova E, Evtushenko EG, Lazarev VN, Kulemin NA, Semina SE, Generozov EV, Govorun VM. Isolation of exosomes by differential centrifugation: Theoretical analysis of a commonly used protocol. *Scientific reports*. 2015;5(1):17319.
24. Montaner-Tarbes S, Pujol M, Jabbar T, Hawes P, Chapman D, Portillo HD, Fraile L, Sánchez-Cordón PJ, Dixon L, Montoya M. Serum-derived extracellular vesicles from African swine fever virus-infected pigs selectively recruit viral and porcine proteins. *Viruses*. 2019;11(10):882.
25. Munagala R, Aqil F, Jeyabalan J, Gupta RC. Bovine milk-derived exosomes for drug delivery. *Cancer letters*. 2016;371(1):48-61.
26. Oskouie MN, Aghili Moghaddam NS, Butler AE, Zamani P, Sahebkar A. Therapeutic use of curcumin-encapsulated and curcumin-primed exosomes. *Journal of cellular physiology*. 2019;234(6):8182-91.
27. Oves M, Qari HA, Felemban NM, Khan AA, Rehan M, Tabrez S, Ahmed F, Haque A, Khan MS, Khan JM, Husain FM. Exosomes: a paradigm in drug development against cancer and infectious diseases. *Journal of Nanomaterials*. 2018;2018.
28. Pan BT, Johnstone RM. Fate of the transferrin receptor during maturation of sheep reticulocytes in vitro: selective externalization of the receptor. *Cell*. 1983;33(3):967-78.
29. Patil SM, Sawant SS, Kunda NK. Exosomes as drug delivery systems: a brief overview and progress update. *European Journal of Pharmaceutics and Biopharmaceutics*. 2020;154:259-69.
30. Qin J, Xu Q. Functions and application of exosomes. *Acta Pol Pharm*. 2014;71(4):537-43.
31. Rout ED, Webb TL, Laurence HM, Long L, Olver CS. Transferrin receptor expression in serum exosomes as a marker of regenerative anaemia in the horse. *Equine veterinary journal*. 2015;47(1):101-6.
32. Schorey JS, Cheng Y, Singh PP, Smith VL. Exosomes and other extracellular vesicles in host-pathogen interactions. *EMBO reports*. 2015;16(1):24-43.
33. Schwartz SL, Cao C, Pylypenko O, Rak A, Wandinger-Ness A. Rab GTPases at a glance. *Journal of cell science*. 2007;120(22):3905-10.
34. Simons M, Raposo G. Exosomes-vesicular carriers for intercellular communication. *Current opinion in cell biology*. 2009;21(4):575-81.
35. Steinbichler TB, Dudás J, Skvortsov S, Ganswindt U, Riechelmann H, Skvortsova II. Therapy resistance mediated by exosomes. *Molecular cancer*. 2019;18(1):1-1.
36. Sun D, Zhuang X, Xiang X, Liu Y, Zhang S, Liu C, Barnes S, Grizzle W, Miller D, Zhang HG. A novel nanoparticle drug delivery system: the anti-inflammatory activity of curcumin is enhanced when encapsulated in exosomes. *Molecular therapy*. 2010;18(9):1606-14.
37. Takov K, Yellon DM, Davidson SM. Comparison of small extracellular vesicles isolated from plasma by ultracentrifugation or size-exclusion chromatography: yield, purity and functional potential. *Journal of extracellular vesicles*. 2019;8(1):1560809.

38. Tauro BJ, Greening DW, Mathias RA, Ji H, Mathivanan S, Scott AM, Simpson RJ. Comparison of ultracentrifugation, density gradient separation, and immunoaffinity capture methods for isolating human colon cancer cell line LIM1863-derived exosomes. *Methods*. 2012;56(2):293-304.
39. Théry C, Amigorena S, Raposo G, Clayton A. Isolation and characterization of exosomes from cell culture supernatants and biological fluids. *Current protocols in cell biology*. 2006;30(1):3-22.
40. Théry C, Witwer KW, Aikawa E, Alcaraz MJ, Anderson JD, Andriantsitohaina R, Antoniou A, Arab T, Archer F, Atkin-Smith GK, Ayre DC. Minimal information for studies of extracellular vesicles 2018 (MISEV2018): a position statement of the International Society for Extracellular Vesicles and update of the MISEV2014 guidelines. *Journal of extracellular vesicles*. 2018;7(1):1535750.
41. Théry C, Zitvogel L, Amigorena S. Exosomes: composition, biogenesis and function. *Nature reviews immunology*. 2002;2(8):569-79.
42. Tian Y, Li S, Song J, Ji T, Zhu M, Anderson GJ, Wei J, Nie G. A doxorubicin delivery platform using engineered natural membrane vesicle exosomes for targeted tumor therapy. *Biomaterials*. 2014;35(7):2383-90.
43. Valadi H, Ekström K, Bossios A, Sjöstrand M, Lee JJ, Lötvall JO. Exosome-mediated transfer of mRNAs and microRNAs is a novel mechanism of genetic exchange between cells. *Nature cell biology*. 2007;9(6):654-9.
44. Vella LJ, Greenwood DL, Cappai R, Scheerlinck JP, Hill AF. Enrichment of prion protein in exosomes derived from ovine cerebral spinal fluid. *Veterinary immunology and immunopathology*. 2008;124(3-4):385-93.
45. Yang T, Martin P, Fogarty B, Brown A, Schurman K, Phipps R, Yin VP, Lockman P, Bai S. Exosome delivered anticancer drugs across the blood-brain barrier for brain cancer therapy in *Danio rerio*. *Pharmaceutical research*. 2015;32:2003-14.
46. Zaborowski MP, Balaj L, Breakefield XO, Lai CP. Extracellular vesicles: composition, biological relevance, and methods of study. *Bioscience*. 2015;65(8):783-97.
47. Zacharias E, Milton G, Momen-Heravi F, Hu J, Zhang X, Wu Y. Therapeutic uses of exosomes. *J Circulat Biomarkers*. 2013;1(1).
48. Zhou C, Cai G, Meng F, Xu Z, He Y, Hu Q, Zheng E, Huang S, Xu Z, Gu T, Hu B. Deep-sequencing identification of MicroRNA biomarkers in serum exosomes for early pig pregnancy. *Frontiers in Genetics*. 2020;11:536.
49. Zhuang X, Xiang X, Grizzle W, Sun D, Zhang S, Axtell RC, Ju S, Mu J, Zhang L, Steinman L, Miller D. Treatment of brain inflammatory diseases by delivering exosome encapsulated anti-inflammatory drugs from the nasal region to the brain. *Molecular Therapy*. 2011;19(10):1769-79.