

Influence of biotic and abiotic factors on pest incidence and natural enemies in guava (*Psidium guajava* L)

ABSTRACT

A study on seasonal incidence of insect pests and their natural enemies in different guava varieties was carried out at Horticultural College and Research Institute, Periyakulam during 2022-23. The experimental results revealed that tea mosquito bug (8.75%), fruit borer (0.75%), mealybugs (12.25% and 2.80 %) and aphids (2.40%) were less in L-46. followed by red flash (1.55- 21.35%), L-49 (0.86- 13.50%) in guava ecosystem. The guava variety, Lalit was found to be more susceptible to tea mosquito bug (22.60%), fruit borer (2.65%), mealy bug (25.60%), spiraling whitefly (23.45%), scales (5.75%) and papaya Mealybug (12.70%). The coccinellids population was high in lalit (1.80/tree) followed by Allahabad safeda (1.70/tree) and L-49 (1.60/tree), whereas the spiders were found high in red flash guava (0.70/tree) but in lalit it was 0.15 per tree. The fruit fly damage, mealybugs, spiraling whitefly, aphids, scales, papaya mealybug and spiders was more during hot weather and was positively correlated with maximum and minimum temperature whereas they are negatively correlated with other weather parameters viz., rainfall, wind speed and relative humidity. The fruit borer incidence was recorded high during rainy days (0.103) with high wind speed (0.344) and minimum temperature (0.004). The coccinellids beetle were more during cloudy and rainy period (0.765), high wind speed (0.554) and relative humidity (0.338).

Keywords: Guava, Pests, Natural enemies, Seasonal incidence, Biotic and abiotic factors

Introduction

Guava is cultivated commercially in India and is popular as all season fruit with rich nutritional and medicinal value. ~~reasonable price, suitability for transport and consumer preference.~~ The ~~guava~~ area under tropical and subtropical regions are expanding due to its high ~~requirement demand~~ for fresh fruits and processed products in global trade. Guava is the fourth widely cultivated fruit crop in India with 0.15 million hectare, producing 1.80 million tonnes of fruit (Mitra *et al.*, 2008) and presently its world's top guava-producing country, with a growing area of 3.14 lakh ha and 4.92 Million tons of production (Amit Kumar Goswami, *et al.*, 2024) . Insect pests infestation is the most significant limiting factor of guava production which deteriorate yield and market value of fruit . Various insect species cause damage to guava and their abundance differ with geographic locations, food sources and weather factors. The major pests include whiteflies, mealy bugs, tea mosquito bugs, fruit flies and castor capsule borer . Besides, mites and birds also cause a least amount of loss. Weather factors play role in multiplication, growth, development and distribution of insects and influence on their seasonal abundance. Temperature, humidity and rainfall are most influential parameters of meteorological factors affecting insect population dynamics. Seasonal population dynamics of any insect pest provide knowledge on relationship between weather factors and insect abundance. ~~It indicates the farmers of a particular area or region about management program of the pest. So, the growers can take proper control measures to prevent loss due to insect attack.~~ Hence, The research was framed to survey and documentation of pests and its natural enemies in different guava varieties and to study the influence of weathers parameters on incidence of pests and natural enemies in guava.

Materials and methods

Studies on the Seasonal incidence of pests and natural enemies in different ~~guava varieties~~ and management of tea mosquito bug using bio – pesticides were carried out at Horticultural College and Research Institute, Periyakulam during 2022. ~~The survey was conducted in different guava varieties for the incidence of pests and natural enemies~~ The survey was conducted in different guava varieties for the incidence of pests and natural enemies at Central farm, Horticultural College and Research institute, Periyakulam during May to July, 2022. The varieties surveyed were Allahabad Safeda , Red fleshed, Lalit,

Lucknow 49 (Sardar) and Lucknow 46. Twenty plants were randomly selected from each Guava varieties were observed by visual counting the pests and natural enemy population and their infestation from 10 randomly selected fruits and twigs per at weekly intervals. The visual inspection of twigs, leaves and fruits was carried out and per cent damage on growing shoots, leaves and fruits were recorded. For fruit fly exit hole or deformation of fruit were taken as identification mark of infestation. The study area was kept free of insecticidal spray during the period of observation. The varietal screening studies data obtained was analyzed statistically using Randomized Block Design after suitable transformation. The weather parameters taken for the study were Maximum temperature, Minimum temperature, Rainfall, Wind velocity and Relative Humidity. These weather data were collected from the weather station of Horticultural College and Research Institute, Periyakulam and was correlated with the pests population ..

Results and Discussion

Insect pests and natural enemies on guava varieties

The incidence of pests and natural enemies in different guava varieties revealed that the tea mosquito bug, fruit fly, castor capsule borer, mealy bug, spiraling whitefly, aphids, scales, papaya Mealybug and predatory coccinellid beetles were found to be the major pests and predators in guava ecosystem of Horticultural College and Research Institute, Periyakulam (Table 1 and Fig 1.). Arifunnaharet *et al.*, (2016) stated that insects like spiraling whitefly, tea mosquito bug, scale insects, mealy bugs, aphids, thrips, coreid bug, fruit fly, fruit borer, stem borer, hairy caterpillar and leaf weevil are affecting guava in south India. Among the different varieties, L-46 was found to be less susceptible for tea mosquito bug (8.75%), fruit borer (0.75%), mealybugs (12.25% and 2.80 %) and aphids (2.40%) followed by red flash (1.55- 21.35%), L-49 (0.86- 13.50%) (Table 2), whereas, in allahabad safeda, the per cent damage by fruit fly (3.50%) and aphids (4.50%) was high compared to other guava varieties. The guava variety, Lalit was found be more susceptible for insect pests, which recorded tea mosquito bug (22.60%) , fruit borer (2.65%), mealy bug (25.60%), spiraling whitefly (23.45%), scales (5.75%) and papaya Mealybug (12.70%), but, Lalit was found be more susceptible for insect pests, which is in accordance with Ganga visalakshy. (2019) and Yasodha *et al.* (2023) who reported that pests causes more than 60 per cent yield loss in

guava. The coccinellids population was high in lalit (1.80/tree) followed by Allahabad safeda (1.70/tree) and L-49 (1.60/tree), whereas the spiders were found high in red flash guava (0.70/tree) but in lalit it was 0.15 per tree (Table 2).

Seasonal incidence of insect pests and their natural enemies in guava

The seasonal incidence of insect pests and their natural enemies in guava revealed that the tea mosquito bug, fruitfly damage, mealybugs, spiraling whitefly, aphids, scales, papaya mealybug and spiders was more during hot weather and was positively correlated with maximum and minimum temperature (0.0249 and 0.0909, 0.269 and 0.390, 0.063 and 0.048, 0.329 and 0.421, 0.140 and 0.007, 0.266 and 0.420 and 0.644 and 0.624, respectively) whereas they are negatively correlated with other weather parameters viz., rainfall, wind speed and relative humidity (Table 3). Shankar *et al.*, (2022) found that weather correlation and negative binomial hurdle model is a good fit for tea mosquito bug in neem. The fruit borer incidence was recorded high during rainy days (0.103) with high wind speed (0.344) and minimum temperature (0.004). The coccinellids beetles were more during cloudy and rainy period (0.765), high wind speed (0.554) and relative humidity (0.338). Baker *et al.*, (2012), Giddi *et al.*, (2021 and 2023) and Yasodha *et al.* (2023) reported that the abundance of insect pest of guava is related with weather factors and the lowest population density is during the winter season. Al-Fwaeer, *et al.*, (2013) and reported that the Guava fruit fly showed most abundance in months April to March and September to November, which coincide with the Guava fruiting season .

Table 1. Insect pests and natural enemies on guava varieties

| S.No. | Insect Pests on guava varieties | | |
|-------|---------------------------------|--|------------------------------|
| | Common Name | Scientific Name | Family and Order |
| 1. | Tea mosquito bug | <i>Helopeltisantonii</i> (Signoret) | Miridae: Hemiptera |
| 2. | Fruit fly | <i>Bactroceradiversus</i> | Tephritidae: Diptera |
| 3. | Fruit borer | <i>Conogethspunctiferalis</i> (Guenée) | Pyraustidae: Lepidoptera |
| 4. | Mealy bug | <i>Ferrisiavirgata</i> (Cockerell) | Pseudococcidae: Hemiptera |
| 5. | | <i>Maconellicoccushirsutus</i> (Green) | |
| 6. | Spiraling whitefly | <i>Aleurodicusdispersus</i> (Russell) | Aleyrodidae: Hemiptera |
| 7. | Aphids | <i>Aphis gossypii</i> (Glover) | Aphididae: Hemiptera |
| 8. | Guava scale | <i>Chloropulivinariapsidii</i> (Maskell) | Coccidae: Hemiptera |
| 9. | Papaya Mealybug | <i>Paracoccusmarginatus</i> (Williams) | Pseudococcidae: Hemiptera |
| 10. | Coccinellid beetles | <i>Coccinellaseptumpunctata</i> C. <i>repanda</i> | Coccinellidae: Coleoptera |




Figure 1. Insect pests and natural enemies on guava varieties

UNDER REVIEW

Table 2. Incidence of insect pests and natural enemies on guava varieties at Central farm, Horticultural College and Research Institute, Periyakulam during May- July, 2022

| Variety | Tea mosquit o bug (%) | Fruit fly (%) | Fruit borer (%) | Mealy bug (%) | Spirali ng whitefl y (%) | Aphids (%) | Guava scale (%) | Papaya Mealybu g (%) | Coccinellid (No./tree) | Spiders (No./tree) |
|----------------------|--------------------------------|------------------------------|-----------------------------|-------------------------------|-----------------------------------|-------------------------------|------------------------------|-------------------------------|-----------------------------|-----------------------------|
| Allahabad safeda | 12.90 (21.04) ^c | 3.50 (10.78) ^d | 1.60 (7.26) ^d | 17.00 (24.34) ^c | 10.65 (19.04) ^a | 4.50 (12.25) ^d | 1.20 (6.29) ^c | 10.75 (19.13) ^c | 1.70 (1.64) ^b | 0.56 (1.25) ^b |
| Red fleshed guava | 11.50 (19.82) ^b | 2.20 (8.52) ^c | 1.55 (7.15) ^c | 21.35 (27.51) ^d | 11.74 (19.94) ^b | 17.80 (24.95) ^e | 3.57 (10.89) ^d | 12.65 (20.83) ^d | 1.20 (1.48) ^d | 0.70 (1.30) ^a |
| L- 49 | 13.55 (21.59) ^d | 1.50 (7.03) ^b | 0.86 (5.31) ^b | 13.50 (21.55) ^b | 18.65 (25.54) ^c | 3.67 (11.04) ^c | 0.90 (5.44) ^a | 3.50 (10.78) ^b | 1.60 (1.61) ^c | 0.45 (1.21) ^c |
| Lalit | 22.60 (28.37) ^e | 1.35 (6.67) ^a | 2.65 (9.36) ^e | 25.60 (30.38) ^e | 23.45 (28.95) ^d | 2.65 (9.40) ^b | 5.75 (13.87) ^e | 12.70 (20.87) ^d | 1.80 (1.67) ^a | 0.15 (1.07) ^e |
| L- 46 | 8.75 (17.20) ^a | 2.20 (8.53) ^{cb} | 0.75 (4.96) ^a | 12.25 (20.48) ^a | 12.00 (20.26) ^b | 2.40 (8.91) ^a | 1.12 (6.07) ^b | 2.80 (9.63) ^a | 0.87 (1.37) ^e | 0.20 (1.20) ^d |
| CD (P=0.05) | 0.427 | 0.085 | 0.169 | 0.538 | 0.431 | 0.216 | 0.196 | 0.259 | 0.016 | 0.006 |

Figures in parentheses are square root transformed values * and arcsine transformed values **

Means followed by a common letter in a column are not significantly different

Values are mean of twenty replications

Table 3. Correlation co-efficient between incidence of insect pests and natural enemies on guava with weather parameters during May- July, 2022

| Weather Parameter | Weather Parameter correlation Coefficient | | | | | | | | | |
|-----------------------|---|-----------|-------------|-----------|--------------------|--------|-------------|-----------------|---------------------|---------|
| | Tea mosquito bug | Fruit fly | Fruit borer | Mealy bug | Spiraling whitefly | Aphids | Guava scale | Papaya Mealybug | Coccinellid beetles | Spiders |
| Max Temp (°C) | 0.479 | 0.025 | -0.009 | 0.269 | 0.063 | 0.329 | 0.140 | 0.266 | -0.278 | 0.644 |
| Min Temp (°C) | 0.577 | 0.091 | 0.004 | 0.390 | 0.048 | 0.421 | 0.007 | 0.420 | -0.298 | 0.624 |
| Rainfall (mm) | -0.351 | -0.392 | 0.103 | -0.634 | -0.687 | -0.295 | -0.169 | -0.165 | 0.765 | -0.651 |
| Wind speed | -0.103 | -0.163 | 0.344 | -0.086 | -0.475 | -0.059 | -0.057 | -0.469 | 0.554 | -0.237 |
| Relative humidity (%) | -0.502 | -0.163 | -0.025 | -0.456 | -0.167 | -0.382 | 0.003 | -0.457 | 0.338 | -0.653 |

Conclusion

The results obtained from this study, showed that major pests of guava viz. tea mosquito bug, fruit borer and mealybugs were maximum in Lalit followed by L-46, red flesh, Allahabad safeda and L-49. But the coccinellids population was high in Lalit followed by Allahabad safeda and L-49, whereas the spiders were found high in red flash guava. The tea mosquito bug, fruit fly damage, mealybugs, spiraling whitefly, aphids, scales, papaya mealybug and spiders was more during hot weather whereas fruit borer incidence was recorded high during rainy days with high wind speed and minimum temperature. The present research will help to develop forecasting models which help the farmers to implement management strategies in appropriate time to prevent yield loss and to avoid pesticide residue issues in export commodities, which are need of the hour.

Disclaimer (Artificial intelligence)

Author(s) hereby declare that **NO generative AI technologies** such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

Reference

1. Al-Fwaeer M, Abo-abied I, Abo-allosh A, Halybih M, Obeidat K, Atawee E and Al-hawamleh H. 2013. Study of pests attacking guava in Jordan. *Angewandten Biologie Forschung*, 1: 43-48
2. Amit Kumar Goswami, Anand Kumar Singh, Sanjay Kumar Singh, A. Nagaraja, Madhubala Thakre and Chavlesh Kumar. 2024. Newly Released Guava Hybrids: Pusa Aarushi and Pusa Pratikshan. *Indian Horticulture*, 69 (1):1cp
3. Arifunnahar M, Z Ferdous, MA Alim and MA Hossain. 2016. Incidence of spiralling whiteflies *Aleurodicus dispersus* Russell (hemiptera: aleyrodidae) and its natural enemies on guava orchards. *J. Bio-Sci.*,67-74.
4. Baker RFA, Mousa SF, Hamouda LS, Badawy RM and Attia SA. 2012. Scale insects infesting guava trees and control measure of *Pulvinari apsidii* (Hemiptera: Coccidae) by using the alternative insecticides. *Egypt Academic Journal of Biological Sciences*, 5: 89-106.

5. Ganga visalakshy.P.N , C. Swathi and Frenitalewis. 2019.Studies on the biological parameters of *Helopeltisantonii* Sign. (Hemiptera: Miridae) on *Psidium guajava* L. *Pest Management in Horticultural Ecosystems*,32-36. Yasodha , P., M. Dhayanandhan , N.Muthukrishnan
6. Giddi Thirumala Devi, Dr. N Emmanuel and Dr. V Sekhar. 2023. Seasonal incidence of sucking pest complex of guava CV. Taiwan white in Andhra Pradesh. *The Pharma Innovation Journal*, 12(3): 4154-4158
7. Giddi Thirumala Devi, N Emmanuel, CP Viji, Dr Salomi Suneetha and V Selham, 2021. Seasonal incidence of fruit borers in guava cv. Taiwan white. *Journal of Entomology and Zoology Studies*, 282-286.
8. Mitra S.K., M.R. Gurung, P.K. Pathak. 2008. Guava production and improvement in India: an overview. *In: International Workshop on Tropical and Subtropical Fruits.* DOI:10.17660/ActaHortic.2008.787.4
9. Shankar R.V., Ajaykumar, R., Prabhakaran, P., Kumaraperumal, R., Guna, M.,2022. Modeling of tea mosquito bug (*Helopeltis theivora*) incidence on neem tree: A zero in lated count data analysis. *Journal of Agrometeorology*, 24 (4) : 409-416.
10. DOI:10.17660/ActaHortic.2008.787.4, A.Krishnaveni. 2023. Seasonal Incidence Of Mango, Guava And Sapota Pests In Tiruvannamalai District Of Tamil Nadu, India. *Agriculture Association of Textile Chemical and Critical Reviews Journal*, 11(4) : 176-182 DOI : <https://doi.org/10.58321/AATCCReview.2023.11.04.176>
11. Mahata, Dibyendu, Mrityunjay Ghosh, B. Panja, K. Karmakar, and S. Banerjee. 2024. “Study on Major Disease-Insect Incidence of Scented Gobindabhog Rice in New Alluvial Zone of West Bengal”. *Journal of Experimental Agriculture International* 46 (5):34-39. <https://doi.org/10.9734/jeai/2024/v46i52353>.
12. Thakur, Kiran, Anand Kumar Panday, Dwarka, and Deepali. 2023. “Study the Seasonal Incidence of Major Sucking Insect-Pests of Sesame (*Sesamum Indicum*, L.) Crop”. *International Journal of Plant & Soil Science* 35 (19):1441-46. <https://doi.org/10.9734/ijpss/2023/v35i193688>.

13. Hooks CR, Valenzuela HR, Defrank J. Incidence of pests and arthropod natural enemies in zucchini grown with living mulches. *Agriculture, ecosystems & environment*. 1998 Jul 6;69(3):217-31.
14. Luck RF. Evaluation of natural enemies for biological control: a behavioral approach. *Trends in Ecology & Evolution*. 1990 Jun 1;5(6):196-9.

UNDER PEER REVIEW