

Original Research Article

Prevalence and Antimicrobial Susceptibility Patterns of *Salmonella*, *Shigella* and *Escherichia coli* among Children suffering from Diarrhoea, Unguja - Zanzibar, Tanzania.

Commented [S1]: Check on your spellings and grammar throughout your work.

Commented [S2]: This title is similar to already published work on <https://pubmed.ncbi.nlm.nih.gov/37193235/>, kindly reframe it please.

ABSTRACT

Aims

This study aimed at providing evidence of the prevalence and antimicrobial susceptibility patterns towards *Salmonella*, *Shigella* and *Escherichia coli* among under-five children in Unguja – Zanzibar, Tanzania.

Study design: Cross sectional design was used to collect samples from stool of children suffering from diarrhoea.

Place and Duration of Study: The study was carried out in Zanzibar west urban region between October, 2019 to February, 2020.

Methodology

A cross-sectional study was conducted from October 2019 to February 2020. Random samples were collected to investigate the prevalence of *Salmonella*, *Shigella* and *Escherichia coli*. The samples were cultured using Hektoen Enteric (HE) and *Salmonella-Shigella* agar. Antibiotic susceptibility testing was done by Kirby–Bauer disc diffusion method.

Results

A total of 159 stool samples were collected in the study; *Salmonella* was identified 12/159 times (7.5%) of the total samples. *Shigella* and *E. coli* were identified in 7/159 samples (4.4%) and 6/159 (3.7%), respectively. Children between 49 and 60 months showed low prevalence, while a high peak prevalence was reported for children between 7–12 months. All *Salmonella*, *Shigella* and *Escherichia coli* species identified were sufficiently susceptible to chloramphenicol and ceftriaxone, with a varying pattern to azithromycin, ciprofloxacin, ampicillin, nalidixic acid and trimethoprim-sulfamethoxazole.

Conclusion

We found *Salmonella*, *Shigella* and *Escherichia coli* isolates in stools of children ≤ 5years from Unguja, Zanzibar, but all the isolates were susceptible to chloramphenicol and ceftriaxone but partially resistant to other tested antibiotics. Identifying resistant bacteria in this age group should be a concern for the public health authorities and trigger research into finding the cause.

Keywords: *Salmonella*, *Shigella*, *Escherichia coli*, Diarrhoea, Prevalence, Antimicrobial Susceptibility and Children under-five years.

Commented [S3]: Give summary as follows

Introduction
Objective
Methodology
Result
Conclusion

Commented [S4]: This one is misplaced, kindly add it under methodology

Commented [S5]: This one is misplaced, kindly add it under methodology

Commented [S6]: Kindly indicate all the tools used to achieve your findings

Commented [S7]: What new are you adding here kindly reframe this part to sound well.
EXAMPLE.

Conclusion: This study showed *E. coli* as the predominant isolate causing diarrheal illness in under five children. The emergence of multidrug resistant phenotypes warrants the continuous monitoring of susceptibility trend of NTS (Non Typhoidal *Salmonella*) in TZ

1. INTRODUCTION

Diarrhoea is a frequent discharge of a watery stool accompanied by abdominal cramps, nausea and vomiting, sometimes with fever and chills. The World Health Organization (WHO) estimates that about 1.7 billion cases of childhood diarrhoeal diseases are responsible for killing 525,000 children yearly (1,2),(3). [1-3] Several countries have diarrhoea outbreaks from *Salmonella*, *Shigella* and *Escherichia coli* (*E. coli*) associated with consumption of contaminated food, water or direct faecal contamination through

Commented [S8]: Adopt this style but you can find out with the journal site.

fingers, the major source of transmission of diarrhoeagenic disease (4,5). It has been reported that most deaths and hospitalization due to diarrhoea occur in the developing countries, particularly in Africa (1,6). *Salmonella*, *Shigella* and *E. coli* are amongst the species that cause diarrhoea in Africa (4,7,8).

The one health approach is the most recommended method to tackle antimicrobial resistance in human health, animal health, food and environment sectors. The common antibiotics for treating diarrhoea caused by *Salmonella*, *Shigella* and *E. coli* are ampicillin, chloramphenicol, and trimethoprim-sulfamethoxazole but there are evidence of resistance to these and other antimicrobials commonly used to treat diarrhoea in developing countries (5,9–11). Unfortunately, limited laboratory techniques to test antimicrobial susceptibility has resulted in a minimum understanding of the resistance burden and reduced therapeutic efficacy (12–15). Thus, there is a need to understand the country-level epidemiology of these diarrhoeagenic bacteria and the efficacy of the available antibiotics to treat the diarrhoea and associated illnesses.

Consumption of contaminated food and water is a common but inadvertent practice in Zanzibar linked to the cultural food system and the environment. For more than 40 years, diarrhoeal diseases have been the common public health problem presenting with outbreaks in Zanzibar (16). It is hypothesized that *Salmonella*, *Shigella* and *E.coli* are responsible for the outbreaks (17). However, there are more microbial organisms causing diarrhoea apart from *Salmonella*, *Shigella* and *E. coli*. This study aimed to investigate the prevalence and antimicrobial susceptibility patterns of *Salmonella*, *Shigella* and *Escherichia coli* among diarrhoea suffering Children in Unguja - Zanzibar. Tanzania.

2. MATERIAL AND METHODS

Study design and site

This was a rapid cross-sectional study, covering public health facilities in the three districts of west urban region, namely urban district, west 'A' district and west 'B' district, from October, 2019 to February, 2020. A random sampling was used to select nine public health facilities in each district. The twenty-seven selected health facilities were Mnazi Mmoja, Chumbuni, Sebleni, Rahaleo, Kwamtipura, Kidongo Chekundu, Kidutani, Shaurimoyo, Mpendae, Fuoni, Kombeni, Magogoni, Kiembe Samaki, Fuoni Kibondeni, Shakani, Bwefum, Chukwani, Kisauni, Mbweni Matrekta, Mtofaani, Selem, Bubwisudi, Chuini, Kizimbani, Kianga, Beit-el-Ras and Kibweni. These public health facilities are government properties, treatment of children is free and hence provides health care to most children.

Sample size determination

Population proportion formula was employed using desired characteristics of 12% (18) of diarrhoea cases, as calculated below.

Fisher's formula: $n = Z^2pq/r^2$ (19).

*

Commented [S9]: How i wish to see a cler background of the study, drug resistance parten on the study, mechanism employed by each bacteria towards the drugs etc

Commented [S10]: How did you overcome the study biasness kindly indicate

Where: n = Desired sample size. p = Proportion of the population with a desired characteristic of 12% (18). $q = 1-p$; z = standard deviation desired degree of accuracy. Where z is 1.96, if the degree of confidence is 95%; r = Degree of error, which was 5%

Therefore: n was found to be 159.

The characteristics of patients whom stool samples were collected from under-five children presenting with diarrhoea frequency > three times per 24 hours, either bloody or watery diarrhoea included suspected cases of cholera. The age and play environment of patients have been shown relation to what has been studied. The samples were collected rainy and dry season with unsuspected and suspected cholera epidemics.

Commented [S11]: Not clear

Sample collection, storage and transportation

One hundred and fifty-nine stool samples were collected from 27 health facilities using sterile plastic containers. All samples collection was performed from a period of October, 2019 – February, 2020, immediately after samples collection, packed in a cool box at 4°C and sent to the microbiology department of the pathology laboratory at Mnazi Mmoja hospital Zanzibar for further analysis.

Bacterial culture, isolation, and identification

In the laboratory, approximately 1 g of stool specimen was placed overnight into 10 ml Selenite-F enrichment broth (Oxoid UK) in a sterile test tube. A loopful of the suspension of the specimen was streaked onto two different media, namely, Hektoen Enteric (HE) Agar and *Salmonella Shigella* (SS) Agar, both from Oxoid, UK. These plates were incubated under 37 °C for 48 hours. A colorless colony with or without a black centre on SS Agar media and a blue-green colony with or without a black centre on HE Agar were isolated as *Salmonella*-like isolates. A colorless colony on SS and a green, moist and raised colony on HE Agar were isolated as *Shigella*-like isolates (20). Colonies exhibiting characteristic reactions of *Salmonella*, *Shigella* and *E. coli*-like were further characterized by the pattern of biochemical reactions after inoculation to Triple sugar iron Agar, lysine iron Agar, Simon's citrate Agar and motility test, Indole and Urease production (MIU) test for final identification using the standard procedures (20).

Commented [S12]: Refer to CLSI 2024 guideline on stool culture
Refer to this
<https://www.microbiologyresearch.org/content/journal/acmi/10.1099/acmi.0.000614.v2>

Antimicrobial Susceptibility Testing

Kirby-Bauer's disc diffusion method was used for antibiotic sensitivity tests. One of the easiest and quickest methods that can be used to test the antibiotic sensitivity of a bacterial isolate. Antimicrobial susceptibility testing with discs is a simple, rapid method and provides a reproducible means of testing bacterial sensitivity to various antibiotics and chemotherapeutic agents. Pure overnight cultures of *Salmonella Shigella* and *E. coli* isolates were mixed with sterile saline and after matching with 0.5 McFarland standards were inoculated in Mueller-Hinton Agar. Antibiotic discs were placed on to the Agar. *Shigella* and *E. coli* isolates were tested against ceftriaxone (CTX, 30 µg), ciprofloxacin (CIP, 5 µg), chloramphenicol (C, 30 µg), ampicillin (AMP, 10 µg), trimethoprim-sulfamethoxazole (SXT, 25 µg) or cotrimoxazole (CO, 25 µg), nalidixic acid (NA, 30 µg), and azithromycin (AZM, 15 µg). Inhibition diameter

*

zone readings recorded according to Clinical and Laboratory Standards Institute (21) and results were reported as sensitive (S), intermediate (I) and resistance (R).

Commented [S13]: <https://www.microbiologyresearch.org/content/journal/acmi/10.1099/acmi.0.000614.v2>

Data handling and statistical analysis

Data were initially compiled in an MS excel spreadsheet, and statistical analyses were performed using Statistical Package for Social Sciences (SPSS) software (16.0 version). Descriptive statistics were calculated and summarized in frequency and proportions. Prevalence and antimicrobial susceptibility patterns of *Salmonella*, *Shigella* and *E. coli* were determined focused on variables potentially magnitude, typical of *Salmonella*, *Shigella* and *E. coli* isolates and their proportions.

3. RESULTS AND DISCUSION

Results

Socio-demographic characteristics

A total of 159 children under 5-year of age with diarrhoea were included in the study. Of the 159 participants, 73 (45.9%) were females, and 86 (54.08%) were males. The specific age categorization was as follow: 17 (10.69%) of them were between 0 to 6 months, 56 (35.2%) were between 7 to 12 months, 44 (27.67%) were between 13 to 24 months, 24 (15.09%) were between 25 to 36 months, 14 (8.8%) were between 37 to 48 months, and 4 (2.5%) were between 49 to 60 months (Figure 1)




Figure 1: Distribution of Participants by Age and Sex

Prevalence of *Salmonella*, *Shigella* and *E. coli*

A total of *Salmonella* 12/159 (7.5%), 7/159 *Shigella* (4.4%) and 6/159 *E. coli* (3.7%) were identified among under-five children with diarrhoea in Unguja - Zanzibar. Tanzania. In relation to the proportion of isolation, *Salmonella* 4/159 (2.5%) were the highest among the age group of between 7 to 12 months, but

*

it was not identified in children above 37 months (Table 1). The *Shigella* 1/159 (0.66%) was identified among 0 to 6 months, 13 to 24 months, 25 to 36 months children, and 37 to 48 months but it was not found in 7 to 12 months and above 49 months children (Table 1). The *E. coli* (1(0.66%)) was identified among 0-6 months, 13-24 months, and 25-36 months children, but not among 7 - 12 months and above 37 months children. Among the 25 culture-positive children, 13 (52%) were female (Table 1).

Table 1: The Distribution of *Salmonella*, *Shigella* and *E. coli* among Children Suffering from Diarrhoea by Age and Sex in Unguja – Zanzibar.

Variables	Patients			
	frequency (%)	<i>Salmonella</i> (%)	<i>Shigella</i> (%)	<i>Escherichia coli</i> (%)
Sex				
Male	86 (54.08%)	6 (3.7)	4 (2.5)	2 (1.3)
Female	73 (45.91%)	6 (3.7)	3 (1.8)	4 (2.5)
Age				
0 – 6months	17 (10.69%)	1 (0.6)	1 (0.6)	1 (0.6)
7 – 12months	56 (35.22%)	4 (2.5)	0(0)	0(0)
13 – 24months	44 (27.67%)	1 (0.6)	1(0.6)	1 (0.6)
25 – 36months	24 (15.09%)	1 (0.6)	1 (0.6)	1 (0.6)
37 – 48months	14(8.8)	0(0)	1(0.6)	0(0)
49 – 60 months	4(2.51)	0(0)	0(0)	0(0)

Antibiotic Susceptibility Test from patients' stool samples

The *Salmonella* isolates displayed different susceptibility rates to the evaluated antibiotics (Table 2). *Salmonella* isolates were highly susceptible to ceftriaxone 12 (100%), chloramphenicol 12 (100%) and trimethoprim-sulfamethoxazole 12 (100%), azithromycin 10 (83.3%), ciprofloxacin 8 (66.6%) and ampicillin 8 (66.6%) while no susceptible to nalidixic acid. They exhibited low resistance to nalidixic acid 5 (41.6%), azithromycin 2 (16.6%) and ampicillin 2 (16.6%). The results indicate no resistance to Chloramphenicol, Ciprofloxacin, Ceftriaxone and Trimethoprim-sulfamethoxazole.

Table 2. Antibiotic Susceptible Pattern of *Salmonella*, *Shigella* and *E. coli* from Patient

Antimicrobials	<i>Salmonella</i>			<i>Shigella</i>			<i>E coli</i>		
	S	I	R	S	I	R	S	I	R
Ampicillin	8(66.6)	2(16.6)	2(16.6)	2(28.5)	5(71.4)	0(0.0)	2(33.3)	1(33.3)	2(33.3)
Ceftriaxone	12(100)	0(0.0)	0(0.0)	7(100)	2(28.5)	0(0.0)	6(100)	0(0.0)	0(0.0)
Chloramphenicol	12(100)	0(0.0)	0(0.0)	7(100)	0(0.0)	0(0.0)	6(100)	0(0.0)	0(0.0)
Nalidixic acid	0(0.0)	7(58.3)	5(41.6)	2(28.5)	3(42.8)	2(28.5)	2(33.3)	2(33.3)	2(33.3)
Ciprofloxacin	8(66.6)	4(33.3)	0(0.0)	4(57.1)	0(0.0)	0(0.0)	4(66.6)	2(33.3)	0(0.0)
Azithromycin	10(83.3)	0(0.0)	2(16.6)	4(57.1)	0(0.0)	3(42.8)	4(66.6)	1(16.6)	1(16.6)

*

Trimethoprim-sulfamethoxazole	12(100)	0(0.0)	0(0.0)	2(28.5)	0(0.0)	5(71.4)	2(33.3)	0(0.0)	4(66.6)
-------------------------------	---------	--------	--------	---------	--------	---------	---------	--------	---------

Abbreviations: **S** - Sensitive, **R** - Resistant and **I** - Intermediate

The *Shigella* isolates were highly susceptible to chloramphenicol 7 (100%) and ceftriaxone 7 (100%) while lowly susceptible to ampicillin 2 (28.5%), nalidixic acid 2 (28.5%) and trimethoprim-sulfamethoxazole 2(28.5%). However, these isolates showed different resistance against trimethoprim-sulfamethoxazole 5 (71.4%), azithromycin 3 (42.8%) and nalidixic acid 2 (28.5%). *Shigella* was found to show no resistance to Chloramphenicol, Ceftriaxone, Ciprofloxacin and Ampicillin, as shown in Table 2

The *E. coli* isolates were highly susceptible to chloramphenicol 6 (100%) and ceftriaxone 6 (100%) and lowly susceptible to ampicillin 2 (33.3%), nalidixic acid 2 (33.3%) and trimethoprim-sulfamethoxazole 2 (33.3%). *E. coli* isolates showed different resistance against trimethoprim-sulfamethoxazole 4 (66.6%), azithromycin 1(16.6%), ampicillin 2 (33.3%) and nalidixic acid 2 (33.3%). These results indicate that *E. coli* was highly resistant to Trimethoprim-sulfamethoxazole 4 (66.6%), as shown in Table 2

The results of multiple drug resistant (MDR) patterns of *Salmonella*, *Shigella* and *E. coli* isolates exhibited multidrug resistance to five antibiotics azithromycin, nalidixic acid, ampicillin, trimethoprim-sulfamethoxazole and ceftriaxone. *Salmonella* isolates exhibited multidrug resistance to three antibiotics, namely azithromycin, nalidixic acid and ampicillin, while *Shigella* isolates exhibited to four antibiotics; trimethoprim-sulfamethoxazole, azithromycin, nalidixic acid and ceftriaxone. On the other hand, *Escherichia coli* isolates exhibited multidrug resistance to three antibiotics: ampicillin, trimethoprim-sulfamethoxazole and nalidixic acid. These results show that *Salmonella* and *Escherichia coli* were less resistant to drugs than *Shigella*, as shown in Tables 2.

Discussion

This study has provided evidence of prevalence and antimicrobial susceptibility patterns towards *Salmonella*, *Shigella* and *Escherichia coli* among under-five children in Unguja – Zanzibar, Tanzania.

Our results have shown a low prevalence to children between 49 and 60 months and high peak for children between 7–12 months. The low prevalence of the target pathogens, especially *Shigella*, in the current study could be attributed to improved awareness of the mothers and caretakers of under-five children and community about personal and environmental hygiene from continuous interventions being made by different stakeholders, including Health extension workers through an educational program like Community Based Education. In our study, children within the age range of 1- 3 years were more susceptible to diarrhoea caused by *Salmonella* 12 (7.5%) than *Shigella* 7 (4.4%) and *Escherichia coli* 6(3.7%) These results indicated that there is no statistically significant relationship between age of patients and identification of *Salmonella*, *Shigella* and *Escherichia coli* (chi-square with four degree of freedom = 2, p = 0.064). These findings were consistent with several previous studies on the matter (5,8,22). Children in this age group normally like to take contaminated soils, food and water into their

*

mouths, eventually leading them to get diseases from the environment easily caused by microbes, including pathogenic *Salmonella*, *Shigella* and *Escherichia coli*.

The results showed that patients' stool had more *Salmonella* isolates, followed by *Shigella* and low isolates of *Escherichia coli*. The *Salmonella* rate of 12 (7.5%) found in this study is higher than the rates found in other studies, including those from retail meat and meat products in China at 3.6% (23), Ethiopia at 6.9% (22), Ethiopia at 1% (5), India at 1% (24) and Qatar at 3.23% (25). Meanwhile, the *Salmonella* rate is still lower compared to the rate found in China at 19.7% (23), China in pork products at 37.3% (23), and China in beef at 16% (23).

On the other side, the isolation rate of *Shigella* 7 (4.4%) was found consistent with the rate in Ethiopia (22), which reported 4.3% of *Shigella* from stools. The high isolates of *Shigella* reported in Tanzania were 16.1% (26) and Ethiopia was 8.3% (5), while low isolates were reported by Jimma 1.1% (27). The isolation rate of *Escherichia coli* 6 (3.7%) was found to be lower than other studies in Tanzania, i.e., 21.6% (26) and 14% in Zanzibar (8).

Looking into treatment options, we found *Salmonella* isolates being highly susceptible to ceftriaxone 100%, chloramphenicol 100%, and trimethoprim-sulfamethoxazole 100%. However, our study reports resistance to nalidixic acid 41.6%, ampicillin 16.6% and azithromycin 16.6%. Similar studies in Ethiopia reported *Salmonella* susceptible to ceftriaxone 100% (22), while ampicillin had high resistance of 100% (5) in Ethiopia. In addition, several studies reported *Salmonella* resistance to ampicillin at 46% in Zanzibar (8), 45.4% in China (23) and 34% in India (24). The resistance patterns to nalidixic acid 43% were reported in Kolkata, India (24). Other resistance patterns in our study were observed in *Salmonella* to nalidixic acid at 35.8% in China (23). The *Salmonella* isolates in the previous study reported resistance patterns to azithromycin 25% (24), which were inconsistent with our study.

The *Shigella* isolates were found to be highly susceptible to chloramphenicol 100% and ceftriaxone 100%. Nevertheless, our findings reported resistance to trimethoprim-sulfamethoxazole 71.4%, azithromycin 42.8% and nalidixic acid 28.5%. Likewise, a study in Zanzibar reported *Shigella* susceptible to chloramphenicol at 46% (8). Other studies reported *Shigella* resistance of trimethoprim-sulfamethoxazole in Bosnia and Herzegovina 70-86% (28) and 68% in Zanzibar (8).

The *Escherichia coli* isolates are highly susceptible to chloramphenicol 100% and ceftriaxone 100%. However, our study reports resistance to trimethoprim-sulfamethoxazole 66.6%, ampicillin 33.3% and nalidixic acid 33.3%. A similar study in Zanzibar reported *Escherichia coli* susceptible to chloramphenicol at 77.3%, which was inconsistent with our study (8). Other studies reported *Escherichia coli* resistance to trimethoprim-sulfamethoxazole at 68% (8) in Zanzibar and 40-86% in Bosnia and Herzegovina (28), which is in agreement with our study. These results indicate a reduced efficacy in treating *Salmonella*, *Shigella* and *Escherichia coli* among under-five children with diarrhoea. The species isolates in our study

*

exhibited multidrug-resistant patterns to at least five antibiotics, namely Trimethoprim-sulfamethoxazole, azithromycin, nalidixic acid, ceftriaxone and ampicillin.

We reported the prevalence and the resistance patterns of *Salmonella*, *Shigella* and *Escherichia coli* in Zanzibar to guide the first-line and second-line drug choice in the treatment of diarrhoea in the subsequent reviews of the Zanzibar Standard Treatment Guideline (ZSTG) for Diarrhoeal diseases, as needed by Ministry of Health, 2016. Moreover, those medicines are easily available in pharmacies and accessible to anyone. These results are important to the Ministry of Health of Zanzibar in relation to the treatment of diarrhoeal diseases. The Ministry needs to initiate a long-term surveillance program to monitor and identify the changes in the rate of antimicrobial patterns of these bacteria of public health concern. Antibiotics remain the most important therapy for successfully managing diarrhoea infections; however, these inexpensive and widely available antimicrobials can no longer be used empirically. There is a need for appropriate control measures for antimicrobial resistance pathogens in Zanzibar.

4. CONCLUSION

We found *Salmonella*, *Shigella* and *Escherichia coli* isolates in stools of children ≤ 5 years from Unguja, Zanzibar, but all the isolates were susceptible to chloramphenicol and ceftriaxone but partially resistant to other tested antibiotics. Identifying resistant bacteria in this age group should be a concern for the public health authorities and trigger research into finding the cause.

8. CONSENT

All authors declare that written informed consent was obtained among Children suffering from Diarrhoea,

9. ETHICAL APPROVAL

Ethical approval was granted from the Zanzibar medical research ethics committee (Ref. No. ZAHREC/02/DEC/2018/6). Permission to conduct the study was sought from the respective health centres' authorities. The written informed consent was obtained from mothers or caretakers of children under five years before collecting of information. The patients result of any investigation remained confidential, while identified organisms were referred to attending physicians for treatment. The samples safe discarded and stored in a deep freezer for further investigation.

REFERENCES

1. FAO and WHO. Risk assessment tools for *Vibrio parahaemolyticus* and *Vibrio vulnificus* associated with seafood. Microbiological Risk Assessment Series (FAO/WHO) eng no. 20. Rome. 2020.

*

Commented [S14]: redo this and compare your finding with other studies

Commented [S15]: can you justify this with a reference

Commented [S16]: have noticed that the citations in your manuscript are not properly formatted. Citations must be listed sequentially from the smallest to the largest, as they appear in the text with no citation missing.

2. IVAC. Pneumonia Progress Report 2020. John Hopkins Bloomberg School of Public Health. 2020;1–22.
3. Troeger C, Forouzanfar M, Rao PC, Khalil I, Brown A, Reiner RC, et al. Estimates of global, regional, and national morbidity, mortality, and aetiologies of diarrhoeal diseases: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet Infect Dis*. 2017 Sep;17(9):909–48.
4. Ugboko HU, Nwinyi OC, Oranusi SU, Oyewale JO. Childhood diarrhoeal diseases in developing countries. *Heliyon*. 2020;6(4):e03690.
5. Abebe W, Earsido A, Taye S, Assefa M, Eyasu A, Godebo G. Prevalence and antibiotic susceptibility patterns of *Shigella* and *Salmonella* among children aged below five years with Diarrhoea attending Nigist Eleni Mohammed memorial hospital, South Ethiopia. *BMC Pediatr*. 2018;18(1):10–5.
6. Ali A, Parisi A, Conversano MC, Iannacci A, D'Emilio F, Mercurio V, et al. Food-borne bacteria associated with seafoods: A brief review. *J Food Qual Hazards Control*. 2020;7(1):4–10.
7. Rabia A, Wambura P, Misinzo G, Kimera S, Mdegela R, Mzula A, et al. Molecular Epidemiology of *Vibrio cholerae* Recovered from Sewage Drains, Captured Fish and Humans in 2015/16 Cholera Outbreak in Zanzibar, Tanzania. *J Adv Microbiol*. 2017;5(3):1–11.
8. Omari MH. Prevalence of Enteric Bacteria Associated With Diarrhoea in Children Less Than Five Years of Age ; and. 2015;
9. Njuguna CK, Mgamb E, Njeru I, Langat D, Makokha A, Ongore D, et al. Treatment of shigella infections: Why sulfamethoxazoletrimethoprim, tetracyclines and ampicillin should no longer be used. *East Afr Med J*. 2016;93(8):318–26.
10. Okorie-Kanu OJ, Ezenduka EV, Okorie-Kanu CO, Ugwu LC, Nnamani UJ. Occurrence and antimicrobial resistance of pathogenic *escherichia coli* and *salmonella* spp. in retail raw table eggs sold for human consumption in enugu state, nigeria. *Vet World*. 2016;9(11):1312–9.
11. Ali Nor BS, Menza NC, Musyoki AM. Multidrug-Resistant Shigellosis among Children Aged below Five Years with Diarrhea at Banadir Hospital in Mogadishu, Somalia. *Canadian Journal of Infectious Diseases and Medical Microbiology*. 2021;2021.
12. Iskandar K, Molinier L, Hallit S, Sartelli M, Hardcastle TC, Haque M, et al. Surveillance of antimicrobial resistance in low- and middle-income countries: a scattered picture. *Antimicrob Resist Infect Control*. 2021;10(1):1–19.
13. Ayukekbong JA, Ntemgwa M, Atabe AN. The threat of antimicrobial resistance in developing countries: Causes and control strategies. *Antimicrob Resist Infect Control*. 2017;6(1):1–8.
14. Marami D, Hailu K, Tolera M. Prevalence and antimicrobial susceptibility pattern of *Salmonella* and *Shigella* species among asymptomatic food handlers working in Haramaya University cafeterias, Eastern Ethiopia. *BMC Res Notes*. 2018;11(1):7–12.
15. Gebrekidan A, Dejene TA, Kahsay G, Wasihun AG. Prevalence and antimicrobial susceptibility patterns of *Shigella* among acute diarrheal outpatients in Mekelle hospital, Northern Ethiopia. *BMC Res Notes*. 2015;8(1):1–7.

*

16. Ministry of Health Zanzibar. Zanzibar Comprehensive Cholera Elimination Plan (ZACCEP): 2018 - 2027. 2018;2018–27.
17. Ministry Of Health Zanzibar. Zanzibar Guidelines for Prevention and Control of Cholera. 2016;1–71.
18. Edwin P, Azage M. Geographical Variations and Factors Associated with Childhood Diarrhea in Tanzania: A National Population Based Survey 2015-16. *Ethiop J Health Sci.* 2019;29(4):513–24.
19. Singh, Ajay S, Masuku MB. Sampling Techniques & Determining Sample Size in Applied Statistics Research: an Overview. *International Journal of Economics, Commerce and Management.* 2014;II(11):1–22.
20. Vandepitte J, Verhaegen, Engbaek K, Rohner P, Piot P, Heuck CC. L a B O R A T O R Y Procedures. *Nutrients.* 2003;3rd ed(92 4 154545 3):188.
21. CLSI. CLSI - Catalogue 2019 -The Highest Standards for Global Health Care. 2019;
22. Assefa A, Girma M. Prevalence and antimicrobial susceptibility patterns of Salmonella and Shigella isolates among children aged below five years with diarrhea attending Robe General Hospital and Goba Referral Hospital, South East Ethiopia. *Trop Dis Travel Med Vaccines.* 2019;5(1):1–11.
23. Yang X, Wu Q, Zhang J, Huang J, Chen L, Wu S, et al. Prevalence, bacterial load, and antimicrobial resistance of salmonella serovars isolated from retail meat and meat products in China. *Front Microbiol.* 2019;10(SEP):1–9.
24. Jain P, Chowdhury G, Samajpati S, Basak S, Ganai A, Samanta S, et al. Characterization of non-typhoidal Salmonella isolates from children with acute gastroenteritis, Kolkata, India, during 2000–2016. *Brazilian Journal of Microbiology.* 2020;51(2):613–27.
25. Al-Dahshan A, Elyamani R, Naja S, Chehab M, Nour M, Elmagboul E, et al. Epidemiological characteristics of a salmonella outbreak among infants in Qatar, 2017. *Qatar Med J.* 2019;2019(3):1–8.
26. Ngoso BE, Namkinga LA, Nkwengulila G. Molecular Characterization of Diarrheagenic Bacteria Isolated from Stool of Under-five Children in Dar Es Salaam. Tanzania. *Journal of Biology and Life Science.* 2015;7(1):71.
27. Lamboro T, Ketema T, Bacha K. Prevalence and Antimicrobial Resistance in Salmonella and Shigella Species Isolated from Outpatients, Jimma University Specialized Hospital, Southwest Ethiopia. *Canadian Journal of Infectious Diseases and Medical Microbiology.* 2016;2016.
28. Vranic S, Uzunovic A. Antimicrobial Resistance of Escherichia Coli Strains Isolated from Urine at Outpatient Population: a Single Laboratory Experience. *Materia Socio Medica.* 2016;28(2):121.

Commented [S17]: what is this for