

Evaluation of green gram (*Vigna radiata* L. var PDM 139) cultivated and harvested soils supplemented with Nano Urea, Rhizobium and Farm yard manure

ABSTRACT

A study was conducted to investigate the “Evaluation of varying levels of Nano urea, FYM, and rhizobium on soil properties and yield characteristics of green gram (*Vigna radiata*. L) var. PDM 139” at the research farm of soil science and agricultural chemistry. The experiment followed a randomized block design (RBD) with three replications. Treatment T₉ (NU at 100% + RZ at 100% + FYM at 100%) showed a slight decrease in pH (6.96 P^H), bulk density (1.27 mg m⁻³), particle density (2.46 mg m⁻³), there was a significant increase in pore space (48.90%), water holding capacity (46.51%), EC (0.20 ds m⁻³), organic carbon (0.58%), available nitrogen (320 kg ha⁻¹), phosphorus (21.4 kg ha⁻¹), potassium (195.12 kg ha⁻¹), and plant growth. Among all the treatment joint use of T₉ (NU at 100% + RZ at 100% + FYM at 100%) showed the most significant impact on green gram growth through healthy soil.

Keywords: Greengram; Soil; Nano urea; Rhizobium; FYM.

1. INTRODUCTION

Soil serves as the primary medium for plant growth and is fundamental to crop production. Various soil properties affect plant growth, including texture (whether coarse or fine), aggregate size, porosity, permeability (aeration), water holding capacity, pH level, bulk density, and particle density. The rate at which water moves into the soil (infiltration) is influenced by soil texture, physical structure (soil structure and tilth), and the amount of vegetation covering the soil surface. Organic matter plays a role in enhancing water retention in all types of soils and can also improve infiltration rates in fine-textured soils. Bulk density indicates the soil's ability to support plant structure, facilitate water and solute movement, and maintain soil aeration (Singh *et al.*, 2008). Soil pH significantly impacts the solubility of essential nutrients required for optimal plant growth and development [20,21,22]. Monitoring soil pH is valuable as it predicts various chemical processes in the soil. It aids in decisions related to plant suitability for specific locations, potential pH adjustments, and provides a rough estimate of nutrient availability to plants in the soil. Plants primarily obtain carbon, oxygen, and hydrogen from air and water, while other essential nutrients, termed plant nutrients, are sourced from soil or supplemented through fertilizers and absorbed

predominantly through roots (**Singh et al., 2016**). Green gram, also known as mung bean, ranks as the third most significant pulse crop among thirteen food legumes cultivated in India. The functional properties of proteins are influenced by various physical and chemical factors such as soil pH, ionic strength, temperature, and pressure, contributing to their variability and potential in food production (**Meena et al., 2016**). Proteins are typically expensive to produce industrially. With a growing global population and improved standards of living, there is increasing pressure on protein availability for food consumption. This underscores the importance of maximizing the use of green gram, particularly its nutritious germinated form (**Suryavanshi et al., 2018**).

Green gram enhances soil physical properties. The productivity of moong bean is hindered by its cultivation on marginal and submarginal lands with inadequate fertilization and suboptimal management practices. Poor yields of moong bean in farmer fields stem from insufficient awareness among farmers regarding optimal sowing dates, effective weed control, balanced fertilizer application, pest management strategies, and proper planting methods. Delayed planting of moong bean results in decreased pod or plant numbers, reduced grain weight, and ultimately lower grain yield. Sowing time is identified as the foremost agronomic factor crucial for realizing the yield potential of improved moong bean varieties. Optimal sowing time ensures synchronization between the vegetative and reproductive stages of the crop, thereby playing a pivotal role in achieving high seed yields (**Rathore et al., 2019**).

Liquid Nano-urea, when applied as a foliar spray, penetrates through stomata and other openings on leaves and is readily absorbed by plant cells. Nano-urea in liquid form efficiently moves through the plant's phloem from source to sink as required. Unused nitrogen is stored in plant vacuoles and released gradually to support proper growth and development. Unlike subsidized urea, liquid Nano-urea is offered to farmers at a 10% lower cost per unit and is easier to transport, with one 500-ml bottle equating to one bag of regular urea. It is safe for humans, animals, birds, rhizosphere organisms, and the environment at recommended application levels. Compared to conventional urea, Nano-urea's uptake efficiency exceeds 80%, reducing the quantity needed to meet plant nitrogen requirements. Foliar application of Nano-urea at critical crop growth stages effectively meets nitrogen needs, enhancing fertilizer efficiency, crop productivity, and quality. Nano-urea can potentially reduce reliance on conventional urea by up to 50% (**Subramani et al., 2023**). Rhizobium, a genus of gram-negative soil bacteria, participates in nitrogen fixation. It forms an endosymbiotic relationship with the roots of legumes and various flowering plants.

Farmyard manure (FYM) contributes significantly to enhancing soil fertility and productivity by positively impacting soil physical, chemical, and biological properties, as well as ensuring balanced plant nutrition (Sharma and Verma, 2011). Fertilizers are crucial for sustaining and enhancing soil fertility by providing plants with readily available nutrients. Therefore, it is important to assess the impact of combining farmyard manure (FYM) with chemical fertilizers on the growth and yield of post-rainy season green gram (Kumar *et al.*, 2017). Applying nano urea through foliar methods alongside organic and inorganic fertilizers enhanced the growth and yield of green gram (Tejaswini *et al.*, 2022). The current study aims to assess soil health and fertility through the application of nano urea (NU), rhizobium (RZ), and farmyard manure (FYM), examining their impact on crop growth and yield.

2. Materials and Methods

2.1 Experimental Site

The experiment took place at the Research Farm of Soil Science and Agricultural Chemistry, located approximately 6 km from Prayagraj city. The farm is situated at latitudes 23°52' N to 31°28' N and longitudes 77°3' to 84°28' E, with an altitude of 98 meters above sea level.

2.2 Climatic conditions in the experimental area

Prayagraj district lies within the subtropical belt of southeast Uttar Pradesh, characterized by hot summers and cold winters. Maximum temperatures can soar to 46-48°C, while occasionally dropping to 4-5°C. Relative humidity ranges from 20% to 94%, and the average annual rainfall is approximately 1100 mm.

Table:1 Physical Parameters

Particulars	Method employed
Sand(%)	Bouyoucos(1927)
Silt(%)	Bouyoucos(1927)
Clay(%)	Bouyoucos(1927)
Textural class	Bouyoucos(1927)
Bulk density(Mg m ⁻³)	Muthuval (1992)
Particle density(Mg m ⁻³)	Muthuval(1992)
Pore space(%)	Muthuval(1992)
Water holding capacity(%)	Black (1965)

Table:2 chemical parameters

Parameters	Method employed	Range
Soil P ^H (1:2)	(Jackson 1958)	(0-14 pH)
Soil EC (ds m ⁻¹)	(Wilcox 1950)	(< 0.75 ds m ⁻¹)
Organic Carbon (%)	(Walkley and Black, 1947)	(0.5-0.75 %)

Available Nitrogen (kg ha ⁻¹)	(Subbiah and Asija, 1956)	(240-480 kg ha ⁻¹)
Available Phosphorus (kg ha ⁻¹)	(Olsen <i>et al.</i> , 1954)	(11.0 -22 kg ha ⁻¹)
Available Potassium (kg ha ⁻¹)	(Toth and Prince., 1949)	(110 -280 kg ha ⁻¹)

Table3: Treatment Combinations

Treatment	Treatment combination
T1	(Absolute control)
T2	NU @ 0% + RZ @ 50% + FYM @ 50%
T3	NU @ 0% + RZ @ 100% + FYM @ 100%
T4	NU @ 50% + RZ @ 0% + FYM @ 0%
T5	NU @ 50% + RZ @ 50% + FYM @ 50%
T6	NU @ 50% + RZ @ 100% + FYM @ 100%
T7	NU @ 100% + RZ @ 0% + FYM @ 0%
T8	NU @ 100% + RZ @ 50% + FYM @ 50%
T9	NU @ 100% + RZ @ 100% + FYM @ 100%

3. RESULTS AND DISCUSSION

3.1 Effect of Nano Urea, FYM, and Rhizobium on the Physical Soil properties post-green gram harvest.

The data showed that the treatment T1 (absolute control) non-significantly influenced the bulk density of soil (1.31), particle density of soil (2.50) at 0–15 cm depth and significantly influenced percentage pore space (48.90), and water holding capacity (46.51) of soil were found to be optimal in treatment T9 (NU @ 100% + RZ @ 100% + FYM @ 100%) over T1 (absolute control) treatment at 0–15 cm and 15–30 cm depth.

3.2 Effect of Nano Urea, FYM, and Rhizobium on Chemical characteristics of soil following green gram harvest.

The data indicated that the treatment T1 (absolute control) non-significantly influenced the soil P^H at its maximum (7.05) at 0–15 cm depth, respectively (Table 5). There was a significantly influenced maximum build-up of electrical conductivity (0.20), percentage organic carbon (0.58), available N (320), available phosphorus (21.4), and available potassium (195.12) observed under the treatment T₉ (NU @ 100% + RZ @ 100% + FYM @ 100%) content in soil; however, minimum values were detected in the treatment T₁ (absolute control) at 0–15 cm depth and at 15–30 cm depth.

CONCLUSION

The conclusion based on the result. The use of liquid inorganic fertilizer and organic fertilizer significantly influenced soil health in relation to green gram cultivation. It can be concluded that application of T₉ (@100% NU + @100% RZ+ @100% FYM) was found to improve soil structure and microbial activity. Consequently, soil properties are positively impacted. nano urea provides efficient nitrogen delivery, FYM improves organic matter content and soil texture, while rhizobium inoculation boosts nitrogen fixation, collectively leading to a soil health. It can be concluded that foliar application of NU to green gram var. PDM 139 treatment (T₉) enhanced soil structure and health by increasing water holding capacity and available N,P,K in the soil.

Table 4: Effect of Nano Urea, FYM, and Rhizobium on Bulk Density, Particle Density, Pore Space, and Water Holding Capacity

S.No.	TREATMENT	B.D (Mg m ⁻³)	P.D (Mg m ⁻³)	Pore Space (%)	WHC (%)
T1	(Absolut control)	1.31	2.50	46.21	43.30
T2	NU @ 0% + RZ @ 50% + FYM @ 50%	1.30	2.50	47.35	44.38
T3	NU @ 0% + RZ @ 100% + FYM @ 100%	1.28	2.48	48.71	45.18
T4	NU @ 50% + RZ @ 0% + FYM @ 0%	1.31	2.49	46.38	43.40
T5	NU @ 50% + RZ @ 50% + FYM @ 50%	1.29	2.49	47.50	44.58
T6	NU @ 50% + RZ @ 100% + FYM @ 100%	1.28	2.47	48.82	45.61
T7	NU @ 100% +RZ @ 0% + FYM @ 0%	1.31	2.49	47.22	44.21
T8	NU @ 100% +RZ @ 50% + FYM @ 50%	1.29	2.48	47.65	45.11
T9	NU @ 100% +RZ @ 100% + FYM @ 100%	1.27	2.46	48.90	46.51
F TEST		NS	NS	S	S
S. Em. (±)		0.015	0.029	1.13	0.54
C.D (P =0.05)		0.047	0.087	3.39	1.62

Where, NU- Nano urea, RZ- Rhizobium, FYM- Farm yard manure B.D- Bulk density, P.D- Particle density, WHC- Water holding capacity

Table 5: Effect of Nano Urea, FYM, and Rhizobium on EC, Organic Carbon, Available Nitrogen, Phosphorus, and Potassium

S.N.	TREATMENT	pH	EC (ds m ⁻³)	OC (%)	Nitrogen (Kg ha ⁻¹)	Phosphorus (Kg ha ⁻¹)	Potassium (Kg ha ⁻¹)
T1	(Absolut Control)	7.05	0.17	0.38	282	17.2	180.21
T2	NU @ 0% + RZ @ 50% + FYM @50%	7.03	0.16	0.42	292	18.7	182.41
T3	NU @ 0% + RZ @ 100% + FYM @100%	6.99	0.18	0.54	298	19.3	189.26
T4	NU @ 50% + RZ @ 0% + FYM @0%	7.04	0.15	0.39	283	16.3	180.31
T5	NU @ 50% + RZ @ 50% + FYM @50%	7.03	0.17	0.46	294	18.6	183.61
T6	NU @ 50% + RZ @ 100%+FYM @100%	6.97	0.19	0.56	310	20.2	190.11
T7	NU @ 100% +RZ @ 0% + FYM @0%	7.04	0.15	0.40	284	16.4	181.11
T8	NU @ 100% +RZ @ 50% + FYM @50%	7.02	0.17	0.48	296	18.5	186.16
T9	NU @100% +RZ @100% +FYM @100%	6.96	0.20	0.58	320	21.4	195.12
F TEST		NS	S	S	S	S	S
S. Em. (±)		0.12	0.013	0.042	4.9	0.30	3.02
C.D (P =0.05)		0.37	0.25	0.126	14.8	0.92	9.06

Where, NU- Nano urea, RZ- Rhizobium, FYM- Farm yard manure,EC-Electrical Conductivity, OC- Organic carb




Fig 1. Graph representation of Av Nitrogen, Av. Phosphorus, Av. Potassium vs Treatment combination




Fig2. graph of representation of organic carbon

Disclaimer (Artificial intelligence)

Option 1:

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

Option 2:

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc have been used during writing or editing of manuscripts. This explanation will include list the name, version, model, and source of the generative AI technology and as well as the all input prompts provided to a generative AI technology

Details of the AI usage are given below:

- 1.
- 2.
- 3.

REFERENCES

1. Black, C.A. (1965) Methods of soil analysis 2, Am. Soc, Agron.madison, Wisconsin, U.S.A.

2. Bouyoucos, (1927) George J. "A rapid method for mechanical analysis of soils." *Science* 65.1692: 549-551.
3. Jackson, M. L. (1958) *Soil chemical analysis*, Second edition Indian Reprint, prentice hall of India, New Delhi. Pp-498.
4. Kumar, Vishal, et al. (2017) "Effect of macro-nutrients and farm yard manure on productivity and profitability of mustard (*Brassica juncea* L.) in Western Uttar Pradesh, India." *Asian Journal of Soil Science and Plant Nutrition* 1.31-6.
5. Meena S, Swaroop N, Dawson J. (2016) Effect of integrated nutrient management on growth and yield of green gram (*Vigna radiata* L.). *Agricultural Science Digest-A Research Journal*.36(1):63-5.
6. Muthuvel, P., Udayasoorian, C., Natesan, R. and Ramaswamy, P. P. (1992) Introduction to Soil Analysis, Tamil Nādu Agricultural University Coimbatore-641002. Nutrient Management on Production Potential and Quality of Summer Mungbean (*Vigna radiata* L.) *J. Krishi Vigyan*, 5(2), 39-45.
7. Olsen, S. R., Cole, C. V., Watanhe, F. S. and Dean, L. A. (1954). Estimation of available Phosphorus in soils by extraction with sodium bicarbonate. U. S. Deptt. Agr. Circ., 939.
8. Rathore, R. S., and R. K. Verma. (2019) "Performance of green gram variety IPM-02-3 in front line demonstrations under rainfed conditions." *International Journal of Agricultural & Statistical Sciences* 15.1.
9. Sharma, R., and Verma, M. L. (2011) Effect of Rhizobium, farm yard manure and chemical fertilizers on sustainable production and profitability of rajma (*Phaseolus vulgaris* l.) and soil fertility in dry temperate region of North-Western Himalayas. *Legume Research-An International Journal*, 34(4), 251-263.
10. Singh NB, Basna R, Verma KK, Yadav RS. (2008) Productivity and feasibility of black gram as influenced by integrated nutrient management in rainfed area of eastern U.P. *National Sym. New paradigms Agron. Res.*87-89.
11. Singh, Priya, M. L. Kewat, and Nisha Sapre (2016) "Effect of tillage and weed management practices on productivity of greengram and physico-chemical properties of soil under soybean-wheat-greengram cropping system." *JNKVV*: 92.
12. Subbiah, B. V. and Asija, E.C., (1956) A rapid procedure for Estimation of available Nitrogen in soil. *Current Science* 25(8): 259-260.
13. Subramani, T., et al. (2023) "Effect of Nano Urea on growth, yield and nutrient use efficiency of Okra under tropical island ecosystem." *International Journal of Agricultural Sciences* 19 (2023): 134-139.
14. Suryavanshi, T., Sharma, A. R., Nandeha, K. L., Lal, S., & Porte, S. S. (2018). Effect of tillage, residue and weed management on soil properties, and crop productivity in greengram (*Vigna radiata* L.) under conservation agriculture. *Journal of Pharmacognosy and Phytochemistry*, 7(1S), 2022-2026.
15. Tejaswini, V., Meyyappan, M., Ganapathy, M., and Angayarkanni, A. (2022) Effect of foliar application of organic and inorganic fertilizers on growth and yield of green gram. *The Pharma Innovation Journal*, 11(9), 939-941.
16. Toth, S. j. and Prince, A. L. (1949) Estimation of cation exchange capacity and exchangeable Ca, k, and Na content of soil by flame photometer technique. *Soil sci.*, 67, 439-445.
17. Walkley, A. and Black, I. A. (1947) Critical examination of rapid method for determining organic carbon in soils, effect of variance in digestion conditions and of inorganic soil constituents. *Soil sci pp.* 632-251.
18. Wilcox, L. V. (1950) Electrical conductivity Am water work Assoc J. 1950; 42: 775-776.

20. Bhadu , Amit, Arun Alfred David, Tarence Thomas, Anil Kumar, and Rohitash. 2023. “Effect of Inorganic Fertilizers, FYM and Nano Urea on Soil Health, Growth and Yield of Cluster Bean (*Cyamopsis Tetragonoloba* L.) Cv. HGS 563”. *International Journal of Plant & Soil Science* 35 (16):404-10. <https://doi.org/10.9734/ijps/2023/v35i163167>.
21. Beniwal , Mahendra, Saket Mishra, and Vijay Bahadur. 2024. “Effect of Foliar Application of Nano Urea, Boron and Zinc Sulphate on Growth Fruit Yield and Quality of Strawberry (*Fragaria × Ananassa* Duch.) Cv. Winter Dawn”. *Journal of Advances in Biology & Biotechnology* 27 (6):725-35. <https://doi.org/10.9734/jabb/2024/v27i6933>.
22. Wang Q, Gao L, Li Y, Shakoor N, Sun Y, Jiang Y, Zhu G, Wang F, Shen Y, Rui Y, Zhang P. Nano-agriculture and nitrogen cycling: Opportunities and challenges for sustainable farming. *Journal of Cleaner Production*. 2023 Aug 17:138489.