

# Understanding Pollen Behaviour under High Temperature for Climate Resilient Breeding

## ABSTRACT

Global warming raise challenges for plant reproduction as pollen development and functioning are among the most heat-sensitive processes. Hence, it is crucial to understand the mechanisms and processes underlying heat-related male sterility in order to maintain food security. Elevated temperatures elicit acclimation responses that permit pollen development under restricted heat stress conditions; and physiological injury leading to failure of pollen development and functioning occurs at higher temperature stress. Pollen and the surrounding anther tissues respond to an increase in temperature at the transcriptome, proteome, and metabolome levels. To counteract the damaging effect of misfolded proteins under heat stress HSPs () accumulate in the cytoplasm and organelles to stabilize, resolubilize, and refold proteins. The pathways leading to the production of carbohydrates, amino acids, phenolic compounds, polyamines, hormones and lipids are interconnected and contribute to metabolic homeostasis required for growth and viability of the pollen. The pathways leading to the production of carbohydrates, amino acids, phenolic compounds, polyamines, hormones and lipids are interconnected and contribute to metabolic homeostasis required for growth and viability of the pollen. Use of molecular markers to introgress specific genomic regions associated with a specific trait along with QTL() fine mapping can identify several candidate genes associated with thermo-tolerance. A deep understanding of the molecular mechanisms and metabolic processes involved in the stress response to high temperatures in flowers and, particularly, in the male reproductive organs will be a major step towards development of effective breeding strategies for high and stable production in crop plants.

Keywords: *Pollen, high temperature, breeding, thermotolerance*

## 1. INTRODUCTION

Plants are sessile organisms that are exposed to changing environments and needs to get adapted to maintain organismal and cellular homeostasis. High or low temperature stresses can be detrimental to all phases of plant development. A clear-cut understanding of behavioural responses of plants to stress during their reproductive phase is critical to manage agricultural productivity as the majority of our food supply is a product of sexual reproduction in flowering plants. The development and functioning of the male gametophyte, or pollen, are known to be among the most temperature-sensitive processes within the plant life cycle [1]. Exposure to high temperature episodes often coincides with the reproductive phase of the plant life cycle. As pollen development and functioning are among the most heat-sensitive processes that impact upon plant fertility, it is crucial to understand the mechanisms and processes underlying heat-related male sterility in order to maintain food security.

## 2. PLANT REPRODUCTION AT HIGH TEMPERATURE

Reproductive growth reduces production potential significantly because it is more susceptible and has a variety of negative impacts, including the loss of buds, flowers, fruits, pods, and seeds [2]. The impact of heat stress on reproductive components during development lowers the harvest index, and the degree and duration of the stress cause different reactions in crops [3]. The development of male and female gametophytes (pollen grain and embryo sac), flower initiation, differentiation of male and female floral parts, micro and megasporogenesis, pollination, micro and megagametogenesis, fertilization, and seed development comprise the reproductive phase. Elevated temperatures cause distinct reactions in each stage; yet, all of these reactions have the same overall impact of decreasing net yield. Heat stress causes problems for both male and female gametophytes. It also inhibits the growth of pollen tubes, reduces ovule function and stigma receptivity, limits embryogenesis, reduces ovule viability, increases ovule abortion, and results in poor seed set [4].

Comment [D1]: Delete it

Comment [D2]: Which temperature i.e (high or low)?

Comment [D3]: Write complete name

Comment [D4]: Correct this word

Comment [D5]: Write complete name

Comment [D6]: Delete it

Comment [D7]: Delete it

Comment [D8]: Delete it

Comment [D9]: Mention (in high temperature)

## 2.1 Pollen development

Sexual reproduction has been shown to be restricted by the sensitivity of the male gametophyte (pollen) to environmental changes and inadequate conditions [5]. Within the anther locules, diploid pollen mother cells undergo meiosis to produce a tetrad of four haploid microspores encased in locular fluid, which is then used to make pollen. The liberated microspores expand and divide asymmetrically (pollen mitosis I) to create a bigger vegetative cell and a smaller generative cell following their liberation from the tetrad. Then, either prior to pollen being expelled from the anther or during the formation of the pollen tube, the generative cell is engulfed by the vegetative cell and goes through a second mitosis (pollen mitosis II) to produce two sperm cells [6]. The tapetum is formed by the innermost layer of the anther wall during the development of the pollen mother cells. This tissue is metabolically active, especially in the early stages of microspore development. It supplies the growing microspores with nutrients, carbohydrates, enzymes, and other substances needed for the creation of the exine, or outer pollen wall. The degeneration of the tapetum happens soon after the microspores are released from the tetrad and is closely synchronized with the development of the microspores. Pollen formation depends on the tapetum's proper operation and timely degradation.

Comment [D10]: Add the reference

## 3. HEAT-INDUCED POLLEN DEFECTS

Pollen growth is adversely affected by both transient high temperatures and persistently modestly higher day and nighttime temperatures. *Meiosis is when the first heat-induced developmental abnormalities appear*, chromosomal behavior and meiotic cell division may be impacted in addition to increased frequency of crossing over and homologous recombination, resulting in uneven chromosomal separation between spores and the creation of diploid dyads [7]. The abnormal spindle orientation caused the chromosomes to behave as they did during cell division. Microtubules and cytoskeleton dynamics have been examined in vegetative tissues and during pollen tube formation, and it is well known that high temperatures have an impact on these structures. While expanding pollen tubes are more susceptible (35°C, 3 h) and damage rises with increasing temperature, this requires temperatures of above 40°C in vegetative cells in tobacco or *Arabidopsis* [8]. Chromosome aberration during meiosis appears to be more common in situations with extreme heat stress [9].

Comment [D11]: Change this sentence and write (Meiosis appears when the first heat-induced developmental abnormalities)

Comment [D12]: Write italic

## 4. THE IMPACT OF HEAT STRESS ON POLLEN QUALITY

### 4.1 Pollen Viability

Pollen is one of the most sensitive organs to heat stress. As a result, there may be less fruit set and sterile pollen [10]. Consequently, the temperature at which pollen develops affects crop yield [11]. Numerous species have previously shown how heat stress affects pollen viability, including tomato [10], rice [12], chickpea [13], and soybean [14].

### 4.2 Pollen Development

Excessive temperatures cause a stop in the early phases of pollen formation. Barley is likewise significantly impacted by heat stress during meiosis [15]. While heat stress at later developmental stages had no effect on pollen viability, it reduced fruit set, seed quantity, and pollen viability in bell peppers during microspore mother cell meiosis [16]. Meiosis is a typical sensitive stage for most crops, and the developmental sensitivity to heat stress in pollen appears to be species-specific.

Comment [D13]: Add the reference

### 4.3 Tapetum

Key to providing metabolites to the pollen is the tapetum, an organ whose growth is highly susceptible to heat stress. For instance, a high temperature of 30 °C in barley caused an early meiotic prophase I and an early tapetum breakdown [17]. There have also been reports of common bean tapetum degradation under heat stress [18]. Ten days prior to anthesis, in common beans, heat stress caused structural anomalies in the endoplasmic reticulum pattern of the tapetum, which in turn caused the tapetum to prematurely degenerate.

Comment [D14]: Add reference

### 4.4 Opening of Loculi

Heat stress in rice impacted the anther loculi's opening, which decreased pollen fertility [12]. Tomatoes also showed the same outcome. Therefore, preventing the anther loculi from opening could prevent pollen grains from being released.

Comment [D15]: Add reference

### 4.5 Pollen Germination

Heat stress can also affect pollen germination, which can stop pollen tubes from growing in addition to pollen development. [19] found that a decrease in pollen germination was mostly responsible for the tomato fruit set drop at increased temperatures. Early pollen development failure may be the cause of reduced germination of pollen under heat stress. Additionally, cotton's in vitro pollen germination decreased in response to high temperatures [20].

Comment [D16]: Add reference

Comment [D17]: Change to (the)

Comment [D18]: Change to (of)

Comment [D19]: Add (,)

Comment [D20]: Add (;)

## 5. IMPACT OF HEAT STRESS ON POLLEN METABOLITES

### 5.1. Carbohydrates

Carbohydrates are essential for plant homeostasis because they provide the energy needed for pollen production and germination, but they also play a part in osmotic balance maintenance, stress signaling, and membrane protection. In tomato plants, blooms that were grown at a temperature higher than 32 °C displayed decreased pollen viability. These flowers displayed an increase in soluble sugars in the locular fluid but a decrease in soluble sugars in the pollen and anther wall [21]. Soluble sugars collected in the locular fluid as a result of pollen being modified by high temperatures and decreasing in requirement of them [22]. A failure in pollen development would likely result from changes in the accumulation of carbohydrates at high temperatures, which would also likely reduce the availability of energy resources and the osmotic power of carbohydrates.

Comment [D21]: Add reference

### 5.2. Proline

Proline has been demonstrated to be involved in preserving cellular homeostasis, protecting membrane integrity, and scavenging reactive oxygen species [19]. By preserving the hydration shells surrounding molecules, proline has the ability to stabilize proteins. Proline content in the anthers of heat-tolerant and heat-sensitive cowpea cultivars grown in high temperatures (45 °C maximum daily temperature) was analyzed. Heat-sensitive cultivars accumulated the largest quantities of proline in their anthers, although tolerant cultivars' mature pollen had a higher proline abundance than sensitive cultivars' pollen.

Comment [D22]: Add reference

### 5.3. Lipids

Membrane fluidity is significantly influenced by lipids, particularly in stressful situations. Bound unsaturated fatty acids became more abundant at high temperatures, but saturated fatty acid quantity fell. The authors hypothesised that an increase in unsaturated fatty acids, which result in an increase in membrane fluidity because of the presence of double bonds, may be the source of the damage to membranes produced by high temperatures. The membrane is more susceptible to ROS() assaults because of the unsaturated fatty acid moiety. A change in phospholipid saturation was linked to a decline in pollen viability [14].

Comment [D23]: Write complete word

They can preserve membrane integrity and function as ROS scavengers. By adding spermidine or spermine to the germination medium, it was possible to increase in vitro pollen germination in tomatoes after incubating pollen extract in the media for 20 hours at 33 °C [23]. This finding revealed that the amount of polyamines was reduced by heat stress. In fact, spermidine and spermine contents in pollen were decreased after 4 hours of incubation at 38 °C, whereas putrescine content rose. These variations were linked to a decline in pollen germination [23].

### 5.5. Hormones

Steroid hormones called brassinosteroids are involved in how the body reacts to a variety of abiotic stressors, including heat, salt, and drought. They participate in ROS scavenging and interact with heat shock proteins [24]. The reaction to heat stress also involves ethylene. When plants were treated with ethylene, the amount of pollen that was non-viable under prolonged heat stress (32 °C/26 °C) decreased due to a mutation in the ethylene receptor (nr) [25]. Auxin levels in the anthers of barley and arabidopsis reduced after heat stress (30 °C/25 °C for 5 days), and this was connected with male sterility [26]. Male sterility was reversed by exogenous auxin treatment. Auxin content was found to be connected with a decrease in pollen viability and germination in the anthers of rice plants that were subjected to 39 °C for 4 hours each day for 5 days [27]. In hot weather, ABA() concentrations rose while GA() concentrations fell. Pollen germination was previously found to be negatively correlated with high ABA levels [28].

Comment [D24]: Write italic

Comment [D25]: Write complete word

Comment [D26]: Write complete word

## 6. HEAT RESPONSES

### 6.1 Acclimation

Like other plant cell types, the pollen and the surrounding anther tissues react to a rise in temperature at the transcriptome, proteome, and metabolome levels. It was discovered that these reproductive tissues exhibited a number of heat-related reactions that have been identified as adaptive in vegetative tissues.

Comment [D27]: Change to (amount)

Comment [D28]: Add reference

#### 6.1.1 Protein Homeostasis in the Cytosol: The Heat Shock Response

One of the main damaging effects of high temperature results from changes in protein structure. This may interfere with protein function, and if more hydrophobic regions are exposed, proteins can aggregate and become cytotoxic. Failure to prevent the accumulation and aggregation of misfolded proteins may eventually lead to cell death. To counteract these effects, the expression of heat shock protein (HSP) chaperones is induced at high temperature in a process known as the heat shock response (HSR). HSPs accumulate in the cytoplasm and organelles to stabilize, resolubilize, and refold proteins [20]. Underlying the HSR is a network of heat stress transcription factors (HSFs) that bind to a palindromic DNA

sequence, the heat shock element, to induce the expression of heat-responsive genes [30]. Proteins of different types of HSPs (i.e. belonging to diverse families, like sHSP, HSP70, HSP90, and HSP100) accumulate in developing anthers and pollen grains after a short period of high temperature stress [31], which points to a capacity for both tissues to activate "classical" thermotolerance mechanisms. One of the main HSFs regulating the HSR is HSFA2, which forms a "superactivator complex" with HSFA1 proteins [32].

### **6.1.2 Protein Homeostasis in the ER: The Unfolded Protein Response**

A second collection of genes, including those encoding chaperones, is activated in response to high temperatures in order to shield cells from harmful concentrations of unfolded proteins in the endoplasmic reticulum (ER) [33]. The unfolded protein response (UPR) is a phenomenon where an overabundance of misfolded proteins in the endoplasmic reticulum (ER) results in increased expression of certain ER protein folding machinery components. This response is triggered by two different pathways: one relies on the ER membrane releasing bZIP28/bZIP17, and the other is triggered by IRE1 splicing bZIP60 in an alternate manner [34]. A number of UPR genes are activated by heat, and it has been demonstrated that seedlings of a bZIP28 knockout mutant are heat-sensitive, indicating that the UPR plays a crucial part in the body's overall response to heat stress and thermotolerance. The UPR is up-regulated at the transcript and protein levels in male reproductive organs during or soon after a heat shock or prolonged exposure to mild heat [35], indicating the UPR has a role in acclimating developing pollen to heat. By examining a double knockout mutant of ire1a and ire1b, which deactivates the RNA-splicing component of the UPR signaling pathway. At room temperature, the mutant was found to be fertile, but at slightly higher temperatures, it became sterile in males due to decreased viability of mature pollen and changed pollen coat composition [36].

Comment [D29]: Change to (the)

### **6.1.3 Reactive Oxygen Species Scavenging**

The buildup of reactive oxygen species is partially blamed for the cellular damage caused by high temperatures (ROS). ROS are hazardous metabolic byproducts that also function as signaling molecules that control stress reactions [37]. Cells feature substantial ROS scavenging and detoxifying machinery, which comprises of enzymes such as catalase, ascorbate peroxidase (APX), and superoxide dismutase, as well as antioxidant compounds including ascorbic acid and flavonoids [38]. Under steady state conditions, there is a controlled equilibrium between production and scavenging because many of the genes linked to ROS scavenging are responsive to ROS levels. This equilibrium can be upset by exposure to high temperatures, as heat quickly causes ROS to build up and causes a secondary oxidative stress. Heat quickly upregulates the expression of ROS scavengers and antioxidant levels to deal with the excess amount of ROS that arises [39].

Comment [D30]: Change to (because of)

Comment [D31]: Add (that the)

## **6.2 Collapse**

### **6.2.1 Defects related to pollen failure under high temperature**

The surrounding tapetal cells and growing pollen have a high sensitivity to heat stress, which frequently causes the tapetal cells to degenerate prematurely and the developing pollen to undergo abnormal developmental or programmed cell death. A number of physiological traits of growing pollen may be connected to pollen failure under hot conditions. Primarily, growing pollen and tapetal cells have a large amount of mitochondria. Consequently, increased respiration in response to heat stress may produce a disproportionate amount of reactive oxygen species (ROS) that are too large for the protective cellular processes to adequately detoxify, leading to damage to various cell components. Secondly, diminished transport of carbohydrates and other components required for optimal pollen growth may arise from premature tapetum degeneration or impacts on particular metabolic enzymes under heat stress. When combined, the decreased availability of carbohydrates and the elevated respiration caused by a high number of mitochondria may result in energy reserve depletion and developmental abnormalities in the future. Thirdly, proteins unfold in response to heat. The classical chaperone heat stress response (HSR) usually reduces this effect. Heat causes HSFs (), crucial HSR signaling components, to activate in pollen; yet, the cell is unable to establish a complete HSR equivalent to vegetative tissues, which is insufficient to shield and refold proteins. Lastly, it is known that heat stress and reactive oxygen species can affect microtubules and the cytoskeleton. Heat alters the spindle apparatus's orientation during meiotic cell division, resulting in abnormal chromosomal behavior and the failure of pollen formation [40].

Comment [D32]: Change to (number)

Comment [D33]: Write complete word

### **6.2.2 Impact of high temperature stress on multiple functions of tapetum**

A number of plant species link stress-induced spore abortion and male sterility to changes in tapetal development and sporadic abnormalities in the endothecium and middle layer, two of the surrounding cell layers [41]. A single layer of endopolyploid cells that encircles the locules of growing microspores and pollen grains makes up the tapetum, an integral component of the male sporangium [42]. The tapetum is very metabolically active during normal sporogenesis and acts as a nutritive source by giving the nearby microspores energy and necessary nutrients [43]. Moreover, the tapetum, a secretory cell layer, supplies cell wall components, such as sporopollenin, for the development of the pollen exine layer and enzymes for the release of microspores from the meiotic tetrad [44]. The tapetal cells undergo programmed cell death (PCD) and then disintegrate at PMI (), a later stage of male gametogenesis [45]. For appropriate microspore formation, pollen maturation, and fertility, this controlled tapetum-specific PCD () and disintegration is necessary.

Comment [D34]: Write complete name

Comment [D35]: Write the complete name

Comment [D36]: Add reference

When anthers are subjected to heat stress, they usually exhibit dramatic changes in microspore formation along with an early loss of the tapetal cell layer. These tapetal abnormalities severely impair the development of male gametogenesis and prevent microspores from completing PMI (due to loss of nutritional supply. Severe subcellular changes are seen in the tapetum and the outer three anther wall layers (endothecium, middle layer, and epidermis) including hypertrophy (increased vacuolization), overdevelopment of chloroplasts, and excessive swelling of mitochondria [46]. Furthermore, elevated temperatures lead to a significant suppression of transcription, which in turn significantly lowers the ability of the tapetum and its surrounding anther layers to divide.

Comment [D37]: Write complete name

Comment [D38]: Add reference

Growing microspores gather photoassimilates like starch and other carbohydrates and form a potent photosynthetic sink. The carbohydrate reserve is rather modest during the uninuclear microspore stage, but after PMI, growing spores usually exhibit a fast phase of starch production and rapidly acquire substantial amounts of starch [47]. The amount of starch gradually drops during final pollen maturation as it is broken down into soluble sugars. Over time, accumulated anther sugars provide energy for pollen tube formation, microsporogenesis, and pollen maturation. They also act as an osmolyte, providing pollen resistance to abiotic stresses such as desiccation (sucrose, for example, preserves membrane integrity). Therefore, carbohydrates play a crucial role in determining pollen viability and germination capacity in addition to being necessary for pollen growth [48].

Comment [D39]: Change this word

#### 6.2.2.1 Sugar metabolism and transport in the tapetal cell layer and microspores

Changes in assimilate supply or modifications in sugar metabolism are the two main causes of the generally decreased accumulation of starch and other carbohydrate components in abiotically stressed microspores. A lower sugar supply to the reproductive tissues is thought to be the cause of the failure of male spore formation during abiotic stress [49]. Rather than changes in sucrose supply, the changed carbohydrate profile in stressed anthers is due to changes in sugar metabolism and use. In anthers under abiotic stress, the primary enzymes responsible for the metabolism of sucrose and starch exhibit decreased activity. Differential expression of genes related to sugar cleavage and utilization, sugar transport, and starch synthesis has been observed in heat-stressed microspores of sorghum [50]. More precisely, as has already been noted in heat-stressed tomato anthers (32/26 °C Day/night), season-long heat stress (36/26 °C Day/night) dramatically lowers the transcript level of cell wall invertase (CWI) in the male gametophyte [51].

Comment [D40]: Change this word

Comment [D41]: Write italic

The transcriptional down-regulation of the invertase b-d-fructofuranosidase is typically the reason for the absence of starch in abiotically challenged microspores. Invertase enzymes are needed in conjunction with sucrose synthase to break the delivered sugar module. Invertase (INV) is the predominant enzyme that regulates the cleavage of sucrose in spores and anthers of wheat and several other species [50]. Moreover, extracellular CWI invertases are essential for the unloading of sucrose from the tapetum into the apoplastic space and the creation of a soluble sugar gradient along the various anther fractions because developing microspores are physically isolated from the surrounding tapetal cell layer (no symplastic PD channels). Due to the specific down-regulation of CWI invertase in anthers under abiotic stress, there are substantial abnormalities in pollen development and viability. This is because it prevents the import of hexose sugar units into growing microspores and hinders the buildup of starch in maturing pollen. Stress-induced down-regulation of INV not only reduces starch accumulation in spores but also modifies the consumption of sucrose in other anther tissues. As a result, it frequently results in the ectopic buildup of sugars in organ types other than microspores, like the endothecium and connective tissue [52]. The morphological abnormalities in anthers under abiotic stress may be attributed to these modifications in the partitioning of anther sugar.

Comment [D42]: Write complete name

Comment [D43]: Write complete name

Comment [D44]: Add reference

Combinatorial effects of two sugar metabolism pathways, such as reduced sucrose transport and altered sucrose metabolism, result in anther-specific changes in sugar content and the corresponding induction of male sterility under abiotic stress.

Comment [D45]: Add reference

#### 6.2.3 Starvation hypothesis of tapetal vulnerability to abiotic stress

According to the starvation theory, abiotic stress results in the downregulation of a tapetal cell wall invertase, which lowers the amount of sugars that reach the growing microspores [42]. In vegetative tissues, abiotic stress causes an increase in the synthesis of ABA. The gene(s) encoding tapetal cell wall invertase(s) are downregulated as a result of the hormone's transportation to the tapetum. There is a decrease in the breakdown of sucrose and a compromise in the supply of hexose to the tapetum and the growing microspores. Reduced GA levels have the potential to affect tapetal growth, delay PCD, and cause tapetal starch mobilization.

Comment [D46]: Write the complete word

Comment [D47]: Write the complete word

Comment [D48]: Add reference

### 7. POLLEN AS A SCREENING TOOL FOR HEAT TOLERANCE.

Fruit set and production will be impacted by brief bursts of extreme weather, such as high temperatures, which are expected to happen more frequently in the future climate. As a result, it would be beneficial for plants to demonstrate increased reproductive resilience in the harsh temperatures typically encountered during plant reproduction as well as for yield-producing processes like fertilization, embryo development, pollen tube growth, pollen grain development, and germination. It has been noted that two phases of pollen development—mature microspores at anthesis and microspore

Comment [D49]: Change this word

mother cell meiosis [51], are extremely susceptible to high temperatures [16]. Reduced pollen germination under high temperatures was the main factor contributing to decreased pollen fertility [53]. Finding out how well pollen germination and pollen tube growth function across a broad temperature range will help predict how the plant will grow and set fruit at various temperatures.

Comment [D50]: Add reference

### 7.1 Pollen Viability an Index of Abiotic Stresses Tolerance

In many crop species, temperatures over or equivalent to 32 °C resulted in reduced pollen viability, retention of pollen in the anthers, and pollen germination [54]. The impact of heat stress on pollen viability is linked to changes in the anther's development-related glucose metabolism. The effects of high temperatures have been well shown in a number of crop types. When wheat florets are developing, excessive temperatures might result in total sterility. When pollen mother cells were dividing in the early reproductive phase of a wheat plant, for example, they were subjected to heat stress for three days straight at 32 °C. This caused pollen sterility, which significantly reduced seed laying. Pollen mother cells fail during meiosis when sorghum pollen is killed at temperatures of > 42 °C and 10 °C at night, although this has no effect on the survivability of the female pollen.

Comment [D51]: Italic

Comment [D52]: Add reference

#### 7.1.1 Pollen viability assessment

Numerous techniques can be used to indicate pollen viability. To evaluate pollen vitality, however, a lot of in-vitro germination and pollen viability data have been used [55]. Pollen viability is also a prerequisite for high crop output and is crucial to the hybridization process [56]. The vigor and fertility of the pollen can be used to determine its quality. The rate of pollen tube germination and pollen germination speed are referred to as vigour. The proportion and vigor of pollen germination have been measured over time using in vitro assays [57]. Assessment of pollen fertility and germination potential are key criteria for pollen evaluation, and genotypes can be regarded good pollinators if their pollen viability is high [58]. Additionally, pollen viability is calculated following the use of chemical hybridizing agents to induce sterility [59]. In the past, several stains, such as pollen viability tests, have been employed to determine the relative estimate of fertilization potential and to evaluate pollen vitality [60]. The assessment of pollen during storage, crop improvement, genetics and fertility research, and crop breeding all depend on the application of trustworthy techniques for determining the functional quality of pollen [61]. Additionally, in a number of fruit species, there is a linear relationship between pollen fertility and germination capacity. Although stain tests were quick and simple, pollen germination techniques were occasionally necessary to determine pollen fertility [62].

#### 7.1.2 Methods for improving pollen viability

To increase the number of viable blooms, treat seed-producing plants with a fungicide from the strobilurin group, preferably pyraclostrobin. There are various methods to administer fungicide, such as directly onto the seeds or into the soil before planting. Alternatively, it could be foliar sprayed following germination. [63] Research aimed at determining the impact of boron on rice reproductive growth, stigma receptivity, pollen viability, and yield qualities revealed that stigma receptivity was reported at 0.4 ppm, and mean pollen vitality was 86% [64]. Moreover, rice plants treated with salicylic acid at 0.1–10 mM under heat stress demonstrated a substantial increase in pollen viability and seed-setting rate in comparison to the NON-SA treatment [65]. The effects of nitrogen and phosphorus, respectively, on Cucurbita pepo (Cucurbitaceae) pollen production and performance. The mean total number of staminate flowers per plant was unaffected by nitrogen treatments, however it did impact the mean size and number of pollen grains per staminate flower. The commercially suggested dosage of pollen produced a higher proportion of seeds and fertilized the ovules with faster-growing pollen tubes. The pollen was developed by the high nitrogen treatment plants. However, phosphorus treatments had a notable impact on both male and female reproductive production. Compared to low treatment soils, pollen originated from the high phosphorus treatment plants, which produced more seeds and met the commercially recommended dose. Nonetheless, it has been determined that variations in pollen performance can cause heterogeneity in the soil's phosphorus and nitrogen content to have a substantial impact on the proportion of seeds [66].

Comment [D53]: Add reference

Comment [D54]: Better delete this paragraph

## 8. BREEDING FOR POLLEN THERMO-TOLERANCE

### 8.1 QTL Mapping for Thermo-Tolerance

Wheat cultivars with markers linked to the length of grain filling during heat stress have already been screened using marker assisted selection [67]. By classical breeding, thermo-tolerance could also be incorporated into a sensitive genotype via a marker linked to pollen thermo-tolerance. QTL studies for thermo-tolerance traits have been performed in two ways: (i) by measuring fruit set in rice [71] and tomato [70], or (ii) by evaluating pollen viability in maize, rice [68], and tomato [69]. Breeders can use QTL identification as the basis for more in-depth analyses, such as QTL fine-mapping, which may identify more closely linked molecular markers. These markers can then be used by breeders to screen seedlings for genotypes that are thermotolerant and don't have too much linkage drag. Fine mapping may eventually even result in the identification of the essential genes causing thermotolerance. By counting the pods per peduncle in a population of recombinant inbred lines, cowpea QTLs associated with thermo-tolerance have been found [72]. As a result,

Comment [D55]: Mention complete name

numerous candidate genes encoding proline transporters, heat shock transcription factors, and heat shock proteins were found.

Comment [D56]: Add reference

Metabolite profiles, especially in pollen, alter at different temperatures in addition to pollen viability and fruit set. An alternate method of phenotyping pollen thermotolerance is to map the variance in levels of particular metabolites linked to thermo-tolerance. Pollen from genotypes that are thermotolerant retains its sugar levels, while sensitive genotypes see a reduction in sugar levels [73]. It was proposed that sugar level, which is easier to detect and more reliable than pollen viability, would be a useful thermotolerance criterion. Therefore, discovering metabolite QTLs (mQTL) linked to thermotolerance may be a helpful method to gain fresh understanding of the processes granting thermotolerance [74]. A mapping population's segregation of these metabolites can be used to pinpoint chromosomal areas that may be connected to pollen thermotolerance. The key to identifying thermo-tolerant QTLs and predictive molecular markers is the creation of a good mapping population and simple, trustworthy phenotyping techniques. There are other methods besides QTL analysis to find potential genes linked to pollen thermotolerance. On the other hand, thorough understanding of the desired characteristic may identify particular candidate genes, whose function in pollen thermotolerance can be examined using reverse genetics techniques (Figure 1) [75], in which the candidate gene is either overexpressed or downregulated by transgenic or tilling techniques. Important genes involved in thermo-tolerance may be identified through over-expression or down-regulation of flavonoid pathway genes. These genes might serve as the foundation for discovering (via diversity screens) or producing (through tilling) unique genetic variation that can be utilized as a marker in pollen thermotolerance breeding.

Comment [D57]: Change this word

Comment [D58]: Add reference




Figure 1: Breeding approaches to improve crop thermo-tolerance

There are two distinct, so-called reverse genetics methods for validating candidate genes. The first involves analyzing the phenotypic of a population that has been tilled (targeted induced local damages in genomes) by focusing on a particular gene. Seeds are mutagenized at random to create a tilling population [76]. The second strategy involves creating transgenic lines where the chosen gene may be either overexpressed or silenced using a constitutive or tissue-specific promoter, for example, or by utilizing an antisense construct. The phenotypic study of these lines might verify that the variation in the population's trait of interest is, in fact, caused by the candidate gene. An over-expression of a potential gene linked to pollen thermo-tolerance, for instance, would be expected to result in higher performance at high temperatures than the wild-type.

Comment [D59]: Delete it

Comment [D60]: Add reference

## 9. CONCLUSION

During the crop growth cycle, the reproductive phase is more susceptible to high temperatures than the vegetative phase. Male reproductive organs are more susceptible to heat stress than female reproductive systems, but the whole reproductive process—from gamete formation to fertilization and seed maturity—is very susceptible to high temperatures. Pollen viability, pollen tube development, and decreased stigma sensitivity all prevent fertilization. Under heat stress, plants respond in a number of ways that provide tolerance, such as by producing signal cascades that significantly alter the expression of specific genes. Making use of "omics" (genomics, transcriptomics, proteomics, and metabolomics) is essential to comprehending the molecular foundations and mechanisms of the plant's heat stress tolerance systems. Because of molecularly linked functional physiology, plants can be developed with increased tolerance to harsh conditions seen in arid and semi-arid regions of the world, resulting in more profitable yields.

Comment [D61]: Change this word

## AVAILABILITY OF DATA AND MATERIALS

The authors confirm that the data supporting the findings of this study are available within the article.

## REFERENCES

1. Zinn KE, Tunc-Ozdemir M, Harper JF. Temperature stress and plant sexual reproduction: uncovering the weakest links. *J Exp Bot.* 2010;61:1959–1968.
2. Thakur P, Kumar S, Malik JA, Berger JD, Nayyar H. Cold stress effects on reproductive development in grain crops: an overview. *EnvironExpBot.* 2010;67:429–443.
3. Harsant J, Pavlovic L, Chiu G, Sultmanis S, Sage TL. High temperature stress and its effect on pollen development and morphological components of harvest index in the C3 model grass *Brachypodium distachyon*. *J Exp Bot.* 2013;64: 2971–2983.
4. Sita K, Sehgal A, Kumar J, Kumar S, Singh S, Siddique K H M, et al. Food Legumes and rising temperatures: Effects, Adaptive Functional Mechanisms Specific to Reproductive Growth Stage and Strategies to Improve Heat Tolerance. *Front Plant Sci.* 2017;8:1658.
5. Kim S, Hong C, Lee I. Heat shock stress causes stage-specific male sterility in *Arabidopsis thaliana*. *J Plant Res.* 2001;114: 301–307. <https://doi.org/10.1007/PL00013991>
6. McCormick S. 2004. Control of male gametophyte development. *Plant Cell (Suppl).* 2004;16:142–153.
7. Omid M, Siahpoosh MR, Mamghani R, Modarresi M. The influence of terminal heat stress on meiosis abnormalities in pollen mother cells of wheat. *Cytologia.* 2014;79:49–58.
8. Parrotta L, Faleri C, Cresti M, Cai G. Heat stress affects the cytoskeleton and the delivery of sucrose synthase in tobacco pollen tubes. *Planta.* 2016;243:43–63.
9. De Storme N, Geelen, D. The impact of environmental stress on male reproductive development in plants: biological processes and molecular mechanisms. *Plant, Cell Environ.* 2014;37:1–18.
10. Abdul-Baki AA, Stommel JR. Pollen viability and fruit set of tomato genotypes under optimum and high-temperature regimes. *HortScience.* 1995;30:115–117.
11. Ploeg Van der A, Heuvelink E. Influence of sub-optimal temperature on tomato growth and yield: A review. *J HortSci Biotech.* 2005;80:652–659.
12. Matsui T, Omasa K. Rice (*Oryza sativa* L.) cultivars tolerant to high temperature at flowering: Anther characteristics. *Ann Bot.* 2002;89:683–687.
13. Devasirvatham V, Gaur PM, Mallikarjuna N, Tokachichu RN, Trethowan RM, Tan DKY. Effect of high temperature on the reproductive development of chickpea genotypes under controlled environments. *Funct. Plant Biol.* 2012; 39:1009–1018.
14. Djanaguiraman M, Prasad PVV, Boyle DL, Schapaugh WT. Soybean pollen anatomy, viability and pod set under high temperature stress. *J Agron Crop Sci.* 2013; 199:171–177.
15. Sakata T, Takahashi H, Nishiyama H, Higashitani A. Effects of high temperature on the development of pollen mother cells and microspores in barley *Hordeum vulgare* L. *J. Plant Res.* 2000;113:395–402.
16. Erickson AN, Markhart AH. Flower developmental stage and organ sensitivity of bell pepper (*Capsicum annuum* L.) to elevated temperature. *Plant Cell Environ.* 2002;25:123–130.
17. Oshino T, Abiko M, Saito R, Ichiishi E, Endo M, Kawagishi-Kobayashi M. et al. Premature progression of anther early developmental programs accompanied by comprehensive alterations in transcription during high-temperature injury in barley plants. *Mol Genet Genomics.* 2007;278:31–42.
18. Suzuki K, Takeda H, Tsukaguchi T, Egawa Y. Ultrastructural study on degeneration of tapetum in anther of snap bean (*Phaseolus vulgaris* L.) under heat stress. *Sex. Plant Reprod.* 2001;13:293–299.
19. Hazra P, Ansary, SH, Dutta AK, Balacheva E, Atanassova B. Breeding tomato tolerant to high temperature stress. *Acta Hort.* 2009; 830:241–248.
20. Kakani VG, Reddy KR, Koti, S, Wallace, TP, Prasad PV, Reddy VR, Zhao D. Differences in in vitro pollen germination and pollen tube growth of cotton cultivars in response to high temperature. *Ann Bot.* 2005;96:59–67.
21. Sawhney VK, Bhadula SK. Microsporogenesis in the normal and male-sterile stamenless-2 mutant of tomato (*Lycopersicon esculentum*). *Can J Bot.* 1988; 66: 2013–2021.

Comment [D62]: Write italic

22. Pressman E, Peet MM, Pharr DM. The effect of heat stress on tomato pollen characteristics is associated with changes in carbohydrates concentration in the developing anthers. *AnnBot.* 2002;631–636
23. Song J, Nada K, Tachibana S. Suppression of S-adenosylmethionine decarboxylase activity is a major cause for high-temperature inhibition of pollen germination and tube growth in tomato (*Lycopersicon esculentum* mill.). *Plant Cell Physiol.* 2002;43:619–627.
24. Mazorra LM. Brassinosteroid action and its relation with heat stress mechanisms in plants. In: Hayat S, Ahmad A, editors. *Brassinosteroids: A Class of Plant Hormone*. Springer: New York, USA ;2011
25. Firon N, Pressman E, Meir S, Khoury R, Altahan L. Ethylene is involved in maintaining tomato (*Solanum lycopersicum*) pollen quality under heat-stress conditions. *AoB Plants*. 2012; doi:10.1093/aobpla/pls024
26. Sakata T, Oshino T, Miura S, Tomabechi M, Tsunaga Y, Higashitani, N, et al. Auxins reverse plant male sterility caused by high temperatures. *Proc Natl Acad Sci.* 2010;107:8569–8574.
27. Tang RS, Zheng JC, Jin, ZQ, Zhang DD, Huang YH, Chen LG. Possible correlation between high temperature-induced floret sterility and endogenous levels of IAA, GA and ABA in rice (*Oryza sativa* L.). *Plant Growth Regul.* 2008;54:37–43.
28. Frascaroli E, Tuberosa R. Effect of abscisic acid on pollen germination and tube growth of maize genotypes. *Plant Breed.* 1993;110:250–254.
29. Hartl FU, Bracher A, Hayer-Hartl M. Molecular chaperones in protein folding and proteostasis. *Nature.* 2011;475:324–332.
30. Ohama N, Sato H, Shinozaki K, Yamaguchi-Shinozaki K. Transcriptional regulatory network of plant heat stress response. *Trends Plant Sci.* 2017;22: 53–65.
31. Chaturvedi P, Doerfler H, Jegadeesan S, Ghatak A, Pressman E, Castillejo MA, Wienkoop S, Egelhofer V, Firon N, Weckwerth W. Heat treatment-responsive proteins in different developmental stages of tomato pollen detected by targeted mass accuracy precursor alignment (tMAPA). *J Proteome Res.* 2015;14:4463–4471.
32. Scharf KD, Berberich T, Ebersberger I, Nover L. The plant heat stress transcription factor (Hsf) family: structure, function and evolution. *Biochim Biophys Acta.* 2012;1819: 104–119.
33. Howell SH. Endoplasmic reticulum stress responses in plants. *Ann. Rev. Plant Biol.* 2013;64: 477–499.
34. Srivastava R, Deng Y, Shah S, Rao AG, Howell SH. BINDING PROTEIN is a master regulator of the endoplasmic reticulum stress sensor/transducer bZIP28 in *Arabidopsis*. *Plant Cell.* 2013;25:1416–1429.
35. Fragkostefanakis S, Mesihovic A, Hu Y, Schleiff E. Unfolded protein response in pollen development and heat stress tolerance. *Plant Reprod.* 2016;29: 81–91.
36. Deng Y, Srivastava R, Quilichini TD, Dong H, Bao, Y, Horner HT, Howell SH. IRE1, a component of the unfolded protein response signaling pathway, protects pollen development in *Arabidopsis* from heat stress. *Plant J.* 2016;88:193–204.
37. Baxter A, Mittler R, Suzuki N. ROS as key players in plant stress signalling. *J Exp Bot.* 2014;65:1229–1240.
38. Mittler R, Vanderauwera S, Gollery M, Van Breusegem F. Reactive oxygen gene network of plants. *Trends Plant Sci.* 2004;9:490–498.
39. Driedonks N, Xu J, Peters JL, Park S, Rieu I. Multi-level interactions between heat shock factors, heat shock proteins, and the redox system regulate acclimation to heat. *Front Plant Sci.* 2015;6: 999.
40. Muller F, Rieu I. Acclimation to high temperature during pollen development. *Plant Reprod.* 2016; 29:107–118
41. Parish RW, Phan HA, Iacuone S, Li SF. Tapetal development and abiotic stress: a centre of vulnerability. *Funct Plant Biol.* 2012;39:553–559.
42. Ma, H. Molecular genetic analyses of microsporogenesis and microgametogenesis in flowering plants. *Ann Rev Plant Biol.* 2005;56:393–434.
43. Scott RJ, Spielman M, Dickinson HG. Stamen structure and function. *The Plant Cell.* 2004;16:46–60.
44. Ariizumi T, Toriyama K. Genetic regulation of sporopollenin synthesis and pollen exine development. *Ann Rev Plant Biol.* 2011;62:437–460.

Comment [D63]: Write italic

Comment [D64]: Write italic

45. Parish RW, Li SF. Death of a tapetum: A programme of developmental altruism. *Plant Sci.*2010;178: 73–89.
46. Oshino T, Miura S, Kikuchi S, Hamada K, Yano K, Watanabe M, et al. Auxin depletion in barley plants under high-temperature conditions represses DNA proliferation in organelles and nuclei via transcriptional alterations. *Plant Cell Environ.* 2011;34:284–290.
47. Datta R, Chamusco KC, Chourey PS. Starch biosynthesis during pollen maturation is associated with altered patterns of gene expression in maize. *Plant Physiology.*2002;130:1645–1656.
48. Speranza A, Calzoni GL, Pacini E. Occurrence of mono- or disaccharides and polysaccharide reserves in mature pollen grains. *Sexual Plant Reproduction.*1997;10:110–115.
49. Saini HS. Effects of water stress on male gametophyte development in plants. *Sex Plant Reprod.* 1997;10:67–73.
50. Jain M, Vara Prasad PV, Boote JK, Hartwell Jr LA, Chourey SP. Effects of season-long high temperature growth conditions on sugar-to-starch metabolism in developing microspores of grain sorghum (*Sorghum bicolor* L. Moench). *Planta.*2007;227:67–79.
51. Sato S, Kamiyama M, Iwata T, Makita N, Furukawa H, Ikeda H. Moderate increase of mean daily temperature adversely affects fruit set of *Lycopersicon esculentum* by disrupting specific physiological processes in male reproductive development. *Ann Bot.*2006;97:731–738.
52. Oliver SN, van Dongen JT, Alfred SC, Mamun EA, Zhao XC, Saini HS, et al. Cold-induced repression of the rice anther-specific cell wall invertase gene OSINV4 is correlated with sucrose accumulation and pollen sterility. *Plant Cell Environ.* 2005;28:1534–1551.
53. Sato S, Peet MM, Thomas JF. Determining critical pre- and post-anthesis periods and physiological processes in *Lycopersicon esculentum* Mill. exposed to moderately elevated temperatures. *J Exp Bot.* 2002; 53:1187–1195.
54. Razzaq MK, Rauf S, Shahzad M, Ashraf I, Shah F. Genetic analysis of pollen viability: an indicator of heat stress in sunflower (*Helianthus annuus* L.). *Int J Innov Appr Agric Res.*2017;1(1) 40-508.
55. Satish D, Ravikumar RL. Standardization of in vitro pollen germination media in selected varieties of cotton and tomato. *Karnataka J Agric Sci.* 2010;23:317-319.
56. Patel RG, Mankad AU. In Vitro Pollen Germination-A Rev. *IJSR.*2014; 358: (5).
57. Chatterjee R, Sarkar S, Rao GN. Improved media for in vitro pollen germination of some species of Apocynaceae. *Int J Environ.*2014; 3:146-153.
58. Gaaliche B, Majdoub A, Trad M, Mars M. Assessment of pollen viability, germination, and tube growth in eight Tunisian caprifig (*Ficus carica* L.) cultivars. *ISRN Agron.*2013.
59. Razzaq MK, Rauf S, Akhtar N. Effect of various chemical hybridizing agents on sunflower (*Helianthus annuus*) seed production. *Seed Technol.* 2015;1:23-31.
60. Frescura VD, Laughinghouse HD, Cantodorow TS. Pollen viability of *Polygala paniculata* L (Polygalaceae) using different staining methods. *Bio. Cell.* 2012;36:143-145.
61. Radicevic S, Nikolic D, Cerovic R, Dordevic M. In vitro pollen germination and pollen grain morphology in some sweet cherry (*Prunus avium* L.) cultivars. *Rom Biotechnol Lett.* 2013;18(3):8341-8349.
62. Sulusoglu M, Cavusoglu A. In vitro Pollen Viability and Pollen Germination in Cherry Laurel (*Prunus laurocerasus* L.). *Sci World J*2014.ID: 657123.
63. Razzaq MK, Rauf S, Khurshid M, Iqbal S, Bhat JA, Farzand, A, Riaz, A. et al. Pollen Viability an Index of Abiotic Stresses Tolerance and Methods for the Improved Pollen Viability. *Pak J Agri Res.* 2019;32:(4) p:609.
64. Guru T, Kunta RT, Rao PR. Improving pollen viability and stigma receptivity by boron application as a physiological approach for high yields in rice. *Bioscan.* 2016;1:79-83.
65. Feng B, Zhang C, Chen T, Zhang X, Tao L, Fu G. Salicylic acid reverses pollen abortion of rice caused by heat stress. *BMC Plant Biol.*2018;18:245.
66. Lau TC, Stephenson AG. Effects of soil phosphorous on pollen production, pollen size, pollen phosphorous contents and the ability to sire seeds in *Cucurbita pepo* (Cucurbitaceae). *Sex Plant Reprod.*1994;7:215-220.

67. SadatS, Saeid KA, Bihanta MR, Torabi S, Salekdeh SGH, AyenehGAL. Marker assisted selection for heat tolerance in bread wheat. *J World Appl Sci.*2013;21:1181–1189.
68. XiaoYH, Pan Y, Luo LH, Deng HB, Zhang GL, Tang WB, Chen LY. Quantitative trait loci associated with pollen fertility under high temperature stress at flowering stage in rice (*Oryza sativa*). *Rice Sci.*2011;18:204–209.
69. KardiveIP. Molecular breeding in vegetable crops-challenges and opportunities. In :Proceeding of 2nd National Workshop on Marker-Assisted Selection for Crop Improvement, ICRISAT, Patancheru, India, 27–29 October; 2010.
70. LinkH, Yeh WL, Chen HM, LoHF. Quantitative trait loci influencing fruit-related characteristics of tomato grown in high-temperature conditions. *Euphytica.*2010;174:119–135.
71. YeC, ArgayosoMA, RedoñaED, Sierra SN, Laza MA, Dilla C et al. Mapping qtl for heat tolerance at flowering stage in rice using snp markers. *Plant Breed.*2012;131: 33–41.
72. Lucas MR, Ehlers JD, Huynh BL, Diop NN, Roberts PA, Close TJ. Markers for breeding heat-tolerant cowpea. *Mol. Breed.*2012; 31:529–536.
73. Firon N, Shaked R, PeetMM, Phari DM, Zamski E, Rosenfeld K, Althan L, Pressman NE. Pollen grains of heat tolerant tomato cultivars retain higher carbohydrate concentration under heat stress conditions. *SciHortic.*2006;109:212–217.
74. Arbona V, Manzi M, Ollas C, Gomez-Cadena A. Metabolomics as a tool to investigate abiotic stress tolerance in plants. *Int J Mol Sci.*2013;14:4885–4911.
75. PaupièreMJ, van HeusdenAW, Bovy AG. The metabolic basis of pollen thermo-tolerance: Perspectives for breeding. *Metabolites.* 2014;4:889-920. doi:10.3390/metabo4040889
76. ColbertT, Till BJ, Tompa R, Reynolds S, SteineMN, Yeung AT, et al. High-throughput screening for induced point mutations. *Plant Physiol.*2001;126:480–484.