

Effect of Concentration of Starter Culture and Incubation Time on Sensory Attributes of RTE Functional Greek Yoghurt

ABSTRACT

Aims: The aim of the current study was to optimize starter culture concentration and incubation time for the preparation of Ready-To- Eat functional Greek yoghurt

Study design: Optimization of culture concentration and incubation time for RTE functional Greek yoghurt

Place and Duration of Study: Sample: Department of Medicine (Medical Unit IV) and Department of Radiology, Services Institute of Medical Sciences (SIMS), Services Hospital Lahore, between June 2009 and July 2010.

Department of Dairy Technology, Dairy Science College, Hebbal, Bengaluru, Karnataka Veterinary, Animal and Fisheries Sciences University, Bidar, Karnataka, India. Between June 2022 and April 2024

Methodology:

Ready-To-eat functional Greek yoghurt was prepared by different levels of starter culture concentration and incubation time period and optimized based on sensory analysis using 9-point hedonic scale

Results: The addition of various levels of yoghurt culture (0.15, 0.20, 0.25, and 0.30%), the yoghurt was evaluated by sensory characteristics. According to trial results, 0.20% yoghurt culture scored higher than the control sample (7.14, 7.16, 7.15, and 7.16) in each category for colour and appearance, body and texture, flavour, and overall acceptability (8.06, 8.44, 8.60, and 8.65 out of 9.00). Effect of varying incubation period was analyzed after inoculation of yoghurt culture for different incubation hours (5, 6, 7 and 8 h). While judging sensory attributes for different parameters such as colour and appearance, body and texture, flavour, and overall acceptability, experimental sample at 6 hours of incubation scored maximum (8.27, 8.55, 8.72 and 8.67) as opposed to the control (7.15, 7.16, 7.17 and 7.16) respectively.

Conclusion: The optimized levels of starter culture concentration and incubation time improved the overall sensory quality of the Ready-To-Eat Greek yoghurt

Keywords: Starter culture, incubation time, Greek yoghurt, Sensory attributes

1. INTRODUCTION

Yoghurt has ability to introduce probiotic microorganisms into human intestine, as well as the high nutritional value of yoghurt constitute growing importance of Greek yoghurt as a potent functional food in food industry. Beyond their nutritional worth, fermented foods have been

shown to provide health benefits. Among the popular exotic fermented milk products, the consumption of yoghurt has surged rapidly in recent decade due to its remarkable nutritional as well as therapeutic properties [1].

Greek yogurt is traditionally obtained by straining normal yogurt, which gives it a thicker texture. It is a concentrated fermented milk as defined by Codex Alimentarius (2003)[2], which specifies that the protein content must be at least 5.6 g per 100 g. During the draining process, the aqueous part of the milk – the whey or lactoserum – is removed, which explains the low lactose content of Greek yogurt.

Greek yoghurt is a highly nutritious fermented product characterized by its higher protein, low sodium and low carbohydrate contents, and has gained immense popularity compared to traditional yoghurt. It has a unique semi-solid texture and rich mouthfeel due to the particular process during manufacturing. Moreover, Greek yogurt has a creamy white color, a soft and smooth body, good spreadability and the product will be slightly acidic. Since it is a concentrated form of yogurt, it possesses sensory attributes that are different from regular yogurt as well as superior nutritional properties make it a good choice to be included in daily diet [3].

The production of yogurt is a basic ecology that depends on the interactions of two bacteria, *Lactobacillus delbrueckii* subsp. *bulgaricus* and *Streptococcus thermophilus*. The ecological term proto-cooperation describes the relationship between *S. thermophilus* and *Lb. bulgaricus* in a yogurt starter culture. Lactic acid bacteria (LAB) are picky microorganisms, and lack of vital nutrients in milk that frequently limits their ability to develop. Thus, the cooperation of *Lactobacillus bulgaricus* and *Streptococcus thermophilus* is often critical to the success of milk fermentation. Proto-cooperation is the term for this indirect positive connection that occurs when both bacteria are able to develop independently in milk. This positive relationship often has a beneficial effect on bacterial growth and on the production on lactic acid and aroma compounds like acetaldehyde [4].

So, in order to maximize process efficiency and consumer acceptance, the current study was done to investigate the impact of culture concentration and incubation time on various characteristics of Greek yoghurt produced.

2. MATERIAL AND METHODS

For the preparation of RTE functional Greek yoghurt, Fresh Cow milk was procured from Students Experimental Dairy Plant (SEDP) of Dairy Science College, Hebbal, Bengaluru and Freeze dried starter culture was procured from Delvo DSL pvt Ltd, Netherlands. The milk was heated to 90°C/5 minutes and cooled to 42°C followed by addition of freeze-dried DVS Yoghurt culture of selected microflora (*Streptococcus thermophilus* and *Lactobacillus bulgaricus*) was added at concentrations of 0.15, 0.20, 0.25, and 0.30 percent. Then further incubation at 42°C for 5, 6, 7 and 8 hours and then subjected for de-wheyng.

The panel of trained judges that was chosen assessed the generated study sample's sensory attributes, such as colour and appearance, body and texture, flavour, and overall acceptability, using a 9-point hedonic scale. Statistical analysis was conducted using the output with the highest score as per Peryam and Pilgrim, (1957) [5]. Using R software (R. version 4.0.3), the data collected for the research investigations was examined to evaluate the significant or non-significant effects of various treatments and trials obtained for the current study. The mean and critical difference was calculated.

3. RESULTS AND DISCUSSION

3.1. Effect of concentration of starter culture on sensory attributes of RTE functional Greek yoghurt.

The colour and appearance scores of the control and other treated samples showed a significant difference ($P < 0.05$). Highest score of 8.06 was obtained for RTE functional Greek yoghurt made with 0.20% yoghurt culture and lowest score of 7.03 was obtained for Greek yoghurt made with 0.30% yoghurt culture respectively. The colour and appearance may have got affected in sample with 0.20% culture addition as the surface was smooth and uniform. The study conducted by Gamage et al. [6] observed the similar effects in drinking yoghurt developed from freeze dried yoghurt culture at 0.2% culture level (4.4, 3.9, 4.0 & 4.0 out of 5 for colour, flavour, texture and acceptability, respectively) against 0.3 % and 0.4%.

Body and texture of yoghurt was significantly affected by yoghurt culture concentration ($P < 0.05$) as the Lactic Acid Bacteria has ability to produce exopolysaccharides and bioactive peptide which could act as the stabilizer, thickener and emulsifier like other polysaccharides. As 0.20% yoghurt culture forming firm gel-like body distributed throughout the product it received the highest sensory score of 8.44, whereas 0.15 % gave weak body. 0.3% yoghurt culture addition resulted in non-uniform body due to free whey separation and obtained least score (7.00) for sensory. Highest texture scores were recorded for yoghurt samples prepared with 2.00 and 2.50 % sub-cultured inoculum in the research findings of Anjum et al. [7].

Flavour score for yoghurt culture addition was highest (8.60) for 0.20% with natural yoghurt flavour with significant differences ($P < 0.05$) whereas 0.30% culture addition resulted acidic flavour scored lowest sensory score (7.10) as starter culture has symbiotic interaction which responsible for aroma compound acetaldehyde. The judges recorded similar level of scores for yoghurts for flavour and taste for the yoghurt culture after sub-culturing concentration of 1% showed higher acceptability as opposed to 0.9% and 1.1% culture level for buffalo milk yoghurt from the research of Iftikhar et al.[8].

Overall acceptability rating was significantly highest ($P < 0.05$) for yoghurt with 0.2% culture addition compared to control and other experimental samples and scored 8.65. The least scored sample was 0.3% yoghurt culture addition with a score of 7.03. The highest overall acceptability was recorded in drinking yoghurt developed from regular yoghurt culture at 0.2% freeze dried yoghurt culture level out of 0.2%, 0.3% and 0.4% culture level in the research study by Gamage et al., 2016. In 2020 Dabija et al. [9] studied the influence of inoculation doses for yoghurt from cow milk on rheological properties and observed that the optimum freeze-dried culture dose is 0.2 %, while at 0.3 % the rheological parameters decrease. The decrease of the rheological parameters at high inoculation doses is because of the powerful proteolysis activity of *Lactobacillus delbrueckii* subsp. *bulgaricus* and *Streptococcus thermophilus*.

The effect of different levels of yoghurt culture on sensory characteristics of RTE functional Greek yoghurt based on 9-point hedonic scale has been mentioned in Table.1 and Fig.1

Table 1: Effect of yoghurt culture on the sensory characteristics of RTE functional Greek yoghurt.

Culture	Colour	Body	Flavour	Overall
---------	--------	------	---------	---------

addition (%)	andappearance	andtexture		acceptability
Control(0.1%)	7.14 ^b	7.16 ^{bc}	7.15 ^b	7.16 ^{cd}
0.15	7.16 ^b	7.57 ^{bc}	7.82 ^{ab}	8.21 ^b
0.2	8.06 ^a	8.44 ^a	8.60 ^a	8.65 ^a
0.25	7.23 ^b	7.60 ^b	7.60 ^b	7.40 ^c
0.30	7.03 ^b	7.00 ^c	7.10 ^b	7.03 ^d
CD (P=0.05)	0.26	0.40	0.61	0.19

*Note:

Inoculation of freeze dried DVS starter culture at 0.1% for control to get 107-1010 cfu/ml of bacterial cell concentration and incubated at 42 °C for 4 hours

All the values are average of three trials

Similar superscripts indicate non - significance at the corresponding critical difference

Sensory analysis – 9-point hedonic scale




Fig. 1. Effect of yoghurt culture on the sensory characteristics of RTE functional Greek yoghurt.

3.2. Effect of incubation time on sensory attributes of RTE functional Greek yoghurt.

There was a substantial difference ($P < 0.05$) between the control and other treated samples' colour and appearance evaluations. The Greek yoghurt that made with 6 hours of incubation had the highest score of 8.27 due to smooth and even surface, while the 4 hours of incubation resulted in getting lowest score (7.15) of control. Eze et al. [10] got the highest score for the set yoghurt made of 5 hours of incubation time (7.35, 6.8, 6.9 & 7.35 for colour, flavour, texture and acceptability, respectively) against 3 and 4 hours.

There is a significant difference ($P < 0.05$) between the body and texture scores of the control and experimental samples. The sample's body and texture ratings raised to 8.55 with 6 hours of incubation and lowest score of 7.00 for 8 hours of incubation. This may be due to longer incubation time tends to weaken the structure of yoghurt due to over-fermentation. The findings are correlated Aman et al., (2021) got soft texture for yoghurt after 8 hours of incubation time.

The research revealed a significant difference ($P < 0.05$) in the flavour ratings between the experimental and control samples of Greek yoghurt. The flavour score increased to 8.72 with 6 hours of incubation time for yoghurt due to fully developed flavour and sample with 8 hours of incubation had the lowest score of 7.16. Shorter incubation time leads to milder and incomplete flavour while incubating for longer time can cause high-acid flavour. The results aligned with the research conducted by Eze et al., (2021), which indicated that scores intensified when set yoghurt made with 5 hours of incubation time.

When compared to the control and other experimental samples, the yoghurt with 6 hours of incubation time had the greatest overall acceptance rating ($P < 0.05$), scoring 8.67. With a score of 7.05, the sample with the lowest score was the experimental sample with 8 hours of incubation time. In Aman et al.'s [11] research, the yoghurt that was developed with 8 hours of incubation time scored more for overall acceptability as opposed to 6 and 10 hours with different processing condition.

Table 2 and Fig. 2 discuss the impact of varying incubation period on the sensory attributes of Greek yoghurt.

Table 2. Effect of incubation time on sensory attributes of RTE functional Greek yoghurt

Incubation hours at 42°C	Colour and appearance	Body and texture	Flavour	Overall acceptability
Control (4 Hr)	7.15 ^c	7.16 ^c	7.17 ^c	7.16 ^c
5	7.70 ^b	7.47 ^b	8.10 ^b	8.07 ^b
6	8.27 ^a	8.55 ^a	8.72 ^a	8.67 ^a
7	7.40 ^{bc}	7.32 ^{bc}	7.30 ^c	7.33 ^c
8	7.20 ^c	7.00 ^c	7.16 ^c	7.05 ^c
CD (P=0.05)	0.29	0.24	0.31	0.36

*Note:

All the values are average of three trials

Similar superscripts indicate non - significance at the corresponding critical difference




Fig. 2: Effect of incubation period on the sensory characteristics of Greek yoghurt.

4. CONCLUSION

The addition of different levels of yoghurt culture the RTE functional Greek yoghurt was evaluated by sensory characteristics. According to research trial results, 0.2% yoghurt culture scored higher than the control for colour and appearance, body and texture, flavour, and overall acceptability. Effect of varying incubation period was analysed after inoculation of yoghurt culture for different incubation hours. While subjected for judging sensory attributes for different parameters such as colour and appearance, body and texture, flavour, and overall acceptability, experimental sample at 6 hours of incubation scored maximum as opposed to the control. The metabolic processes of the starter cultures are the source of some of the beneficial effects such as rise in acidity and typical flavour and production of exopolysaccharide which helps in suitable body formation.

REFERENCES

1. Vijay L, Karthikeyan N, Kumaresan G and Kathirvelan C. Sensory properties of functional Greek yoghurt incorporated with *Moringa oleifera* leaf powder. *J. Pharm. Innov.* 2022; (4): 251-254.
2. Codex Alimentarius. Codex Standard for Fermented Milks. Codex stan.2003;243-2013.
3. Mani F. and Hannachi C. Quality mentoring of Greek yoghurt fortified with pomegranate juice and arils (*Punica granatum L.*) during storage. *J. New Sci.* 2017; 44 (5): 2430-2437.

4. Mihail A, Kostov G, Simova E, Beshkova D, Koprinkova-Hristova P. Proto-cooperation factors in yoghurt starter cultures. *Revue de Génie Industriel*. 2009; 3: 4-12.
5. Peryam DR and Pilgrim FJ. Hedonic scale method of measuring food preferences. *Food Technol*. 1957; 15 (6): 9-14.
6. Gamage, G G A P, Adikari A M J B, Nayananjali W, Prasanna P, Jayawardena N W I A and Wathsala R H G R. Physicochemical, microbiological and sensory properties of probiotic drinking yoghurt developed with goat milk. *Int. J. Sci. Res. Publ.* 2016; 6(6): 203-208.
7. Anjum R R, Zahoor T and Akhtar S. Comparative study of yoghurt prepared by using local isolated and commercial imported starter culture. *J. Res. Sci.* 2007;18 (1): 35-41.
8. Iftikhar M A, Pasha T N, Inayat S, Javed K and Ullah R. Effect of concentration of culture on incubation time, storage and sensory characteristics of set yoghurt. *Pak. J. Agric. Sci.* 2023; 60 (1): 149-161.
9. Dabija A, Oroian M, Codină GG and Rusu L. Assessment the influence of the main technological factors on yogurt quality. *Scientific Study & Research. Chemistry & Chemical Engineering, Biotechnology, Food Industry.* 2020; 21(1): 83-94.
10. Eze CM, Aremu K O, Alamu E O and Okonkwo T M. Impact of type and level of stabilizers and fermentation period on the nutritional, microbiological, and sensory properties of short-set Yoghurt. *Food Sci. Nutr.* 2021; 9 (10): 5477-5492.
11. Aman, Santos R, Rini P, Vina W, Anugrah A and Tiara S. The effect of starter concentration and incubation time on yogurt characteristics. *AIP Conference Proceedings*. 2021; 2353 (1): 1-6