

Impact of integrated nitrogen management practices on nutrient content and their uptake by transplanted rice crop (*Oryza sativa* L.)

ABSTRACT

Aim: Nitrogen is very crucial for production of rice and recently the uses of chemical fertilizers are increased for rice production, which are ruining our soils and environment. So, to evaluate the effect of integrated nitrogen management practices on nutrient content and their uptake by transplanted rice crop this study was conducted.

Study design: Randomized block design.

Place and duration of study: One year field experiment at Research farm, School of Agriculture, Abhilashi University, Chail Chowk, Mandi, (H.P.).

Methodology: The field study was conducted with eight treatments and three replications. The different treatments combinations were T₁ (absolute control), T₂ [100% RDN through Chemical Fertilizer (CF)], T₃ (75% RDN through CF + 25% N through FYM), T₄ (75% RDN through CF + 25% N through poultry manure), T₅ (75% RDN through CF + 25% N through vermicompost), T₆ (50% RDN through CF + 50% N through FYM), T₇ (50% RDN through CF + 50% N through poultry manure), T₈ (50% RDN through CF + 50% N through vermicompost).

Results: The study of results revealed that the nitrogen, phosphorus and potassium content in grains and straw of rice crop showed non-significant differences with the application of nitrogen through various treatments, while the maximum contents of these nutrients were higher in treatment T₂. However, the significantly highest uptake of

nitrogen, phosphorus and potassium by grains and straw and total uptake of these nutrients by rice was noted in treatment T₂ which was statistically ($P = .05$) at par with treatment T₄ and T₅. Whereas, the minimum content in grains and straw and uptake of these nutrients by grains and straw along with their total uptake was found under control treatment.

Conclusion: This study suggests that the use of integrated nitrogen management enhances the nitrogen, phosphorus and potassium content and their uptake by rice crop.

Keyword: Rice, chemical fertilizer, poultry manure, vermicompost, nutrient content and nutrient uptake.

1. INTRODUCTION

“Rice (*Oryza sativa* L.) is the most important crop in the developing world and is the staple food of over half the world’s population. For 40% of the world’s population, it is their primary source of calories” (Virdia and Mehta, 2009) [1]. “Almost 90% of the total rice is produced and consumed in Asia. Half of the world’s population depends primarily on rice, which is farmed in more than hundred countries” (Fukagawa et al. 2019) [2]. “Rice is an excellent food and an excellent source of carbohydrates and energy. It is a high calorie food which contains carbohydrates, protein, fat, cellulose and ash. In Asia, more than two billion people are getting 60-70% of their energy requirement from rice and its derived products” (Tomar et al. 2018) [3,24,25,26]. “Rice is cultivated in approximately 120 countries; China grows 214 million tons and

India (about 173 million tons)” (Asenso et al. 2021) [4]. “China was the world leader in rice production in 2019-20, producing 146.73 million metric tons, followed by India with 115.00 million metric tons” (Anonymous, 2020) [5]. “In India, rice occupies an area of 45.77 million hectares with production of 124.37 million tons with an average productivity of 2717 kg ha⁻¹” (Anonymous, 2021) [6].

“Nitrogen (N) is the most essential and limiting nutrient element required in large amounts for rice production and provision of adequate supply of N throughout the growing period is essential for realizing potential yields” (Surekha et al. 2016) [7]. Nitrogen promotes rapid plant growth and improves grain yield and quality through higher tillering, leaf area development, grain formation, grain filling and protein synthesis (Reddy et al. 2019) [8] and plays a crucial role in enhancing the yield. “Nitrogen plays a vital role in plant tissues and is a constituent of protein, enzyme, hormone, vitamins, alkaloids and chlorophyll etc. Nitrogen is highly mobile both in the plants and soil. Within a certain range, increased nitrogen fertilizer increase net photosynthesis rate. Nevertheless, too much nitrogen fertilizer causes the photosynthetic capacity decreases. Increasing nitrogen fertilization can enhance the nutritional quality and processing quality of rice” (Li et al. 2019) [9].

“Nitrogenous fertilizers are one of the main factors responsible for contributing to increase in rice yield in India. In rice production, farmers apply large amounts of N fertilizer to maximize yield, but only 20-50% of N is taken up by the crop” (Chivenge et al. 2021) [10]. “The resulting loss of the applied N, which is a mobile nutrient leads to increased water and land pollution and greenhouse gas (GHG) emissions” (Zhang et al. 2013) [11]. “Continuous use of chemical fertilizers alone causes soil organic matter degradation, soil acidity and environmental pollution” (Shipra et al. 2019) [12]. The increasing fertility of soil and crop productivity through use of chemical or synthetic fertilizers has often negatively affected biogeochemical cycles. Also, usage of fertilizer cause leaching and run-off of nutrients, especially nitrogen and phosphorus results in degradation of environment. Nitrogen (N) can be lost from the soil by leaching, de-nitrification and volatilization. Practice like deep placement of urea super granule (USG), soil incorporation, use of slow release N fertilizer etc., that minimize N concentration in flood water can reduce its loss through runoff.

“Inorganic fertilizer has been shown to degrade soil and improve production rapidly when used during cropping, but not over a long period of time” (Satynarayana et al. 2002) [13]. “The integration of organic manure with inorganic fertilizers is essential to achieving sustained production of rice and maintains the soil fertility for a longer period” (Gill and Walia, 2014) [14]. “Numerous research findings suggest that using nitrogen fertilizer in combination with organic sources can accelerate the process of organic N mineralization and immobilization, hence, lowering nitrogen losses” (Wang et al. 2023) [15]. “Manures improve soil structure, water holding capacity of soil, organic matter contents as well as microbial populations. Integrated use of organics especially vermicompost, poultry manure, farmyard manure, green manure, neem cake and bio-fertilizers as sources of nitrogen application with inorganic fertilizers improve the growth, yield and yield attributing traits in rice. Poultry manure contains major, secondary and micro nutrients that can support crop production and enhance the physical and chemical properties of soil. It improves soil retention and uptake of plant nutrients” (Mohammed et al. (2010) [16]. “Vermicompost is rich in plant nutrients like- nitrogen, phosphorus, potassium, calcium, vitamins, natural phyto-regulators and micro flora in balanced form that helps in re-establishment of the natural fertility of the soil” (Arancon and Edwards, 2009) [17]. “Optimal crop yields and long term soil productivity may be maintained with the combined application of organic manures and inorganic fertilizers” (Puli et al. 2016) [18]. “Integration of nutrients from organic manures and chemical fertilizers is necessary to maintain soil fertility, amount of nutrients in soil, production sustainability and good crop yield” (Shankar et al. 2020) [19].

2. MATERIAL AND METHODS

2.1 Details of the experiment site and plant material

This research was carried out at the research farm of the School of Agriculture, Abhilashi University, Chail Chowk, Mandi (H.P.) during the *Kharif* season of 2023. The experimental farm is situated at 30° 32' N latitude and 74° 53' E longitudes, with an elevation of 1391 m above mean sea level. The soil of the experimental field was slightly acidic in reaction, medium in organic carbon, low in available nitrogen and medium in available phosphorus and potassium. The pH (5.5) of the experimental soil was slightly acidic in reaction with an electrical conductivity of 0.011 dS m⁻¹, medium in organic carbon (0.70%), low in available nitrogen (218.37 kg ha⁻¹), medium in available phosphorus (17.89 kg ha⁻¹) and potassium (226.15 kg ha⁻¹). For the determination of the content of nitrogen, phosphorus and potassium in grains and straw of rice crop, the sample of grains and straw of the rice were collected after harvest of the crop from the each treated plots, separately.

2.2 Design of the experiment and cultural practices

The spacing for the tested variety (Hybrid paddy super-120) was 20 × 10 cm row to row and plant to plant. The experiment was laid out in a randomized block design (RBD) with eight treatments and three replications. The treatments include:

List 1: List of treatments

S.N.	Treatments
T ₁	Absolute control
T ₂	100% RDN (Recommended dose of nitrogen) through CF (chemical fertilizer)
T ₃	75% RDN through CF + 25% N through FYM
T ₄	75% RDN through CF + 25% N through poultry manure
T ₅	75% RDN through CF + 25% N through vermicompost
T ₆	50% RDN through CF + 50% N through FYM
T ₇	50% RDN through CF + 50% N through poultry manure
T ₈	50% RDN through CF + 50% N through vermicompost

Recommended doses of N, P and K were applied in the form of urea, diammonium phosphate (DAP) and muriate of potash (MOP). Well decomposed farm yard manure, poultry manure and vermicompost were used as organic sources for nitrogen. Urea was the source of nitrogen, and it was administered in three separate doses during the transplanting, tillering and panicle initiation stages. Equal doses of potassium and phosphorus (50 kg ha⁻¹) were applied to each treatment as basal application in the form of muriate of potash and diammonium phosphate (DAP), respectively. After harvest, plant samples from each treatments were gathered, cleaned and then allowed to dry in the shade at room temperature. A mixer was used to finely powder the samples after they had been shade-dried and oven-dried for 24 to 48 hours at 60 ± 2°C. N, P and K content as well as rice crop uptake were examined using the finely powdered plant samples. The modified Kjeldahl digestion and distillation method, which was used to estimate the amount of nitrogen in the plant sample described by (Jackson, 1973) [20]. Using the vanadomolybdate phosphoric yellow color method, the phosphorus content of the plant was ascertained, and the phosphorus content of the plant sample was estimated as per (Jackson, 1973) [20] instructions. A flame photometer was used to evaluate the potassium content of plants (Jackson, 1973) [20]. The N, P and K

absorption (kg ha^{-1}) in each treatment was calculated by multiplying the N, P and K content (%) by the grain and straw yields (q ha^{-1}) of transplanted rice crop at harvest.

2.3 Analysis of data with statistical method

The collected data were analyzed statistically following the one factor analysis of variance (ANOVA) technique and the mean differences were adjusted with operational statistics (OPSTAT) software.

3. RESULTS AND DISCUSSION

3.1 Nitrogen content (%) and uptake (kg ha^{-1})

The nitrogen (N) content and their uptake by grains and straw of transplanted rice is displayed in (Table-1) and predicted in the (Fig.-1). The study showed no significant differences of nitrogen concentration present in the grains and straw of the transplanted rice crop by various nitrogen management treatments. However, the maximum nitrogen content in grains and straw of transplanted rice crop was noticed in treatment of T_2 (100% RDN through CF) and minimum in treatment T_1 (absolute control).

The application N by organic manure and nitrogenous chemical fertilizer had a substantial impact on the uptake of N by grains and straw as well as the total N uptake. During the study, the highest nitrogen uptake by grains and straw of transplanted rice crop was recorded under treatment T_2 (100% RDN through CF), which was statistically ($P = .05$) at par with treatment T_4 (75% RDN through CF + 25% N through poultry manure) and T_5 (75% RDN through CF + 25% N through vermicompost). Though, treatment T_1 (absolute control) showed the minimum amount of N uptake by grains and straw of transplanted rice crop. The treatment T_2 also recorded the total uptake of nitrogen and was it was on par with treatments T_4 and T_5 and the minimum total nitrogen uptake was found under treatment T_1 .

Table 1. Effect of integrated nitrogen management on nitrogen content and their uptake by grain and straw of transplanted rice crop

S.N.	Treatments	Nitrogen content (%)		Nitrogen uptake (kg ha^{-1})		
		Grain	Straw	Grain	Straw	Total
T ₁	Absolute control	1.31	0.40	37.74	15.87	53.61
T ₂	100% RDN through CF (Chemical Fertilizer)	1.39	0.44	69.12	25.15	93.27
T ₃	75% RDN through CF + 25% N through FYM	1.36	0.42	61.30	21.16	82.46
T ₄	75% RDN through CF + 25% N through poultry manure	1.38	0.43	66.78	23.07	89.85
T ₅	75% RDN through CF + 25% N through vermicompost	1.37	0.43	64.55	23.57	87.12
T ₆	50% RDN through CF + 50% N through FYM	1.34	0.41	53.56	19.11	72.67
T ₇	50% RDN through CF + 50% N through poultry manure	1.36	0.42	58.18	20.39	78.57
T ₈	50% RDN through CF + 50% N through vermicompost	1.35	0.42	55.42	20.01	75.43
	S.Em.±	0.04	0.01	1.89	0.61	2.51
	C.D. (P = .05)	NS	NS	5.78	1.88	7.70

The nitrogen uptake by rice grains and straw increased significantly with the combined application of organic manure and chemical fertilizers as compared to control. The reason for the increased uptake of N with incremental dosages could be that the plant absorbed the nutrients proportionately as the amount of

accessible nutrients in the soil solution increased. Integrated application of nutrients might enhance nitrogen uptake in rice by providing a controlled release of nutrients, ensuring sustained availability to rice crop. This might be due to only larger fertilizer doses are able to fill the absorption sites in the soil and increase the nutrient status. Organic manure nutrient release rate is usually very slow and high depends on microbial activity. Whereas, chemical fertilizer are highly concentrated with less leaching and quick release of N compared to organic manure can be attributed for higher uptake of nitrogen in grains and straw of transplanted rice crop. Similar results also reported by **Sahu and Chaubey (2020) [21]**.




Fig.1. Effect of integrated nitrogen management on nitrogen content and their uptake by grain and straw of transplanted rice crop

3.2 Phosphorus content (%) and uptake (kg ha⁻¹)

Along with their uptake by grains, straw and total uptake, the P content in grains and straw of the transplanted rice crop is provided in (Table- 2) and illustrated in (Fig.- 2). The study of data shows that the amount of P in the grains and straw of the transplanted rice crop did not vary significantly ($P = .05$) by the application of the various nitrogen management treatments. While, the treatment T₂= 100% RDN through CF had the highest P content in grains and straw of transplanted rice crop, however, the minimum P content in grain and straw of transplanted rice crop was reported under treatment T₁ (absolute control).

Nitrogen application through organic and inorganic sources had an adequate effect on the uptake of phosphorus by grains and straw. The study of the data shows that treatment T₂= (100% RDN through CF) recorded the maximum nitrogen uptake by grains and straw of transplanted rice crop and it was statistically ($P = .05$) on par with treatment T₄ (75% RDN through CF + 25% N through poultry manure) and T₅ (75% RDN through CF + 25% N through vermicompost). However, the minimum uptake of P by grains and straw of transplanted rice crop was recorded under treatment T₁ (absolute control). Likewise, uptake of P by grains and straw, the total P uptake by transplanted rice crop was highest under treatment T₂, which was statistically comparable with treatment T₄ and T₅. While, the lowest P uptake by transplanted rice crop were found in treatment T₁.

The integration of the chemical fertilizers with organic manure noted the higher content and uptake of phosphorus and was more effective than control treatment in promoting phosphorus uptake in rice crop.

Nitrogen might enhance the efficacy of phosphorus-absorbing mechanisms and encourages root growth, which in turn helps in uptake of phosphorus by transplanted rice crop. Phosphorus moves to the root surface through diffusion. However, the presence of the organic matters in the different organic manures increases the activity and population of the microbes into the soil, which might increase the absorption of phosphorus in grains and straw of transplanted rice crop. Eventually, integrated nitrogen application enhancing root development, soil characteristics, nutrient interaction and efficiency promotes rice to absorb the maximum amount of phosphorus available. Tiwari et al. (2020) [22] also noted the similar finding related to this study.

Table 2: Effect of integrated nitrogen management on phosphorus content and their uptake by grain and straw of transplanted rice crop

S.N.	Treatments	Phosphorus content (%)		Phosphorus uptake (kg ha ⁻¹)		
		Grain	Straw	Grain	Straw	Total
T ₁	Absolute control	0.21	0.100	6.05	3.97	10.02
T ₂	100% RDN through CF (Chemical Fertilizer)	0.24	0.108	11.94	5.93	17.87
T ₃	75% RDN through CF + 25% N through FYM	0.23	0.105	10.37	5.29	15.66
T ₄	75% RDN through CF + 25% N through poultry manure	0.24	0.106	11.61	5.69	17.30
T ₅	75% RDN through CF + 25% N through vermicompost	0.23	0.106	11.31	5.56	16.87
T ₆	50% RDN through CF + 50% N through FYM	0.21	0.102	8.39	4.76	13.15
T ₇	50% RDN through CF + 50% N through poultry manure	0.22	0.104	9.41	5.05	14.46
T ₈	50% RDN through CF + 50% N through vermicompost	0.22	0.103	9.03	4.91	13.94
	S.Em.±	0.01	0.003	0.28	0.16	0.49
	C.D. (P = .05)	NS	NS	0.86	0.50	1.51




Fig.2. Effect of integrated nitrogen management on phosphorus content and their uptake by grain and straw of transplanted rice crop

3.3 Potassium content (%) and uptake (kg ha⁻¹)

The K content and their uptake by grains and straw of transplanted rice are presented in (Table- 3) and depicted in (Fig.- 3). The analysis of the data of the research trial showed that there are no significant ($P = .05$) differences of the K content in the grains and straw of the transplanted rice crop. However, the highest K content in grains and straw of the transplanted rice crop was recorded under treatment T₂= 100% RDN through CF (chemical fertilizer), whereas, the lowest was observed to be in treatment T₁ (absolute control).

The uptake of potassium (K) by transplanted rice crop grains and straw was found to be significantly ($P = .05$) impacted by integrated nitrogen management. In this study, T₂ (100% RDN through CF) recorded the maximum K uptake by grains and straw of transplanted rice crop, which was statistically ($P = .05$) at par with treatment T₄ (75% RDN through CF + 25% N through poultry manure) and T₅ (75% RDN through CF + 25% N through vermicompost). While, the treatment T₁ (absolute control) noticed the lowest K uptake by grains and straw. The total uptake of K by transplanted rice crop was also recorded under treatment T₂, which was statistically ($P = .05$) on par with treatment T₄ and T₅. Even so, treatment T₁ (absolute control) had the minimum uptake of K by transplanted rice crop during the research.

Table 3: Effect of integrated potassium management on potassium content and their uptake by grain and straw of transplanted rice crop

S.N.	Treatments	Potassium content (%)		Potassium uptake (kg ha ⁻¹)		
		Grain	Straw	Grain	Straw	Total
T ₁	Absolute control	0.27	1.30	7.78	51.58	59.36
T ₂	100% RDN through CF (Chemical Fertilizer)	0.31	1.36	15.42	74.64	90.06
T ₃	75% RDN through CF + 25% N through FYM	0.29	1.34	13.07	67.50	80.57
T ₄	75% RDN through CF + 25% N through poultry manure	0.31	1.35	15.00	72.41	87.57
T ₅	75% RDN through CF + 25% N through vermicompost	0.30	1.34	14.14	70.32	87.41
T ₆	50% RDN through CF + 50% N through FYM	0.28	1.31	11.19	61.07	84.46
T ₇	50% RDN through CF + 50% N through poultry manure	0.29	1.33	12.41	64.56	72.26
T ₈	50% RDN through CF + 50% N through vermicompost	0.28	1.32	11.49	62.90	74.39
	S.Em.±	0.01	0.04	0.43	2.12	2.40
	C.D. (P = .05)	NS	NS	1.32	6.51	7.36

The application of organic manures and chemical fertilizer might have changed the physical, chemical and most importantly biological conditions of the soil and assisted in the transportation and absorption of nutrients from the soil. Nitrogen fertilization encourages root growth, which increases the surface area available for the absorption of nutrients, which include potassium. Rice plants with longer root systems are able to reach more soils, resulting in they get into interaction with more potassium ions. Increased nitrogen can help the movement of potassium from the roots to various areas of the plant facilitating its uptake and effective utilization. Adequate nitrogen and potassium supply are necessary for enzymatic activity and the metabolism of carbohydrates, increased photosynthesis activity which all might result in a higher uptake of potassium. The application of organic and inorganic sources of nutrients in combination remarkably improved potassium uptake by (promoting root development, ensuring balanced nutrient

uptake and transport and physico-chemical properties) in grains and straw of transplanted rice crop. Similar results related to this study were also reported by Sharma and Singh (2021) [23].




Fig.3. Effect of integrated nitrogen management on potassium content and their uptake by grain and straw of transplanted rice crop

4. CONCLUSION

In conclusion, this study shows that the application of the chemical fertilizers alone or integration of the chemical fertilizers with organic manures, to provide the nitrogen to rice crop has been found significant to attain the higher nutrients content and uptake (*viz.*-nitrogen, phosphorus and potassium) by rice crop. The content of the nitrogen, phosphorus and potassium were non-significant in grains and straw of rice, while, found maximum under 100% RDN by chemical fertilizers, however, uptake of these nutrients by grains, straw and total uptake by rice crop were also noted maximum with 100% RDN through chemical fertilizers and it was statistically at par with alteration of 75% RDF by chemical fertilizers with 25% poultry manure and 25% vermicompost, respectively. This study concluded that the alteration of the nitrogen supply can be done through chemical fertilizers with 25 or 50% organic manures for higher nutrients content and their uptake by rice crop. This is very important because the rice crop is most important cereal crop for most of the world's population, which can be produced with integration of inorganic fertilizers and organic manure and it has significance for maintaining the soil fertility and sustaining the environment by less use of the chemical fertilizers for the production of the rice crop.

ACKNOWLEDGEMENT

I would like to express my deepest gratitude to my major advisor (Dr. Jay Nath Patel) for their guidance, support and encouragement throughout the entire duration of this study. The course of this research has been greatly influenced by their knowledge and helpful guidance. I am also like to thank member of my research committee (Dr. Mohd Shah Alam) for their suggestions and helpful advice. The authors are thankful to Department of Agronomy, School of Agriculture, Abhilashi University, Chail Chowk, Mandi, H.P. for providing necessary laboratory facilities.

AUTHOR'S CONTRIBUTION

The first author of this research paper is M.Sc. Research Scholar of Agronomy Department, School of Agriculture, Abhilashi University, Chail Chowk, Mandi, (H.P.) India. The author conducted the field study, collected the all data and performed the laboratorial methods under the guidance of **Dr. Jay Nath Patel (major advisor of author)** who has supervised the author for every technical and laboratorial works during the experiment and writing the literature. He has also provided the necessary equipment's to the author to complete their research work. Dr. Mohd Shah Alam is member of advisory committee of author and helped author a lot.

Disclaimer (Artificial intelligence)

Option 1:

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

Option 2:

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc have been used during writing or editing of manuscripts. This explanation will include list the name, version, model, and source of the generative AI technology and as well as the all input prompts provided to a generative AI technology

Details of the AI usage are given below:

- 1.
- 2.
- 3.

REFERENCES

1. Virdia HM, Mehta HD. Integrated Nutrient Management in Transplanted Rice (*Oryza sativa* L.). Journal of Rice Research, Vol.2, No.2.pp. 2009;99-104.
2. **Fukagawa NK and Ziska LH. Rice: Importance for global nutrition. Journal of Nutritional Science and Vitaminology. 2019;65:82-83.**
3. Tomar R, Singh NB, Singh V, Kumar D. Effect of planting method and integrated nutrient management on growth parameters, yield and economic of rice. Journal of Pharmacognosy and Phytochemistry. 2018;7(2):520-527.
4. **Asenso E, Wang Z, Li J and Hu L. Puddling, direct seeding, mechanical transplanting for rice: effects on soil characteristics and productivity of rice, 2021.**
5. Anonymous. (USDA) World Agricultural Production. Foreign Agricultural Service Circular Series WAP 2020;1-20.

6. Anonymous. 4th Advance Estimates, Agricultural statistics at a glance, Department of Agriculture and Farmers Welfare Economics & Statistics Division. Government of India Ministry of Agriculture & Farmers Welfare. 2021.
7. Surekha KR, Kumar M, Nagendra N, Sailaja T, Satyanarayana V. 4R nitrogen management for sustained rice production better crops. *South Asia*, pp. 2016;18-19.
8. Reddy P, Rao BB, Surekha K, Hussain S. Transplanted rice as influenced by different enriched nitrogen sources - An economic appraisal. *International Journal of Current Microbiology and Applied Sciences*. 2019;8(06):3229-3231.
9. Li S, Pu S, Deng F, Wang L, Hu H, Liao S, Li W and Ren W. Influence of optimized management on the quality of medium hybrid rice under different ecological conditions. *Chinese Journal of Eco-Agriculture*. 2019;27(7):1042-1052.
10. Chivenge P, Sharma S, Bunquin MA and Hellin J. Improving nitrogen use efficiency – A key for sustainable rice production systems. *Frontiers in Sustainable Food Systems*. 2021;5:737412.
11. Zhang WF, Dou ZX, He P, Ju XT, Powlson D and Chadwick D. New technologies reduce greenhouse gas emissions from nitrogenous fertilizer in China. *Proceedings of the National Academy of Sciences of the United States of America*. 2013;110:8375.
12. Shipra Y, Lal M, Naresh RK, Yadav RB, Yadav A, Yadav KG, Kumar R, Sharath Chandra M, and Rajput, P. Effect of organic and inorganic nutrient sources on productivity, grain quality of rice and rice health in North Western IGP: A Review. *International Journal of Current Microbiology and Applied Sciences*. 2019;8(12):2488-2514.
13. Satyanarayana V. Murthy VRK, Prasad V, Boote KJ. Influence of integrated use of farmyard manure and inorganic fertilizers on yield and yield components of irrigated lowland rice. *Journal of Plant Nutrition*. 2002;25(10), 2081-2090.
14. Gill JS, Walia SS. Influence of FYM, brown manuring and nitrogen levels on direct seeded and transplanted rice (*Oryza sativa* L.) A Review. *Research Journal of Agriculture and Environmental Management*. 2014;3(9): 417- 426.
15. Wang Wu L, Xiao Q, Huang Y, Liu K, Wu Y, Li D, Duan Y and Zhang W. Long-term manuring enhances soil gross nitrogen mineralization and ammonium immobilization in subtropical area. *Agriculture, Ecosystems and Environment*. 2023;348:108439.
16. Mohammed MA, Sekar S, Muthukrishnan P. Prospects and Potential of Poultry Manure. *Asian Journal of Plant Sciences*. 2010;9(4):172-182.
17. Arancon QN, Edwards CA. The utilization of vermicompost in Horticulture and Agriculture. In: Edwards CA, Jeyaraaj R, Indira AJ (Eds.) *Vermitechnology in Human welfare*. Rohini Achagam, Coimbatore, Tamil Nadu, India, pp 2009;98-108.
18. Puli M.R, Prasad PRK, Ravindra PB, Jayalakshmi M, Burla SR. (2016). Effect of organic and inorganic sources of nutrients on rice crop (*Oryza sativa* L.). *International Journal of Bio-resource and stress management*. 2016;53:151-159.
19. Shankar T, Maitra S, Sai RM, Mahapatra R. Influence of integrated nutrient management on growth and yield attribution of summer rice (*Oryza sativa* L.). *Crop Research*. 2020;55(1&2):1-5.

20. Jackson ML. Soil chemical analysis, Prentice hall of India, Pvt. Ltd., New Delhi 1973.
21. Sahu YK and Chaubey AK. Effect of integrated nutrient management on nutrient content, uptake and yield of rice crop in inceptisol. 2020;9(3):2531-2541
22. Tiwari H, Singh AK, Pandey SR, Tiwari A. Effect of Integrated nutrient management practices on nutrient content and uptake by rice (*Oryza sativa* L.). Journal of Pharmacognosy and Phytochemistry. 2020;9(6):2131- 2134.
23. Sharma S and Singh J. Split application of potassium improves yield and potassium uptake of rice under deficient soils. Journal of Soil and Water Conservation. 2021;20(2):213-220.
24. Baria, V. K., H. H. Sitapara, A. S. Thounaojam, and P. B. Goswami. 2023. "Influence of Integrated Nitrogen Management on Growth and Yield of Beet Root (*Beta Vulgaris* L.) Var. Crimson Globe". International Journal of Plant & Soil Science 35 (24):123-30. <https://doi.org/10.9734/ijpps/2023/v35i244304>.
25. Kaur, Gurvinder, Ishwar Singh, R.K. Behl, and Amit Dhankar. 2024. "Effect of Different Integrated Nutrient Management Approaches on Growth, Yield Attributes and Yield of Wheat (*Triticum Aestivum* L.) Crop: A Review". Asian Journal of Soil Science and Plant Nutrition 10 (1):457-68. <https://doi.org/10.9734/ajssp/2024/v10i1252>.
26. Rathke GW, Behrens T, Diepenbrock W. Integrated nitrogen management strategies to improve seed yield, oil content and nitrogen efficiency of winter oilseed rape (*Brassica napus* L.): a review. Agriculture, ecosystems & environment. 2006 Nov 1;117(2-3):80-108.